

Complete Semigroups of Binary Relations Defined by Semilattices of the Class $\Sigma_1(X, 10)$

Shota Makharadze¹, Neşet Aydın², Ali Erdoğan³

¹Shota Rustaveli University, Batumi, Georgia

²Çanakkale Onsekiz Mart University, Çanakkale, Turkey

³Hacettepe University, Ankara, Turkey

Email: shota59@mail.ru, neseta@comu.edu.tr, alier@hacettepe.edu.tr

Received 10 January 2015; accepted 28 January 2015; published 4 February 2015

Copyright © 2015 by authors and Scientific Research Publishing Inc.

This work is licensed under the Creative Commons Attribution International License (CC BY).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

Abstract

In this paper we give a full description of idempotent elements of the semigroup $B_X(D)$, which are defined by semilattices of the class $\Sigma_1(X, 10)$. For the case where X is a finite set we derive formulas by means of which we can calculate the numbers of idempotent elements of the respective semigroup.

Keywords

Semilattice, Semigroup, Binary Relation

1. Introduction

Let X be an arbitrary nonempty set, D be an X -semilattice of unions, i.e. such a nonempty set of subsets of the set X that is closed with respect to the set-theoretic operations of unification of elements from D , f be an arbitrary mapping of the set X in the set D . To each such a mapping f we put into correspondence a binary relation α_f on the set X that satisfies the condition

$$\alpha_f = \bigcup_{x \in X} (\{x\} \times f(x))$$

The set of all such α_f ($f: X \rightarrow D$) is denoted by $B_X(D)$. It is easy to prove that $B_X(D)$ is a semigroup with respect to the operation of multiplication of binary relations, which is called a complete semigroup of binary relations defined by an X -semilattice of unions D .

Recall that we denote by \emptyset an empty binary relation or empty subset of the set X . The condition $(x, y) \in \alpha$ will be written in the form $x\alpha y$. Further let $x, y \in X$, $Y \subseteq X$, $\alpha \in B_X(D)$, $T \in D$, $\emptyset \neq D' \subseteq D$, $\bar{D} = \cup D$ and $t \in \bar{D}$. Then by symbols we denoted the following sets:

$$\begin{aligned} y\alpha &= \{x \in X \mid y\alpha x\}, \quad Y\alpha = \bigcup_{y \in Y} y\alpha, \quad 2^X = \{Y \mid Y \subseteq X\}, \quad X^* = 2^X \setminus \{\emptyset\}, \\ V(D, \alpha) &= \{Y\alpha \mid Y \in D\}, \quad D'_T = \{T' \in D' \mid T \subseteq T'\}, \quad \ddot{D}'_T = \{T' \in D' \mid T' \subseteq T\}, \\ D'_t &= \{Z' \in D' \mid t \in Z'\}, \quad l(D', T) = \cup(D' \setminus D'_T). \end{aligned}$$

By symbol $\Lambda(D, D')$ is denoted an exact lower bound of the set D' in the semilattice D .

Definition 1. We say that the complete X -semilattice of unions D is an XI -semilattice of unions if it satisfies the following two conditions:

- $\Lambda(D, D_t) \in D$ for any $t \in \bar{D}$;
- $Z = \bigcup_{t \in Z} \Lambda(D, D_t)$ for any nonempty element Z of the semilattice D .

Definition 2. We say that a nonempty element T is a nonlimiting element of the set D' if $T \setminus l(D', T) \neq \emptyset$ and a nonempty element T is a limiting element of the set D' if $T \setminus l(D', T) = \emptyset$.

Definition 3. Let $\alpha \in B_X(D)$, $T \in V(X^*, \alpha)$, $Y_T^\alpha = \{y \in X \mid y\alpha = T\}$. A representation of a binary relation α of the form $\alpha = \bigcup_{T \in V(X^*, \alpha)} (Y_T^\alpha \times T)$ is called quasinormal.

Note that, if $\alpha = \bigcup_{T \in V(X^*, \alpha)} (Y_T^\alpha \times T)$ is a quasinormal representation of the binary relation α , then the following conditions are true:

- $X = \bigcup_{T \in V(X^*, \alpha)} Y_T^\alpha$;
- $Y_T^\alpha \cap Y_{T'}^\alpha = \emptyset$ for $T, T' \in V(X^*, \alpha)$ and $T \neq T'$.

Let $\sum_n(X, m)$ denote the class of all complete X -semilattices of unions where every element is isomorphic to a fixed semilattice D .

The following Theorems are well know (see [1] and [3]).

Theorem 4. Let X be a finite set; δ and q be respectively the number of basic sources and the number of all automorphisms of the semilattice D . If $|X| = n \geq \delta$ and $|\sum_n(X, m)| = s$, then

$$s = \frac{1}{q} \cdot \sum_{p=\delta}^m \left(\sum_{i=1}^{p+1} \frac{(-1)^{p+i+1} \cdot C_{m-\delta}^{p-\delta} \cdot C_p^\delta \cdot (\delta!) \cdot ((p-\delta)!) \cdot i^n}{(i-1)! \cdot (p-i+1)!} \right)$$

where $C_j^k = \frac{j!}{(k!) \cdot (j-k)!}$ (see Theorem 11.5.1 [1]).

Theorem 5. Let D be a complete X -semilattice of unions. The semigroup $B_X(D)$ possesses right unit iff D is an XI -semilattice of unions (see Theorem 6.1.3 [1]).

Theorem 6. Let X be a finite set and $D(\alpha)$ be the set of all those elements T of the semilattice $Q = V(D, \alpha) \setminus \{\emptyset\}$ which are nonlimiting elements of the set \ddot{Q}_T . A binary relation α having a quasinormal representation $\alpha = \bigcup_{T \in V(D, \alpha)} (Y_T^\alpha \times T)$ is an idempotent element of this semigroup iff

- $V(D, \alpha)$ is complete XI -semilattice of unions;
- $\bigcup_{T' \in \ddot{D}(\alpha)_T} Y_{T'}^\alpha \supseteq T$ for any $T \in D(\alpha)$;
- $Y_T^\alpha \cap T \neq \emptyset$ for any nonlimiting element of the set $\ddot{D}(\alpha)_T$ (see Theorem 6.3.9 [1]).

Theorem 7. Let D , $\Sigma(D)$, $E_X^{(r)}(D')$ and I denote respectively the complete X -semilattice of unions, the set of all XI -subsemilattices of the semilattice D , the set of all right units of the semigroup $B_X(D')$ and the set of all idempotents of the semigroup $B_X(D)$. Then for the sets $E_X^{(r)}(D')$ and I the following statements are true:

- if $\emptyset \in D$ and $\Sigma_\emptyset(D) = \{D' \in \Sigma(D) \mid \emptyset \in D'\}$ then
 - $E_X^{(r)}(D') \cap E_X^{(r)}(D'') = \emptyset$ for any elements D' and D'' of the set $\Sigma_\emptyset(D)$ that satisfy the condition $D' \neq D''$;
 - $I = \bigcup_{D' \in \Sigma_\emptyset(D)} E_X^{(r)}(D')$
- the equality $|I| = \sum_{D' \in \Sigma_\emptyset(D)} |E_X^{(r)}(D')|$ is fulfilled for the finite set X .

2) if $\emptyset \notin D$, then
 a) $E_X^{(r)}(D') \cap E_X^{(r)}(D'') = \emptyset$ for any elements D' and D'' of the set $\Sigma(D)$ that satisfy the condition $D' \neq D''$;

b) $I = \bigcup_{D' \in \Sigma(D)} E_X^{(r)}(D')$

c) the equality $|I| = \sum_{D' \in \Sigma(D)} |E_X^{(r)}(D')|$ is fulfilled for the finite set X (see Theorem 6.2.3 [1]).

Corollary 1. Let $Y = \{y_1, y_2, \dots, y_k\}$ and $D_j = \{T_1, T_2, \dots, T_j\}$ be some sets, where $k \geq 1$ and $j \geq 1$. Then the number $s(k, j)$ of all possible mappings of the set Y into any such subset D'_j of the set D_j that $T_j \in D'_j$ can be calculated by the formula $s(k, j) = j^k - (j-1)^k$ (see Corollary 1.18.1 [1]).

2. Idempotent Elements of the Semigroups $B_X(D)$ Defined by Semilattices of the Class $\Sigma_1(X, 10)$

Let X and $\Sigma_1(X, 10)$ be respectively an arbitrary nonempty set and a class X -semilattices of unions, where each element is isomorphic to some X -semilattice of unions $D = \{Z_9, Z_8, Z_7, Z_6, Z_5, Z_4, Z_3, Z_2, Z_1, \check{D}\}$ that satisfies the conditions:

$$\begin{aligned}
 & Z_9 \subset Z_4 \subset Z_1 \subset \check{D}, Z_9 \subset Z_5 \subset Z_1 \subset \check{D}, \\
 & Z_9 \subset Z_6 \subset Z_1 \subset \check{D}, Z_9 \subset Z_6 \subset Z_2 \subset \check{D}, \\
 & Z_9 \subset Z_6 \subset Z_3 \subset \check{D}, Z_9 \subset Z_7 \subset Z_3 \subset \check{D}, \\
 & Z_9 \subset Z_8 \subset Z_3 \subset \check{D}, Z_1 \setminus Z_2 \neq \emptyset, Z_2 \setminus Z_1 \neq \emptyset, \\
 & Z_1 \setminus Z_3 \neq \emptyset, Z_3 \setminus Z_1 \neq \emptyset, Z_2 \setminus Z_3 \neq \emptyset, \\
 & Z_3 \setminus Z_2 \neq \emptyset, Z_4 \setminus Z_5 \neq \emptyset, Z_5 \setminus Z_4 \neq \emptyset, \\
 & Z_4 \setminus Z_6 \neq \emptyset, Z_6 \setminus Z_4 \neq \emptyset, Z_4 \setminus Z_7 \neq \emptyset, \\
 & Z_7 \setminus Z_4 \neq \emptyset, Z_4 \setminus Z_8 \neq \emptyset, Z_8 \setminus Z_4 \neq \emptyset, \\
 & Z_5 \setminus Z_6 \neq \emptyset, Z_6 \setminus Z_5 \neq \emptyset, Z_5 \setminus Z_7 \neq \emptyset, \\
 & Z_7 \setminus Z_5 \neq \emptyset, Z_5 \setminus Z_8 \neq \emptyset, Z_8 \setminus Z_5 \neq \emptyset, \\
 & Z_6 \setminus Z_7 \neq \emptyset, Z_7 \setminus Z_6 \neq \emptyset, Z_6 \setminus Z_8 \neq \emptyset, \\
 & Z_8 \setminus Z_6 \neq \emptyset, Z_7 \setminus Z_8 \neq \emptyset, Z_8 \setminus Z_7 \neq \emptyset, \\
 & Z_1 \cup Z_2 = Z_1 \cup Z_3 = Z_2 \cup Z_3 = Z_4 \cup Z_2 \\
 & \quad = Z_4 \cup Z_3 = Z_4 \cup Z_7 = Z_4 \cup Z_8 = Z_5 \cup Z_2 \\
 & \quad = Z_5 \cup Z_3 = Z_5 \cup Z_7 = Z_5 \cup Z_8 = Z_7 \cup Z_1 \\
 & \quad = Z_7 \cup Z_2 = Z_8 \cup Z_1 = Z_8 \cup Z_2 = \check{D}, \\
 & Z_4 \cup Z_5 = Z_4 \cup Z_6 = Z_5 \cup Z_6 = Z_1, \\
 & Z_6 \cup Z_7 = Z_6 \cup Z_8 = Z_7 \cup Z_8 = Z_3.
 \end{aligned} \tag{1}$$

An X -semilattice that satisfies conditions (1) is shown in Figure 1.

Let $C(D) = \{P_0, P_1, P_2, P_3, P_4, P_5, P_6, P_7, P_8, P_9\}$ be a family of sets, where $P_0, P_1, P_2, P_3, P_4, P_5, P_6, P_7, P_8, P_9$




Figure 1. Diagram of D .

are pairwise disjoint subsets of the set X and $\varphi = \begin{pmatrix} \tilde{D} & Z_1 & Z_2 & Z_3 & Z_4 & Z_5 & Z_6 & Z_7 & Z_8 & Z_9 \\ P_0 & P_1 & P_2 & P_3 & P_4 & P_5 & P_6 & P_7 & P_8 & P_9 \end{pmatrix}$ be a mapping of the semilattice D onto the family sets $C(D)$. Then for the formal equalities of the semilattice D we have a form:

$$\begin{aligned} \tilde{D} &= P_0 \cup P_1 \cup P_2 \cup P_3 \cup P_4 \cup P_5 \cup P_6 \cup P_7 \cup P_8 \cup P_9, \\ Z_1 &= P_0 \cup P_2 \cup P_3 \cup P_4 \cup P_5 \cup P_6 \cup P_7 \cup P_8 \cup P_9, \\ Z_2 &= P_0 \cup P_1 \cup P_3 \cup P_4 \cup P_5 \cup P_6 \cup P_7 \cup P_8 \cup P_9, \\ Z_3 &= P_0 \cup P_1 \cup P_2 \cup P_4 \cup P_5 \cup P_6 \cup P_7 \cup P_8 \cup P_9, \\ Z_4 &= P_0 \cup P_2 \cup P_3 \cup P_5 \cup P_6 \cup P_7 \cup P_8 \cup P_9, \\ Z_5 &= P_0 \cup P_2 \cup P_3 \cup P_4 \cup P_6 \cup P_7 \cup P_8 \cup P_9, \\ Z_6 &= P_0 \cup P_4 \cup P_5 \cup P_7 \cup P_8 \cup P_9, \\ Z_7 &= P_0 \cup P_1 \cup P_2 \cup P_4 \cup P_5 \cup P_6 \cup P_8 \cup P_9, \\ Z_8 &= P_0 \cup P_1 \cup P_2 \cup P_4 \cup P_5 \cup P_6 \cup P_7 \cup P_9, \\ Z_9 &= P_0. \end{aligned} \quad (2)$$

Here the elements $P_1, P_2, P_3, P_4, P_5, P_6, P_7, P_8$ are basis sources, the elements P_0, P_6, P_9 are sources of completeness of the semilattice D . Therefore $|X| \geq 7$ and $\delta = 7$ (see [2]).

Lemma 1. Let $D \in \Sigma_1(X, 10)$, $|\Sigma_1(X, 10)| = s$ and $|X| \geq \delta \geq 7$. If X is a finite set, then

$$s = \frac{1}{8} \left((-1) \times 4^n + 7 \times 5^n - 21 \times 6^n + 35 \times 7^n - 35 \times 8^n + 21 \times 9^n + 11^n \right).$$

Proof. In this case we have: $m = 10$, $\delta = 7$. Notice that an X -semilattice given in Figure 1 has eight automorphisms. By Theorem 1.1 it follows that

$$s = \frac{1}{8} \cdot \sum_{p=7}^{10} \left(\sum_{i=1}^{p+1} \left(\frac{(-1)^{p+i+1} \cdot C_3^{p-7} \cdot C_p^7 \cdot (7!) \cdot ((p-7)!) \cdot i^n}{(i-1)! \cdot (p-i+1)!} \right) \right),$$

where $C_j^k = \frac{j!}{k! \cdot (j-k)!}$ and that

$$s = \frac{1}{8} \left((-1) \times 4^n + 7 \times 5^n - 21 \times 6^n + 35 \times 7^n - 35 \times 8^n + 21 \times 9^n + 11^n \right).$$

Example 8. Let $n = 7, 8, 9, 10$ Then:

$$|B_X(D)| = 10^7, 10^8, 10^9, 10^{10}.$$

Lemma 2. Let $D \in \Sigma_1(X, 10)$. Then the following sets are all proper subsemilattices of the semilattice $D = \{Z_9, Z_8, Z_7, Z_6, Z_5, Z_4, Z_3, Z_2, Z_1, \tilde{D}\}$:

- 1) $\{Z_9\}, \{Z_8\}, \{Z_7\}, \{Z_6\}, \{Z_5\}, \{Z_4\}, \{Z_3\}, \{Z_2\}, \{Z_1\}, \{\tilde{D}\}$
(see diagram 1 of the Figure 2);
- 2) $\{Z_9, Z_8\}, \{Z_9, Z_7\}, \{Z_9, Z_6\}, \{Z_9, Z_5\}, \{Z_9, Z_4\}, \{Z_9, Z_3\}, \{Z_9, Z_2\}, \{Z_9, Z_1\}, \{Z_9, \tilde{D}\}, \{Z_8, Z_3\},$
 $\{Z_8, \tilde{D}\}, \{Z_7, Z_3\}, \{Z_7, \tilde{D}\}, \{Z_6, Z_3\}, \{Z_6, Z_2\}, \{Z_6, Z_1\}, \{Z_6, \tilde{D}\}, \{Z_5, Z_1\}, \{Z_5, \tilde{D}\}, \{Z_4, Z_1\},$
 $\{Z_4, \tilde{D}\}, \{Z_3, \tilde{D}\}, \{Z_2, \tilde{D}\}, \{Z_1, \tilde{D}\}$
(see diagram 2 of the Figure 2);
- 3) $\{Z_9, Z_8, Z_3\}, \{Z_9, Z_8, \tilde{D}\}, \{Z_9, Z_7, Z_3\}, \{Z_9, Z_7, \tilde{D}\}, \{Z_9, Z_6, Z_3\}, \{Z_9, Z_6, Z_2\}, \{Z_9, Z_6, Z_1\},$
 $\{Z_9, Z_6, \tilde{D}\}, \{Z_9, Z_5, Z_1\}, \{Z_9, Z_5, \tilde{D}\}, \{Z_9, Z_4, Z_1\}, \{Z_9, Z_4, \tilde{D}\}, \{Z_9, Z_3, \tilde{D}\}, \{Z_9, Z_2, \tilde{D}\},$

- $\{Z_9, Z_1, \bar{D}\}, \{Z_8, Z_3, \bar{D}\}, \{Z_7, Z_3, \bar{D}\}, \{Z_6, Z_3, \bar{D}\}, \{Z_6, Z_2, \bar{D}\}, \{Z_6, Z_1, \bar{D}\}, \{Z_5, Z_1, \bar{D}\}, \{Z_4, Z_1, \bar{D}\}$
(see diagram 3 of the **Figure 2**);
- 4) $\{Z_9, Z_4, Z_1, \bar{D}\}, \{Z_9, Z_5, Z_1, \bar{D}\}, \{Z_9, Z_6, Z_1, \bar{D}\}, \{Z_9, Z_6, Z_2, \bar{D}\}, \{Z_9, Z_6, Z_3, \bar{D}\},$
 $\{Z_9, Z_7, Z_3, \bar{D}\}, \{Z_9, Z_8, Z_3, \bar{D}\}$
(see diagram 4 of the **Figure 2**);
- 5) $\{Z_9, Z_5, Z_4, Z_1\}, \{Z_9, Z_6, Z_4, Z_1\}, \{Z_9, Z_6, Z_5, Z_1\}, \{Z_9, Z_7, Z_6, Z_3\}, \{Z_9, Z_8, Z_6, Z_3\}, \{Z_9, Z_8, Z_7, Z_3\},$
 $\{Z_9, Z_8, Z_4, \bar{D}\}, \{Z_9, Z_8, Z_5, \bar{D}\}, \{Z_9, Z_7, Z_2, \bar{D}\}, \{Z_9, Z_7, Z_4, \bar{D}\}, \{Z_9, Z_7, Z_5, \bar{D}\}, \{Z_9, Z_8, Z_1, \bar{D}\},$
 $\{Z_9, Z_8, Z_2, \bar{D}\}, \{Z_9, Z_4, Z_2, \bar{D}\}, \{Z_9, Z_4, Z_3, \bar{D}\}, \{Z_9, Z_5, Z_2, \bar{D}\}, \{Z_9, Z_5, Z_3, \bar{D}\}, \{Z_9, Z_7, Z_1, \bar{D}\},$
 $\{Z_9, Z_2, Z_1, \bar{D}\}, \{Z_9, Z_3, Z_1, \bar{D}\}, \{Z_9, Z_3, Z_2, \bar{D}\}, \{Z_6, Z_2, Z_1, \bar{D}\}, \{Z_6, Z_3, Z_1, \bar{D}\}, \{Z_6, Z_3, Z_2, \bar{D}\};$
(see diagram 5 of the **Figure 2**);
- 6) $\{Z_9, Z_5, Z_4, Z_1, \bar{D}\}, \{Z_9, Z_6, Z_4, Z_1, \bar{D}\}, \{Z_9, Z_6, Z_5, Z_1, \bar{D}\}, \{Z_9, Z_7, Z_6, Z_3, \bar{D}\},$
 $\{Z_9, Z_8, Z_6, Z_3, \bar{D}\}, \{Z_9, Z_8, Z_7, Z_3, \bar{D}\}$
(see diagram 6 of the **Figure 2**);
- 7) $\{Z_9, Z_6, Z_2, Z_1, \bar{D}\}, \{Z_9, Z_6, Z_3, Z_1, \bar{D}\}, \{Z_9, Z_6, Z_3, Z_2, \bar{D}\}$
(see diagram 7 of the **Figure 2**);
- 8) $\{Z_9, Z_8, Z_6, Z_3, Z_2, \bar{D}\}, \{Z_9, Z_8, Z_6, Z_3, Z_1, \bar{D}\}, \{Z_9, Z_7, Z_6, Z_3, Z_2, \bar{D}\}, \{Z_9, Z_7, Z_6, Z_3, Z_1, \bar{D}\},$
 $\{Z_9, Z_6, Z_5, Z_3, Z_1, \bar{D}\}, \{Z_9, Z_6, Z_5, Z_2, Z_1, \bar{D}\}, \{Z_9, Z_6, Z_4, Z_3, Z_1, \bar{D}\}, \{Z_9, Z_6, Z_4, Z_2, Z_1, \bar{D}\}$
(see diagram 8 of the **Figure 2**);
- 9) $\{Z_8, Z_7, Z_3\}, \{Z_8, Z_6, Z_3\}, \{Z_8, Z_6, \bar{D}\}, \{Z_8, Z_5, \bar{D}\}, \{Z_8, Z_4, \bar{D}\}, \{Z_8, Z_2, \bar{D}\}, \{Z_8, Z_1, \bar{D}\}, \{Z_7, Z_6, Z_3\},$
 $\{Z_7, Z_5, \bar{D}\}, \{Z_7, Z_4, \bar{D}\}, \{Z_7, Z_2, \bar{D}\}, \{Z_7, Z_1, \bar{D}\}, \{Z_6, Z_5, Z_1\}, \{Z_6, Z_4, Z_1\}, \{Z_5, Z_4, Z_1\}, \{Z_5, Z_3, \bar{D}\},$
 $\{Z_5, Z_2, \bar{D}\}, \{Z_4, Z_3, \bar{D}\}, \{Z_4, Z_2, \bar{D}\}, \{Z_3, Z_2, \bar{D}\}, \{Z_3, Z_1, \bar{D}\}, \{Z_2, Z_1, \bar{D}\}$
(see diagram 9 of the **Figure 2**);
- 10) $\{Z_8, Z_6, Z_3, \bar{D}\}, \{Z_8, Z_7, Z_3, \bar{D}\}, \{Z_7, Z_6, Z_3, \bar{D}\}, \{Z_5, Z_4, Z_1, \bar{D}\}, \{Z_6, Z_4, Z_1, \bar{D}\}, \{Z_6, Z_5, Z_1, \bar{D}\}$
(see diagram 10 of the **Figure 3**);
- 11) $\{Z_6, Z_5, Z_3, Z_1, \bar{D}\}, \{Z_6, Z_5, Z_2, Z_1, \bar{D}\}, \{Z_6, Z_4, Z_3, Z_1, \bar{D}\}, \{Z_6, Z_4, Z_2, Z_1, \bar{D}\},$
 $\{Z_7, Z_6, Z_3, Z_2, \bar{D}\}, \{Z_7, Z_6, Z_3, Z_1, \bar{D}\}, \{Z_8, Z_6, Z_3, Z_2, \bar{D}\}, \{Z_8, Z_6, Z_3, Z_1, \bar{D}\}$
(see diagram 11 of the **Figure 2**);
- 12) $\{Z_6, Z_5, Z_4, Z_1\}, \{Z_8, Z_7, Z_6, Z_3\}, \{Z_8, Z_2, Z_1, \bar{D}\}, \{Z_3, Z_2, Z_1, \bar{D}\}, \{Z_4, Z_3, Z_2, \bar{D}\}, \{Z_5, Z_3, Z_2, \bar{D}\},$
 $\{Z_7, Z_2, Z_1, \bar{D}\}, \{Z_7, Z_4, Z_2, \bar{D}\}, \{Z_7, Z_5, Z_2, \bar{D}\}, \{Z_8, Z_4, Z_2, \bar{D}\}, \{Z_8, Z_5, Z_2, \bar{D}\}$
(see diagram 12 of the **Figure 2**);
- 13) $\{Z_7, Z_5, Z_3, \bar{D}\}, \{Z_4, Z_2, Z_1, \bar{D}\}, \{Z_8, Z_3, Z_1, \bar{D}\}, \{Z_8, Z_3, Z_2, \bar{D}\}, \{Z_8, Z_4, Z_1, \bar{D}\}, \{Z_4, Z_3, Z_1, \bar{D}\},$
 $\{Z_5, Z_2, Z_1, \bar{D}\}, \{Z_5, Z_3, Z_1, \bar{D}\}, \{Z_7, Z_3, Z_1, \bar{D}\}, \{Z_7, Z_3, Z_2, \bar{D}\}, \{Z_7, Z_4, Z_1, \bar{D}\}, \{Z_7, Z_4, Z_3, \bar{D}\},$
 $\{Z_7, Z_5, Z_1, \bar{D}\}, \{Z_8, Z_4, Z_3, \bar{D}\}, \{Z_8, Z_5, Z_1, \bar{D}\}, \{Z_8, Z_5, Z_3, \bar{D}\}$
(see diagram 13 of the **Figure 2**);
- 14) $\{Z_9, Z_6, Z_5, Z_4, Z_1\}, \{Z_9, Z_8, Z_7, Z_6, Z_3\}, \{Z_6, Z_3, Z_2, Z_1, \bar{D}\}, \{Z_9, Z_3, Z_2, Z_1, \bar{D}\}, \{Z_9, Z_4, Z_3, Z_2, \bar{D}\},$

- $\{Z_9, Z_5, Z_3, Z_2, \bar{D}\}, \{Z_9, Z_7, Z_2, Z_1, \bar{D}\}, \{Z_9, Z_7, Z_4, Z_2, \bar{D}\}, \{Z_9, Z_7, Z_4, Z_3, \bar{D}\}, \{Z_9, Z_7, Z_5, Z_2, \bar{D}\},$
 $\{Z_9, Z_8, Z_2, Z_1, \bar{D}\}, \{Z_9, Z_8, Z_4, Z_2, \bar{D}\}, \{Z_9, Z_8, Z_5, Z_2, \bar{D}\}$
 (see diagram 14 of the **Figure 2**);
- 15) $\{Z_9, Z_4, Z_2, Z_1, \bar{D}\}, \{Z_9, Z_4, Z_3, Z_1, \bar{D}\}, \{Z_9, Z_5, Z_2, Z_1, \bar{D}\}, \{Z_9, Z_5, Z_3, Z_1, \bar{D}\}, \{Z_9, Z_7, Z_3, Z_1, \bar{D}\},$
 $\{Z_9, Z_7, Z_3, Z_2, \bar{D}\}, \{Z_9, Z_7, Z_4, Z_1, \bar{D}\}, \{Z_9, Z_7, Z_5, Z_1, \bar{D}\}, \{Z_9, Z_7, Z_5, Z_3, \bar{D}\}, \{Z_9, Z_8, Z_3, Z_1, \bar{D}\},$
 $\{Z_9, Z_8, Z_3, Z_2, \bar{D}\}, \{Z_9, Z_8, Z_4, Z_1, \bar{D}\}, \{Z_9, Z_8, Z_4, Z_3, \bar{D}\}, \{Z_9, Z_8, Z_5, Z_1, \bar{D}\}, \{Z_9, Z_8, Z_5, Z_3, \bar{D}\}$
 (see diagram 15 of the **Figure 2**);
- 16) $\{Z_5, Z_4, Z_3, Z_1, \bar{D}\}, \{Z_5, Z_4, Z_2, Z_1, \bar{D}\}, \{Z_7, Z_5, Z_4, Z_1, \bar{D}\}, \{Z_8, Z_5, Z_4, Z_1, \bar{D}\}, \{Z_8, Z_7, Z_3, Z_2, \bar{D}\},$
 $\{Z_8, Z_7, Z_3, Z_1, \bar{D}\}, \{Z_8, Z_7, Z_5, Z_3, \bar{D}\}, \{Z_8, Z_7, Z_4, Z_3, \bar{D}\}$
 (see diagram 16 of the **Figure 2**);
- 17) $\{Z_5, Z_3, Z_2, Z_1, \bar{D}\}, \{Z_4, Z_3, Z_2, Z_1, \bar{D}\}, \{Z_7, Z_4, Z_2, Z_1, \bar{D}\}, \{Z_7, Z_3, Z_2, Z_1, \bar{D}\}, \{Z_7, Z_5, Z_3, Z_2, \bar{D}\},$
 $\{Z_7, Z_5, Z_2, Z_1, \bar{D}\}, \{Z_7, Z_4, Z_3, Z_2, \bar{D}\}, \{Z_8, Z_3, Z_2, Z_1, \bar{D}\}, \{Z_8, Z_5, Z_3, Z_2, \bar{D}\}, \{Z_8, Z_5, Z_2, Z_1, \bar{D}\},$
 $\{Z_8, Z_4, Z_3, Z_2, \bar{D}\}, \{Z_8, Z_4, Z_2, Z_1, \bar{D}\}$
 (see diagram 17 of the **Figure 2**);
- 18) $\{Z_7, Z_4, Z_3, Z_1, \bar{D}\}, \{Z_7, Z_5, Z_3, Z_1, \bar{D}\}, \{Z_8, Z_5, Z_3, Z_1, \bar{D}\}, \{Z_8, Z_4, Z_3, Z_1, \bar{D}\}$
 (see diagram 18 of the **Figure 2**);
- 19) $\{Z_6, Z_5, Z_4, Z_1, \bar{D}\}, \{Z_8, Z_7, Z_6, Z_3, \bar{D}\}.$
 (see diagram 19 of the **Figure 2**);
- 20) $\{Z_8, Z_7, Z_5, Z_3, Z_2, \bar{D}\}, \{Z_8, Z_7, Z_4, Z_3, Z_2, \bar{D}\}, \{Z_8, Z_5, Z_4, Z_2, Z_1, \bar{D}\}, \{Z_7, Z_5, Z_4, Z_2, Z_1, \bar{D}\},$
 $\{Z_8, Z_7, Z_3, Z_2, Z_1, \bar{D}\}, \{Z_5, Z_4, Z_3, Z_2, Z_1, \bar{D}\}$
 (see diagram 20 of the **Figure 2**);
- 21) $\{Z_6, Z_5, Z_4, Z_2, Z_1, \bar{D}\}, \{Z_8, Z_7, Z_6, Z_3, Z_2, \bar{D}\}, \{Z_8, Z_7, Z_6, Z_3, Z_1, \bar{D}\}, \{Z_6, Z_5, Z_4, Z_3, Z_1, \bar{D}\}$
 (see diagram 21 of the **Figure 2**);
- 22) $\{Z_9, Z_5, Z_4, Z_2, Z_1, \bar{D}\}, \{Z_9, Z_8, Z_7, Z_4, Z_3, \bar{D}\}, \{Z_9, Z_8, Z_7, Z_3, Z_1, \bar{D}\}, \{Z_9, Z_8, Z_5, Z_4, Z_1, \bar{D}\},$
 $\{Z_9, Z_7, Z_5, Z_4, Z_1, \bar{D}\}, \{Z_9, Z_5, Z_4, Z_3, Z_1, \bar{D}\}, \{Z_9, Z_8, Z_7, Z_3, Z_2, \bar{D}\}, \{Z_9, Z_8, Z_7, Z_5, Z_3, \bar{D}\}$
 (see diagram 22 of the **Figure 2**);
- 23) $\{Z_9, Z_8, Z_7, Z_6, Z_3, \bar{D}\}, \{Z_9, Z_6, Z_5, Z_4, Z_1, \bar{D}\}$
 (see diagram 23 of the **Figure 2**);
- 24) $\{Z_9, Z_8, Z_5, Z_3, Z_2, \bar{D}\}, \{Z_9, Z_8, Z_5, Z_2, Z_1, \bar{D}\}, \{Z_9, Z_8, Z_4, Z_3, Z_2, \bar{D}\}, \{Z_9, Z_8, Z_4, Z_2, Z_1, \bar{D}\},$
 $\{Z_9, Z_8, Z_3, Z_2, Z_1, \bar{D}\}, \{Z_9, Z_7, Z_5, Z_3, Z_2, \bar{D}\}, \{Z_9, Z_7, Z_5, Z_2, Z_1, \bar{D}\}, \{Z_9, Z_7, Z_4, Z_3, Z_2, \bar{D}\},$
 $\{Z_9, Z_7, Z_4, Z_2, Z_1, \bar{D}\}, \{Z_9, Z_7, Z_3, Z_2, Z_1, \bar{D}\}, \{Z_9, Z_5, Z_3, Z_2, Z_1, \bar{D}\}, \{Z_9, Z_4, Z_3, Z_2, Z_1, \bar{D}\}$
 (see diagram 24 of the **Figure 2**);
- 25) $\{Z_9, Z_8, Z_5, Z_3, Z_1, \bar{D}\}, \{Z_9, Z_8, Z_4, Z_3, Z_1, \bar{D}\}, \{Z_9, Z_7, Z_5, Z_3, Z_1, \bar{D}\}, \{Z_9, Z_7, Z_4, Z_3, Z_1, \bar{D}\}$
 (see diagram 25 of the **Figure 2**);
- 26) $\{Z_9, Z_6, Z_3, Z_2, Z_1, \bar{D}\}$
 (see diagram 26 of the **Figure 2**);

- 27) $\{Z_8, Z_7, Z_5, Z_3, Z_1, \bar{D}\}, \{Z_8, Z_7, Z_4, Z_3, Z_1, \bar{D}\}, \{Z_8, Z_5, Z_4, Z_3, Z_1, \bar{D}\}, \{Z_7, Z_5, Z_4, Z_3, Z_1, \bar{D}\}$
(see diagram 27 of the **Figure 2**);
- 28) $\{Z_8, Z_6, Z_5, Z_3, Z_1, \bar{D}\}, \{Z_8, Z_6, Z_4, Z_3, Z_1, \bar{D}\}, \{Z_8, Z_5, Z_3, Z_2, Z_1, \bar{D}\}, \{Z_7, Z_6, Z_5, Z_3, Z_1, \bar{D}\},$
 $\{Z_7, Z_6, Z_4, Z_3, Z_1, \bar{D}\}$
(see diagram 28 of the **Figure 2**);
- 29) $\{Z_8, Z_6, Z_3, Z_2, Z_1, \bar{D}\}, \{Z_7, Z_6, Z_3, Z_2, Z_1, \bar{D}\}, \{Z_6, Z_5, Z_3, Z_2, Z_1, \bar{D}\}, \{Z_6, Z_4, Z_3, Z_2, Z_1, \bar{D}\}$
(see diagram 29 of the **Figure 2**);
- 30) $\{Z_8, Z_4, Z_3, Z_2, Z_1, \bar{D}\}, \{Z_7, Z_5, Z_3, Z_2, Z_1, \bar{D}\}, \{Z_7, Z_4, Z_3, Z_2, Z_1, \bar{D}\}$
(see diagram 30 of the **Figure 2**);
- 31) $\{Z_8, Z_7, Z_5, Z_4, Z_3, Z_1, \bar{D}\}$
(see diagram 31 of the **Figure 2**);
- 32) $\{Z_6, Z_5, Z_4, Z_3, Z_2, Z_1, \bar{D}\}, \{Z_8, Z_7, Z_6, Z_3, Z_2, Z_1, \bar{D}\}$
(see diagram 32 of the **Figure 2**);
- 33) $\{Z_7, Z_5, Z_4, Z_3, Z_2, Z_1, \bar{D}\}, \{Z_8, Z_5, Z_4, Z_3, Z_2, Z_1, \bar{D}\}, \{Z_8, Z_7, Z_4, Z_3, Z_2, Z_1, \bar{D}\},$
 $\{Z_8, Z_7, Z_5, Z_3, Z_2, Z_1, \bar{D}\}$
(see diagram 33 of the **Figure 2**);
- 34) $\{Z_7, Z_6, Z_5, Z_4, Z_3, Z_1, \bar{D}\}, \{Z_8, Z_6, Z_5, Z_4, Z_3, Z_1, \bar{D}\}, \{Z_8, Z_7, Z_6, Z_4, Z_3, Z_1, \bar{D}\},$
 $\{Z_8, Z_7, Z_6, Z_5, Z_3, Z_1, \bar{D}\}$
(see diagram 34 of the **Figure 2**);
- 35) $\{Z_9, Z_6, Z_4, Z_3, Z_2, Z_1, \bar{D}\}, \{Z_9, Z_6, Z_5, Z_3, Z_2, Z_1, \bar{D}\}, \{Z_9, Z_7, Z_6, Z_3, Z_2, Z_1, \bar{D}\},$
 $\{Z_9, Z_8, Z_6, Z_3, Z_2, Z_1, \bar{D}\}, \{Z_9, Z_8, Z_6, Z_3, Z_2, Z_1, \bar{D}\}$
(see diagram 35 of the **Figure 2**);
- 36) $\{Z_9, Z_6, Z_5, Z_4, Z_3, Z_1, \bar{D}\}, \{Z_9, Z_8, Z_7, Z_6, Z_3, Z_1, \bar{D}\}, \{Z_9, Z_8, Z_7, Z_6, Z_3, Z_1, \bar{D}\}$
(see diagram 36 of the **Figure 2**);
- 37) $\{Z_9, Z_7, Z_4, Z_3, Z_2, Z_1, \bar{D}\}, \{Z_9, Z_7, Z_5, Z_3, Z_2, Z_1, \bar{D}\}, \{Z_9, Z_8, Z_4, Z_3, Z_2, Z_1, \bar{D}\},$
 $\{Z_9, Z_8, Z_5, Z_3, Z_2, Z_1, \bar{D}\}$
(see diagram 37 of the **Figure 2**);
- 38) $\{Z_9, Z_7, Z_5, Z_4, Z_3, Z_1, \bar{D}\}, \{Z_9, Z_8, Z_5, Z_4, Z_3, Z_1, \bar{D}\}, \{Z_9, Z_8, Z_7, Z_4, Z_3, Z_1, \bar{D}\}$
(see diagram 38 of the **Figure 2**);
- 39) $\{Z_9, Z_7, Z_6, Z_5, Z_3, Z_1, \bar{D}\}$
(see diagram 39 of the **Figure 2**);
- 40) $\{Z_9, Z_7, Z_5, Z_4, Z_2, Z_1, \bar{D}\}, \{Z_9, Z_8, Z_5, Z_4, Z_2, Z_1, \bar{D}\}, \{Z_9, Z_8, Z_7, Z_4, Z_3, Z_2, \bar{D}\},$
 $\{Z_9, Z_8, Z_7, Z_5, Z_3, Z_2, \bar{D}\}, \{Z_9, Z_5, Z_4, Z_3, Z_2, Z_1, \bar{D}\}$
(see diagram 40 of the **Figure 2**);
- 41) $\{Z_9, Z_6, Z_5, Z_4, Z_2, Z_1, \bar{D}\}, \{Z_9, Z_8, Z_7, Z_6, Z_3, Z_2, \bar{D}\}$
(see diagram 41 of the **Figure 2**);

- 42) $\{Z_7, Z_6, Z_5, Z_4, Z_3, Z_2, Z_1, \bar{D}\}$
(see diagram 42 of the **Figure 2**);
- 43) $\{Z_8, Z_6, Z_5, Z_4, Z_3, Z_2, Z_1, \bar{D}\}, \{Z_8, Z_7, Z_6, Z_4, Z_3, Z_2, Z_1, \bar{D}\}, \{Z_8, Z_7, Z_6, Z_5, Z_3, Z_2, Z_1, \bar{D}\}$
(see diagram 43 of the **Figure 2**);
- 44) $\{Z_8, Z_7, Z_5, Z_4, Z_3, Z_2, Z_1, \bar{D}\}$
(see diagram 44 of the **Figure 2**);
- 45) $\{Z_9, Z_8, Z_7, Z_5, Z_4, Z_3, Z_1, \bar{D}\}$
(see diagram 45 of the **Figure 2**);
- 46) $\{Z_9, Z_6, Z_5, Z_4, Z_3, Z_2, Z_1, \bar{D}\}, \{Z_9, Z_8, Z_7, Z_6, Z_3, Z_2, Z_1, \bar{D}\}$
(see diagram 46 of the **Figure 2**);
- 47) $\{Z_9, Z_7, Z_5, Z_4, Z_3, Z_2, Z_1, \bar{D}\}, \{Z_9, Z_8, Z_7, Z_5, Z_3, Z_2, Z_1, \bar{D}\}, \{Z_9, Z_7, Z_5, Z_4, Z_3, Z_2, Z_1, \bar{D}\},$
 $\{Z_9, Z_8, Z_7, Z_4, Z_3, Z_2, Z_1, \bar{D}\}$
(see diagram 47 of the **Figure 2**);
- 48) $\{Z_9, Z_7, Z_6, Z_5, Z_4, Z_3, Z_1, \bar{D}\}, \{Z_9, Z_8, Z_6, Z_5, Z_4, Z_3, Z_1, \bar{D}\}, \{Z_9, Z_8, Z_7, Z_6, Z_4, Z_3, Z_1, \bar{D}\},$
 $\{Z_9, Z_8, Z_7, Z_6, Z_5, Z_3, Z_1, \bar{D}\}$
(see diagram 48 of the **Figure 2**);




Figure 2. Diagram of all subsemilattices of D .

- 49) $\{Z_8, Z_7, Z_6, Z_5, Z_4, Z_3, Z_2, Z_1, \bar{D}\}$
(see diagram 49 of the **Figure 2**);
- 50) $\{Z_9, Z_7, Z_6, Z_5, Z_4, Z_3, Z_2, Z_1, \bar{D}\}, \{Z_9, Z_8, Z_6, Z_5, Z_4, Z_3, Z_2, Z_1, \bar{D}\}, \{Z_9, Z_8, Z_7, Z_6, Z_4, Z_3, Z_2, Z_1, \bar{D}\},$
 $\{Z_9, Z_8, Z_7, Z_6, Z_5, Z_3, Z_2, Z_1, \bar{D}\}$
(see diagram 50 of the **Figure 2**);
- 51) $\{Z_9, Z_8, Z_7, Z_5, Z_4, Z_3, Z_2, Z_1, \bar{D}\}$
(see diagram 51 of the **Figure 2**);
- 52) $\{Z_9, Z_8, Z_7, Z_6, Z_5, Z_4, Z_3, Z_1, \bar{D}\}$
(see diagram 52 of the **Figure 2**);

Diagrams of subsemilattices of the semilattice D .

Lemma 3. Let $D \in \Sigma_1(X, 10)$. Then the following sets are all XI-subsemi-lattices of the given semilattice D :

- 1) $\{Z_9\}, \{Z_8\}, \{Z_7\}, \{Z_6\}, \{Z_5\}, \{Z_4\}, \{Z_3\}, \{Z_2\}, \{Z_1\}, \{\bar{D}\}$
(see diagram 1 of the **Figure 2**);
- 2) $\{Z_9, \bar{D}\}, \{Z_9, Z_8\}, \{Z_9, Z_7\}, \{Z_9, Z_6\}, \{Z_9, Z_5\}, \{Z_9, Z_4\}, \{Z_9, Z_3\}, \{Z_9, Z_2\}, \{Z_9, Z_1\}, \{Z_8, Z_3\},$
 $\{Z_8, \bar{D}\}, \{Z_7, Z_3\}, \{Z_7, \bar{D}\}, \{Z_6, Z_3\}, \{Z_6, Z_2\}, \{Z_6, Z_1\}, \{Z_6, \bar{D}\}, \{Z_5, Z_1\}, \{Z_5, \bar{D}\}, \{Z_4, Z_1\},$
 $\{Z_4, \bar{D}\}, \{Z_3, \bar{D}\}, \{Z_2, \bar{D}\}, \{Z_1, \bar{D}\}$
(see diagram 2 of the **Figure 2**);
- 3) $\{Z_9, Z_8, \bar{D}\}, \{Z_9, Z_7, \bar{D}\}, \{Z_9, Z_6, \bar{D}\}, \{Z_9, Z_5, \bar{D}\}, \{Z_9, Z_4, \bar{D}\}, \{Z_9, Z_3, \bar{D}\}, \{Z_9, Z_2, \bar{D}\}, \{Z_9, Z_1, \bar{D}\},$
 $\{Z_9, Z_8, Z_3\}, \{Z_9, Z_7, Z_3\}, \{Z_9, Z_6, Z_3\}, \{Z_9, Z_6, Z_2\}, \{Z_9, Z_6, Z_1\}, \{Z_9, Z_5, Z_1\}, \{Z_9, Z_4, Z_1\},$
 $\{Z_8, Z_3, \bar{D}\}, \{Z_7, Z_3, \bar{D}\}, \{Z_6, Z_3, \bar{D}\}, \{Z_6, Z_2, \bar{D}\}, \{Z_6, Z_1, \bar{D}\}, \{Z_5, Z_1, \bar{D}\}, \{Z_4, Z_1, \bar{D}\}$
(see diagram 3 of the **Figure 2**);
- 4) $\{Z_9, Z_4, Z_1, \bar{D}\}, \{Z_9, Z_5, Z_1, \bar{D}\}, \{Z_9, Z_6, Z_1, \bar{D}\}, \{Z_9, Z_6, Z_2, \bar{D}\}, \{Z_9, Z_6, Z_3, \bar{D}\},$
 $\{Z_9, Z_7, Z_3, \bar{D}\}, \{Z_9, Z_8, Z_3, \bar{D}\}$
(see diagram 4 of the **Figure 2**);
- 5) $\{Z_9, Z_5, Z_4, Z_1\}, \{Z_9, Z_6, Z_4, Z_1\}, \{Z_9, Z_6, Z_5, Z_1\}, \{Z_9, Z_7, Z_6, Z_3\}, \{Z_9, Z_8, Z_6, Z_3\},$
 $\{Z_9, Z_8, Z_7, Z_3\}, \{Z_9, Z_8, Z_4, \bar{D}\}, \{Z_9, Z_8, Z_5, \bar{D}\}, \{Z_9, Z_7, Z_2, \bar{D}\}, \{Z_9, Z_7, Z_4, \bar{D}\},$
 $\{Z_9, Z_7, Z_5, \bar{D}\}, \{Z_9, Z_8, Z_1, \bar{D}\}, \{Z_9, Z_8, Z_2, \bar{D}\}, \{Z_9, Z_4, Z_2, \bar{D}\}, \{Z_9, Z_4, Z_3, \bar{D}\},$
 $\{Z_9, Z_5, Z_2, \bar{D}\}, \{Z_9, Z_5, Z_3, \bar{D}\}, \{Z_9, Z_7, Z_1, \bar{D}\}, \{Z_9, Z_2, Z_1, \bar{D}\}, \{Z_9, Z_3, Z_1, \bar{D}\},$
 $\{Z_9, Z_3, Z_2, \bar{D}\}, \{Z_6, Z_2, Z_1, \bar{D}\}, \{Z_6, Z_3, Z_1, \bar{D}\}, \{Z_6, Z_3, Z_2, \bar{D}\}$
(see diagram 5 of the **Figure 2**);
- 6) $\{Z_9, Z_5, Z_4, Z_1, \bar{D}\}, \{Z_9, Z_6, Z_4, Z_1, \bar{D}\}, \{Z_9, Z_6, Z_5, Z_1, \bar{D}\}, \{Z_9, Z_7, Z_6, Z_3, \bar{D}\},$
 $\{Z_9, Z_8, Z_6, Z_3, \bar{D}\}, \{Z_9, Z_8, Z_7, Z_3, \bar{D}\}$
(see diagram 6 of the **Figure 2**);
- 7) $\{Z_9, Z_6, Z_2, Z_1, \bar{D}\}, \{Z_9, Z_6, Z_3, Z_1, \bar{D}\}, \{Z_9, Z_6, Z_3, Z_2, \bar{D}\}$
(see diagram 7 of the **Figure 2**);

- 8) $\{Z_9, Z_8, Z_6, Z_3, Z_2, \bar{D}\}, \{Z_9, Z_8, Z_6, Z_3, Z_1, \bar{D}\}, \{Z_9, Z_7, Z_6, Z_3, Z_2, \bar{D}\}, \{Z_9, Z_7, Z_6, Z_3, Z_1, \bar{D}\},$
 $\{Z_9, Z_6, Z_5, Z_3, Z_1, \bar{D}\}, \{Z_9, Z_6, Z_5, Z_2, Z_1, \bar{D}\}, \{Z_9, Z_6, Z_4, Z_3, Z_1, \bar{D}\}, \{Z_9, Z_6, Z_4, Z_2, Z_1, \bar{D}\}$
 (see diagram 8 of the **Figure 2**);

Proof. It is well known (see [1]), that the semilattices 1 to 8, which are given by lemma 2 are always *XI*-semilattices. The semilattices 9 and 10 which are given by Lemma 2

$$\begin{aligned} &\{Z_8, Z_7, Z_3\}, \{Z_8, Z_6, Z_3\}, \{Z_8, Z_6, \bar{D}\}, \{Z_8, Z_5, \bar{D}\}, \{Z_8, Z_4, \bar{D}\}, \\ &\{Z_8, Z_2, \bar{D}\}, \{Z_8, Z_1, \bar{D}\}, \{Z_7, Z_6, Z_3\}, \{Z_7, Z_5, \bar{D}\}, \{Z_7, Z_4, \bar{D}\}, \\ &\{Z_7, Z_2, \bar{D}\}, \{Z_7, Z_1, \bar{D}\}, \{Z_6, Z_5, Z_1\}, \{Z_6, Z_4, Z_1\}, \{Z_5, Z_4, Z_1\}, \\ &\{Z_5, Z_3, \bar{D}\}, \{Z_5, Z_2, \bar{D}\}, \{Z_4, Z_3, \bar{D}\}, \{Z_4, Z_2, \bar{D}\}, \{Z_3, Z_2, \bar{D}\}, \\ &\{Z_3, Z_1, \bar{D}\}, \{Z_2, Z_1, \bar{D}\}. \end{aligned}$$

(see diagram 9 of the **Figure 2**);

$$\{Z_8, Z_6, Z_3, \bar{D}\}, \{Z_8, Z_7, Z_3, \bar{D}\}, \{Z_7, Z_6, Z_3, \bar{D}\}, \{Z_5, Z_4, Z_1, \bar{D}\}, \{Z_6, Z_4, Z_1, \bar{D}\}, \{Z_6, Z_5, Z_1, \bar{D}\}$$

(see diagram 10 of the **Figure 2**);

are *XI*-semilattices iff the intersection of minimal elements of the given semilattices is empty set. From the formal equalities (1) of the given semilattice D we have

$$\begin{aligned} Z_8 \cap Z_7 &= (P_0 \cup P_1 \cup P_2 \cup P_4 \cup P_5 \cup P_6 \cup P_7 \cup P_9) \cup (P_0 \cup P_1 \cup P_2 \cup P_4 \cup P_5 \cup P_6 \cup P_8 \cup P_9) \neq \emptyset \\ Z_8 \cap Z_6 &= (P_0 \cup P_1 \cup P_2 \cup P_4 \cup P_5 \cup P_6 \cup P_7 \cup P_9) \cup (P_0 \cup P_4 \cup P_5 \cup P_7 \cup P_8 \cup P_9) \neq \emptyset \\ Z_8 \cap Z_5 &= (P_0 \cup P_1 \cup P_2 \cup P_4 \cup P_5 \cup P_6 \cup P_7 \cup P_9) \cup (P_0 \cup P_2 \cup P_3 \cup P_4 \cup P_6 \cup P_7 \cup P_8 \cup P_9) \neq \emptyset \\ Z_8 \cap Z_4 &= (P_0 \cup P_1 \cup P_2 \cup P_4 \cup P_5 \cup P_6 \cup P_7 \cup P_9) \cup (P_0 \cup P_2 \cup P_3 \cup P_5 \cup P_6 \cup P_7 \cup P_8 \cup P_9) \neq \emptyset \\ Z_8 \cap Z_2 &= (P_0 \cup P_1 \cup P_2 \cup P_4 \cup P_5 \cup P_6 \cup P_7 \cup P_9) \cup (P_0 \cup P_1 \cup P_3 \cup P_4 \cup P_5 \cup P_6 \cup P_7 \cup P_8 \cup P_9) \neq \emptyset \\ Z_8 \cap Z_1 &= (P_0 \cup P_1 \cup P_2 \cup P_4 \cup P_5 \cup P_6 \cup P_7 \cup P_9) \cup (P_0 \cup P_2 \cup P_3 \cup P_4 \cup P_5 \cup P_6 \cup P_7 \cup P_8 \cup P_9) \neq \emptyset \\ Z_7 \cap Z_6 &= (P_0 \cup P_1 \cup P_2 \cup P_4 \cup P_5 \cup P_6 \cup P_8 \cup P_9) \cup (P_0 \cup P_4 \cup P_5 \cup P_7 \cup P_8 \cup P_9) \neq \emptyset \\ Z_7 \cap Z_5 &= (P_0 \cup P_1 \cup P_2 \cup P_4 \cup P_5 \cup P_6 \cup P_8 \cup P_9) \cup (P_0 \cup P_2 \cup P_3 \cup P_4 \cup P_6 \cup P_7 \cup P_8 \cup P_9) \neq \emptyset \\ Z_7 \cap Z_4 &= (P_0 \cup P_1 \cup P_2 \cup P_4 \cup P_5 \cup P_6 \cup P_8 \cup P_9) \cup (P_0 \cup P_2 \cup P_3 \cup P_5 \cup P_6 \cup P_7 \cup P_8 \cup P_9) \neq \emptyset \\ Z_7 \cap Z_2 &= (P_0 \cup P_1 \cup P_2 \cup P_4 \cup P_5 \cup P_6 \cup P_8 \cup P_9) \cup (P_0 \cup P_1 \cup P_3 \cup P_4 \cup P_5 \cup P_6 \cup P_7 \cup P_8 \cup P_9) \neq \emptyset \\ Z_7 \cap Z_1 &= (P_0 \cup P_1 \cup P_2 \cup P_4 \cup P_5 \cup P_6 \cup P_8 \cup P_9) \cup (P_0 \cup P_2 \cup P_3 \cup P_4 \cup P_5 \cup P_6 \cup P_7 \cup P_8 \cup P_9) \neq \emptyset \\ Z_6 \cap Z_5 &= (P_0 \cup P_4 \cup P_5 \cup P_7 \cup P_8 \cup P_9) \cup (P_0 \cup P_2 \cup P_3 \cup P_4 \cup P_6 \cup P_7 \cup P_8 \cup P_9) \neq \emptyset \\ Z_6 \cap Z_4 &= (P_0 \cup P_4 \cup P_5 \cup P_7 \cup P_8 \cup P_9) \cup (P_0 \cup P_2 \cup P_3 \cup P_5 \cup P_6 \cup P_7 \cup P_8 \cup P_9) \neq \emptyset \\ Z_5 \cap Z_4 &= (P_0 \cup P_2 \cup P_3 \cup P_4 \cup P_6 \cup P_7 \cup P_8 \cup P_9) \cup (P_0 \cup P_2 \cup P_3 \cup P_5 \cup P_6 \cup P_7 \cup P_8 \cup P_9) \neq \emptyset \\ Z_5 \cap Z_3 &= (P_0 \cup P_2 \cup P_3 \cup P_4 \cup P_6 \cup P_7 \cup P_8 \cup P_9) \cup (P_0 \cup P_1 \cup P_2 \cup P_4 \cup P_5 \cup P_6 \cup P_7 \cup P_8 \cup P_9) \neq \emptyset \\ Z_5 \cap Z_2 &= (P_0 \cup P_2 \cup P_3 \cup P_4 \cup P_6 \cup P_7 \cup P_8 \cup P_9) \cup (P_0 \cup P_1 \cup P_3 \cup P_4 \cup P_5 \cup P_6 \cup P_7 \cup P_8 \cup P_9) \neq \emptyset \\ Z_4 \cap Z_3 &= (P_0 \cup P_2 \cup P_3 \cup P_5 \cup P_6 \cup P_7 \cup P_8 \cup P_9) \cup (P_0 \cup P_1 \cup P_2 \cup P_4 \cup P_5 \cup P_6 \cup P_7 \cup P_8 \cup P_9) \neq \emptyset \end{aligned}$$

$$\begin{aligned}
 Z_4 \cap Z_2 &= (P_0 \cup P_2 \cup P_3 \cup P_5 \cup P_6 \cup P_7 \cup P_8 \cup P_9) \cup (P_0 \cup P_1 \cup P_3 \cup P_4 \cup P_5 \cup P_6 \cup P_7 \cup P_8 \cup P_9) \neq \emptyset \\
 Z_3 \cap Z_2 &= (P_0 \cup P_1 \cup P_2 \cup P_4 \cup P_5 \cup P_6 \cup P_7 \cup P_8 \cup P_9) \cup (P_0 \cup P_1 \cup P_3 \cup P_4 \cup P_5 \cup P_6 \cup P_7 \cup P_8 \cup P_9) \neq \emptyset \\
 Z_3 \cap Z_1 &= (P_0 \cup P_1 \cup P_2 \cup P_4 \cup P_5 \cup P_6 \cup P_7 \cup P_8 \cup P_9) \cup (P_0 \cup P_2 \cup P_3 \cup P_4 \cup P_5 \cup P_6 \cup P_7 \cup P_8 \cup P_9) \neq \emptyset \\
 Z_2 \cap Z_1 &= (P_0 \cup P_1 \cup P_3 \cup P_4 \cup P_5 \cup P_6 \cup P_7 \cup P_8 \cup P_9) \cup (P_0 \cup P_2 \cup P_3 \cup P_4 \cup P_5 \cup P_6 \cup P_7 \cup P_8 \cup P_9) \neq \emptyset
 \end{aligned}$$

From the equalities given above it follows that the semilattices 9 and 10 are not *XI*-semilattices. \square
 The semilattices 11

$$\begin{aligned}
 &\{Z_6, Z_5, Z_3, Z_1, \bar{D}\}, \{Z_6, Z_5, Z_2, Z_1, \bar{D}\}, \{Z_6, Z_4, Z_3, Z_1, \bar{D}\}, \{Z_6, Z_4, Z_2, Z_1, \bar{D}\}, \\
 &\{Z_7, Z_6, Z_3, Z_2, \bar{D}\}, \{Z_7, Z_6, Z_3, Z_1, \bar{D}\}, \{Z_8, Z_6, Z_3, Z_2, \bar{D}\}, \{Z_8, Z_6, Z_3, Z_1, \bar{D}\}.
 \end{aligned}$$

(see diagram 1-8 of the **Figure 3**);
 are not *XI*-semilattice since we have the following inequalities

$$\begin{aligned}
 &Z_5 \cap Z_3 \neq \emptyset, Z_5 \cap Z_2 \neq \emptyset, Z_4 \cap Z_3 \neq \emptyset, Z_4 \cap Z_2 \neq \emptyset, \\
 &Z_7 \cap Z_2 \neq \emptyset, Z_7 \cap Z_1 \neq \emptyset, Z_8 \cap Z_2 \neq \emptyset, Z_8 \cap Z_1 \neq \emptyset.
 \end{aligned}$$

The semilattices 12 to 52 are never *XI*-semilattices. We prove that the semilattice, diagram 52 of the **Figure 2**, is not an *XI*-semilattice (see **Figure 4**). Indeed, let $Q = \{T_0, T_1, T_2, T_3, T_4, T_5, T_6, T_7, T_8\}$ and

$$C(Q) = \{P'_0, P'_1, P'_2, P'_3, P'_4, P'_5, P'_6, P'_7, P'_8\}$$

be a family of sets, where $P'_0, P'_1, P'_2, P'_3, P'_4, P'_5, P'_6, P'_7, P'_8$ are pairwise disjoint subsets of the set X . Let

$$\varphi = \begin{pmatrix} T_0 & T_1 & T_2 & T_3 & T_4 & T_5 & T_6 & T_7 & T_8 \\ P'_0 & P'_1 & P'_2 & P'_3 & P'_4 & P'_5 & P'_6 & P'_7 & P'_8 \end{pmatrix}$$

be a mapping of the semilattice Q onto the family of sets $C(Q)$. Then for the formal equalities of the semilattice Q we have a form:




Figure 3. Diagram of all subsemilattices which are isomorphic to 11 in **Figure 2**.




Figure 4. Diagram of subsemilattice 52 in **Figure 2**.

$$\begin{aligned}
T_0 &= P'_0 \cup P'_1 \cup P'_2 \cup P'_3 \cup P'_4 \cup P'_5 \cup P'_6 \cup P'_7 \cup P'_8, \\
T_1 &= P'_0 \cup P'_2 \cup P'_3 \cup P'_4 \cup P'_5 \cup P'_6 \cup P'_7 \cup P'_8, \\
T_2 &= P'_0 \cup P'_1 \cup P'_3 \cup P'_4 \cup P'_5 \cup P'_6 \cup P'_7 \cup P'_8, \\
T_3 &= P'_0 \cup P'_2 \cup P'_4 \cup P'_5 \cup P'_6 \cup P'_7 \cup P'_8, \\
T_4 &= P'_0 \cup P'_2 \cup P'_3 \cup P'_5 \cup P'_6 \cup P'_7 \cup P'_8, \\
T_5 &= P'_0 \cup P'_3 \cup P'_4 \cup P'_6 \cup P'_7 \cup P'_8, \\
T_6 &= P'_0 \cup P'_1 \cup P'_3 \cup P'_4 \cup P'_5 \cup P'_7 \cup P'_8, \\
T_7 &= P'_0 \cup P'_1 \cup P'_3 \cup P'_4 \cup P'_5 \cup P'_6 \cup P'_8, \\
T_8 &= P'_0.
\end{aligned} \tag{3}$$

Here the elements $P'_1, P'_2, P'_3, P'_4, P'_6, P'_7$ are basis sources, the elements P'_0, P'_5, P'_8 are sources of completeness of the semilattice D . Therefore $|X| \geq 6$ and $\delta = 7$ (see [2]). Then of the formal equalities we have:

$$Q_t = \begin{cases} Q, & \text{if } t \in P'_0, \\ \{T_7, T_6, T_2, T_0\}, & \text{if } t \in P'_1, \\ \{T_4, T_3, T_1, T_0\}, & \text{if } t \in P'_2, \\ \{T_7, T_6, T_5, T_4, T_2, T_1, T_0\}, & \text{if } t \in P'_3, \\ \{T_7, T_6, T_5, T_3, T_2, T_1, T_0\}, & \text{if } t \in P'_4, \\ \{T_7, T_6, T_4, T_3, T_2, T_1, T_0\}, & \text{if } t \in P'_5, \\ \{T_7, T_5, T_4, T_3, T_2, T_1, T_0\}, & \text{if } t \in P'_6, \\ \{T_6, T_5, T_4, T_3, T_2, T_1, T_0\}, & \text{if } t \in P'_7, \\ \{T_8, T_7, T_6, T_5, T_4, T_3, T_2, T_1, T_0\}, & \text{if } t \in P'_8. \end{cases}$$

$$\Lambda(Q, Q_t) = \begin{cases} T_8, & \text{if } t \in P'_0, \\ T_8, & \text{if } t \in P'_1, \\ T_8, & \text{if } t \in P'_2, \\ T_8, & \text{if } t \in P'_3, \\ T_8, & \text{if } t \in P'_4, \\ T_8, & \text{if } t \in P'_5, \\ T_8, & \text{if } t \in P'_6, \\ T_8, & \text{if } t \in P'_7, \\ T_8, & \text{if } t \in P'_8. \end{cases}$$

We have, that $Q^\wedge = \{T_8\}$ and $\Lambda(Q, Q_t) \in Q$ for any $t \in Q$. But elements $T_7, T_6, T_5, T_4, T_3, T_2, T_1, T_0$ are not union of some elements of the set Q^\wedge . Therefore from the Definition 1 it follows that Q is not an XI-semilattice of unions. Statements 12 to 51 can be proved analogously.

We denoted the following semilattices by symbols:

- $Q_1 = \{T\}$, where $T \in D$ (see diagram 1 of the Figure 5);
- $Q_2 = \{T, T'\}$, where $T, T' \in D$ and $T \subset T'$ (see diagram 2 of the Figure 5);
- $Q_3 = \{T, T', T''\}$, where $T, T', T'' \in D$ and $T \subset T' \subset T''$ (see diagram 3 of the Figure 5);
- $Q_4 = \{Z_9, T, T', \tilde{D}\}$, where $T, T' \in D$ and $Z_9 \subset T \subset T' \subset \tilde{D}$ (see diagram 4 of the Figure 5);
- $Q_5 = \{T, T', T'', T' \cup T''\}$ where $T, T', T'' \in D$, $T \subset T'$, $T \subset T''$, $T' \setminus T'' \neq \emptyset$, $T'' \setminus T' \neq \emptyset$, (see diagram 5 of the Figure 5);
- $Q_6 = \{Z_9, T, T', T \cup T', \tilde{D}\}$, where $T, T' \in D$, $Z_9 \subset T$, $Z_9 \subset T'$, $T \setminus T' \neq \emptyset$, $T' \setminus T \neq \emptyset$ (see diagram 6 of the Figure 5);
- $Q_7 = \{Z_9, Z_6, T, T', \tilde{D}\}$, where $T, T' \in D$, $Z_6 \subset T$, $Z_6 \subset T'$, $T \setminus T' \neq \emptyset$, $T' \setminus T \neq \emptyset$, $T \cup T' = \tilde{D}$ (see diagram 7 of the Figure 5);




Figure 5. Diagram of all XI-subsemilattices of D .

h) $Q_8 = \{Z_9, T, T', T \cup T', Z, \bar{D}\}$, where $Z_9 \subset T' \subset Z$, $T \setminus T' \neq \emptyset$, $T' \setminus T \neq \emptyset$, $(T \cup T') \setminus Z \neq \emptyset$, $Z \setminus (T \cup T') \neq \emptyset$ (see diagram 8 of the **Figure 5**);

Note that the semilattices in **Figure 5** are all XI-semilattices (see [1] and Lemma 1.2.3).

Definition 9. Let us assume that by the symbol $\Sigma'_{XI}(X, D)$ denote a set of all XI-subsemilattices of X -semilattices of unions D that every element of this set contains an empty set if $\emptyset \in D$ or denotes a set of all XI-subsemilattices of D .

Further, let $D', D'' \in \Sigma'_{XI}(X, D)$ and $\mathcal{G}_{XI} \subseteq \Sigma'_{XI}(X, D) \times \Sigma'_{XI}(X, D)$. It is assumed that $D' \mathcal{G}_{XI} D''$ iff there exists some complete isomorphism φ between the semilattices D' and D'' . One can easily verify that the binary relation \mathcal{G}_{XI} is an equivalence relation on the set $\Sigma'_{XI}(X, D)$.

By the symbol $Q_i \mathcal{G}_{XI}$ denote the \mathcal{G}_{XI} -equivalence class of the set $\Sigma'_{XI}(X, D)$, where every element is isomorphic to the X -semilattice Q_i ($i = 1, 2, \dots, 8$).

Let D' be an XI-subsemilattice of the semilattice D . By $I(D')$ we denoted the set of all right units of the semigroup $B_X(D')$, and

$$|I^*(Q_i)| = \sum_{D' \in Q_i, \mathcal{G}_{XI}} |I(D')|$$

where $i = 1, 2, \dots, 8$.

Lemma 4. If X is a finite set, then the following equalities hold

- $|I(Q_1)| = 1$
- $|I(Q_2)| = (2^{|T' \setminus T|} - 1) \cdot 2^{|X \setminus T'|}$
- $|I(Q_3)| = (2^{|T' \setminus T|} - 1) \cdot (3^{|T'' \setminus T'|} - 2^{|T'' \setminus T'|}) \cdot 3^{|X \setminus T''|}$
- $|I(Q_4)| = (2^{|T \setminus Z_9|} - 1) \cdot (3^{|T' \setminus T|} - 2^{|T' \setminus T|}) \cdot (4^{|T \setminus T'|} - 3^{|T \setminus T'|}) \cdot 4^{|X \setminus T|}$
- $|I(Q_5)| = (2^{|T' \setminus T''|} - 1) \cdot (2^{|T'' \setminus T'|} - 1) \cdot 4^{|X \setminus (T' \cup T'')|}$
- $|I(Q_6)| = (2^{|T' \setminus T''|} - 1) \cdot (2^{|T'' \setminus T'|} - 1) \cdot (5^{|D \setminus (T \cup T'')|} - 4^{|D \setminus (T \cup T'')|}) \cdot 5^{|X \setminus D|}$
- $|I(Q_7)| = (2^{|Z_6 \setminus Z_9|} - 1) \cdot 2^{|(T \cap T') \setminus Z_6|} \cdot (3^{|T \setminus T'|} - 2^{|T \setminus T'|}) \cdot (3^{|T' \setminus T|} - 2^{|T' \setminus T|}) \cdot 5^{|X \setminus D|}$
- $|I(Q_8)| = (2^{|T \setminus Z|} - 1) \cdot (2^{|T' \setminus T|} - 1) \cdot (3^{|Z \setminus (T \cup T')|} - 2^{|Z \setminus (T \cup T')|}) \cdot 6^{|X \setminus D|}$

Proof. This lemma immediately follows from Theorem 13.1.2, 13.3.2, and 13.7.2 of the [1]. \square

Theorem 10. Let $D \in \Sigma_1(X, 10)$ and $\alpha \in B_X(D)$. Binary relation α is an idempotent relation of the semigroup $B_X(D)$ iff binary relation α satisfies only one conditions of the following conditions:

- $\alpha = X \times T$, where $T \in D$;
- $\alpha = (Y_T^\alpha \times T) \cup (Y_{T'}^\alpha \times T')$, where $T, T' \in D$, $T \subset T'$, $Y_T^\alpha, Y_{T'}^\alpha \notin \{\emptyset\}$, and satisfies the conditions: $Y_T^\alpha \supseteq T$, $Y_{T'}^\alpha \cap T' \neq \emptyset$;
- $\alpha = (Y_T^\alpha \times T) \cup (Y_{T'}^\alpha \times T') \cup (Y_{T''}^\alpha \times T'')$, where $T, T', T'' \in D$, $T \subset T' \subset T''$, $Y_T^\alpha, Y_{T'}^\alpha, Y_{T''}^\alpha \notin \{\emptyset\}$, and satisfies the conditions: $Y_T^\alpha \supseteq T$, $Y_T^\alpha \cup Y_{T'}^\alpha \supseteq T'$, $Y_{T'}^\alpha \cap T' \neq \emptyset$, $Y_{T''}^\alpha \cap T'' \neq \emptyset$;

- d) $\alpha = (Y_9^\alpha \times Z_9) \cup (Y_T^\alpha \times T) \cup (Y_{T'}^\alpha \times T') \cup (Y_0^\alpha \times \bar{D})$, where $T, T' \in D$, $Z_9 \subset T \subset T' \subset \bar{D}$, $Y_9^\alpha, Y_T^\alpha, Y_{T'}^\alpha, Y_0^\alpha \notin \{\emptyset\}$, and satisfies the conditions: $Y_9^\alpha \supseteq Z_9$, $Y_9^\alpha \cup Y_T^\alpha \supseteq T$, $Y_9^\alpha \cup Y_{T'}^\alpha \cup Y_T^\alpha \supseteq T'$, $Y_T^\alpha \cap T \neq \emptyset$, $Y_{T'}^\alpha \cap T' \neq \emptyset$, $Y_0^\alpha \cap \bar{D} \neq \emptyset$;
- e) $\alpha = (Y_T^\alpha \times T) \cup (Y_{T'}^\alpha \times T') \cup (Y_{T''}^\alpha \times T'') \cup (Y_{T' \cup T''}^\alpha \times (T' \cup T''))$, where $T, T', T'' \in D$, $T \subset T'$, $T \subset T''$, $T' \setminus T'' \neq \emptyset$, $T'' \setminus T' \neq \emptyset$, $Y_T^\alpha, Y_{T'}^\alpha, Y_{T''}^\alpha \notin \{\emptyset\}$ and satisfies the conditions: $Y_T^\alpha \cup Y_{T'}^\alpha \supseteq T'$, $Y_T^\alpha \cup Y_{T''}^\alpha \supseteq T''$, $Y_{T'}^\alpha \cap T' \neq \emptyset$, $Y_{T''}^\alpha \cap T'' \neq \emptyset$;
- f) $\alpha = (Y_9^\alpha \times Z_9) \cup (Y_T^\alpha \times T) \cup (Y_{T'}^\alpha \times T') \cup (Y_{T \cup T'}^\alpha \times (T \cup T')) \cup (Y_0^\alpha \times \bar{D})$, where $Z_9 \subset T$, $Z_9 \subset T'$, $T \setminus T' \neq \emptyset$, $T' \setminus T \neq \emptyset$, $Y_T^\alpha, Y_{T'}^\alpha, Y_0^\alpha \notin \{\emptyset\}$ and satisfies the conditions: $Y_9^\alpha \cup Y_T^\alpha \supseteq T$, $Y_9^\alpha \cup Y_{T'}^\alpha \supseteq T'$, $Y_T^\alpha \cap T \neq \emptyset$, $Y_{T'}^\alpha \cap T' \neq \emptyset$, $Y_0^\alpha \cap \bar{D} \neq \emptyset$;
- g) $\alpha = (Y_9^\alpha \times Z_9) \cup (Y_6^\alpha \times Z_6) \cup (Y_T^\alpha \times T) \cup (Y_{T'}^\alpha \times T') \cup (Y_0^\alpha \times \bar{D})$, where $T, T' \in D$, $Z_6 \subset T$, $Z_6 \subset T'$, $T \setminus T' \neq \emptyset$, $T' \setminus T \neq \emptyset$, $T \cup T' = \bar{D}$, $Y_6^\alpha, Y_T^\alpha, Y_{T'}^\alpha \notin \{\emptyset\}$ and satisfies the conditions: $Y_9^\alpha \supseteq Z_9$, $Y_9^\alpha \cup Y_6^\alpha \supseteq Z_6$, $Y_9^\alpha \cup Y_6^\alpha \cup Y_T^\alpha \supseteq T$, $Y_9^\alpha \cup Y_6^\alpha \cup Y_{T'}^\alpha \supseteq T'$, $Y_6^\alpha \cap Z_6 \neq \emptyset$, $Y_T^\alpha \cap T \neq \emptyset$, $Y_{T'}^\alpha \cap T' \neq \emptyset$;
- h) $\alpha = (Y_9^\alpha \times Z_9) \cup (Y_T^\alpha \times T) \cup (Y_{T'}^\alpha \times T') \cup (Y_{T \cup T'}^\alpha \times (T \cup T')) \cup (Y_Z^\alpha \times Z) \cup (Y_0^\alpha \times \bar{D})$, where $Z_9 \subset T' \subset Z$, $T \setminus T' \neq \emptyset$, $T' \setminus T \neq \emptyset$, $(T \cup T') \setminus Z \neq \emptyset$, $Z \setminus (T \cup T') \neq \emptyset$, $Y_T^\alpha, Y_{T'}^\alpha, Y_Z^\alpha \notin \{\emptyset\}$ and satisfies the conditions: $Y_9^\alpha \cup Y_T^\alpha \supseteq T$, $Y_9^\alpha \cup Y_{T'}^\alpha \supseteq T'$, $Y_9^\alpha \cup Y_{T'}^\alpha \cup Y_Z^\alpha \supseteq Z$, $Y_T^\alpha \cap T \neq \emptyset$, $Y_{T'}^\alpha \cap T' \neq \emptyset$, $Y_Z^\alpha \cap Z \neq \emptyset$.

Proof. By Lemma 3 we know that 1 to 8 are an XL -semilattices. We prove only statement g. Indeed, if

$$\alpha = (Y_9^\alpha \times Z_9) \cup (Y_6^\alpha \times Z_6) \cup (Y_T^\alpha \times T) \cup (Y_{T'}^\alpha \times T') \cup (Y_0^\alpha \times \bar{D}),$$

where $Y_6^\alpha, Y_T^\alpha, Y_{T'}^\alpha \notin \{\emptyset\}$, then it is easy to see, that the set $D(\alpha) = \{Z_9, Z_6, T, T'\}$ is a generating set of the semilattice $\{Z_9, Z_6, T, T', \bar{D}\}$. Then the following equalities hold

$$\ddot{D}(\alpha)_{Z_9} = \{Z_9\}, \quad \ddot{D}(\alpha)_{Z_6} = \{Z_9, Z_6\},$$

$$\ddot{D}(\alpha)_T = \{Z_9, Z_6, T\}, \quad \ddot{D}(\alpha)_{T'} = \{Z_9, Z_6, T'\}.$$

By statement a of the Theorem 6.2.1 (see [1]) we have:

$$Y_9^\alpha \supseteq Z_9, \quad Y_9^\alpha \cup Y_6^\alpha \supseteq Z_6, \quad Y_9^\alpha \cup Y_6^\alpha \cup Y_T^\alpha \supseteq T, \quad Y_9^\alpha \cup Y_6^\alpha \cup Y_{T'}^\alpha \supseteq T'.$$

Further, one can see, that the equalities are true:

$$l(\ddot{D}(\alpha)_{Z_6}, Z_6) = \cup(\ddot{D}(\alpha)_{Z_6} \setminus \{Z_6\}) = Z_9, \quad Z_6 \setminus l(\ddot{D}(\alpha)_{Z_6}, Z_6) = Z_6 \setminus Z_9 \neq \emptyset,$$

$$l(\ddot{D}(\alpha)_T, T) = \cup(\ddot{D}(\alpha)_T \setminus \{T\}) = Z_6, \quad T \setminus l(\ddot{D}(\alpha)_T, T) = T \setminus Z_6 \neq \emptyset,$$

$$l(\ddot{D}(\alpha)_{T'}, T') = \cup(\ddot{D}(\alpha)_{T'} \setminus \{T'\}) = Z_6, \quad T' \setminus l(\ddot{D}(\alpha)_{T'}, T') = T' \setminus Z_6 \neq \emptyset,$$

We have the elements Z_6, T, T' are nonlimiting elements of the sets $\ddot{D}(\alpha)_{Z_6}, \ddot{D}(\alpha)_T, \ddot{D}(\alpha)_{T'}$ respectively.

By statement b of the Theorem 6.2.1 [1] it follows, that the conditions $Y_6^\alpha \cap Z_6 \neq \emptyset$, $Y_T^\alpha \cap T \neq \emptyset$, $Y_{T'}^\alpha \cap T' \neq \emptyset$ hold. Therefore, the statement g is proved. Rest of statements can be proved analogously.

Lemma 5. Let $D \in \Sigma_1(X, 10)$ and $Z_9 \neq \emptyset$. If X is a finite set, then the number $|I^*(Q_1)|$ may be calculated by the formula $|I^*(Q_1)| = 10$.

Lemma 6. Let $D \in \Sigma_1(X, 10)$ and $Z_9 \neq \emptyset$. If X is a finite set, then the number $|I^*(Q_2)|$ may be calculated by formula

$$\begin{aligned} |I^*(Q_2)| = & \left(2^{|Z_8 \setminus Z_9|} - 1 \right) \cdot 2^{|X \setminus Z_8|} + \left(2^{|Z_7 \setminus Z_9|} - 1 \right) \cdot 2^{|X \setminus Z_7|} + \left(2^{|Z_6 \setminus Z_9|} - 1 \right) \cdot 2^{|X \setminus Z_6|} + \left(2^{|Z_5 \setminus Z_9|} - 1 \right) \cdot 2^{|X \setminus Z_5|} \\ & + \left(2^{|Z_4 \setminus Z_9|} - 1 \right) \cdot 2^{|X \setminus Z_4|} + \left(2^{|Z_3 \setminus Z_9|} + 2^{|Z_3 \setminus Z_8|} + 2^{|Z_3 \setminus Z_7|} + 2^{|Z_3 \setminus Z_6|} - 4 \right) \cdot 2^{|X \setminus Z_3|} \\ & + \left(2^{|Z_2 \setminus Z_9|} + 2^{|Z_2 \setminus Z_6|} - 2 \right) \cdot 2^{|X \setminus Z_2|} + \left(2^{|Z_1 \setminus Z_9|} + 2^{|Z_1 \setminus Z_6|} + 2^{|Z_1 \setminus Z_5|} + 2^{|Z_1 \setminus Z_4|} - 4 \right) \cdot 2^{|X \setminus Z_1|} \\ & + \left(2^{|D \setminus Z_9|} + 2^{|D \setminus Z_8|} + 2^{|D \setminus Z_7|} + 2^{|D \setminus Z_6|} + 2^{|D \setminus Z_5|} + 2^{|D \setminus Z_4|} + 2^{|D \setminus Z_3|} + 2^{|D \setminus Z_2|} + 2^{|D \setminus Z_1|} - 9 \right) \cdot 2^{|X \setminus D|}. \end{aligned}$$

$$\begin{aligned}
|I^*(Q_5)| = & (2^{|Z_5 \setminus Z_4|} - 1) \cdot (2^{|Z_4 \setminus Z_5|} - 1) \cdot 4^{|X \setminus Z_1|} + (2^{|Z_6 \setminus Z_4|} - 1) \cdot (2^{|Z_4 \setminus Z_6|} - 1) \cdot 4^{|X \setminus Z_1|} \\
& + (2^{|Z_6 \setminus Z_5|} - 1) \cdot (2^{|Z_5 \setminus Z_6|} - 1) \cdot 4^{|X \setminus Z_1|} + (2^{|Z_7 \setminus Z_6|} - 1) \cdot (2^{|Z_6 \setminus Z_7|} - 1) \cdot 4^{|X \setminus Z_3|} \\
& + (2^{|Z_8 \setminus Z_6|} - 1) \cdot (2^{|Z_6 \setminus Z_8|} - 1) \cdot 4^{|X \setminus Z_3|} + (2^{|Z_8 \setminus Z_7|} - 1) \cdot (2^{|Z_7 \setminus Z_8|} - 1) \cdot 4^{|X \setminus Z_3|} \\
& + (2^{|Z_8 \setminus Z_4|} - 1) \cdot (2^{|Z_4 \setminus Z_8|} - 1) \cdot 4^{|X \setminus \bar{D}|} + (2^{|Z_8 \setminus Z_5|} - 1) \cdot (2^{|Z_5 \setminus Z_8|} - 1) \cdot 4^{|X \setminus \bar{D}|} \\
& + (2^{|Z_7 \setminus Z_2|} - 1) \cdot (2^{|Z_2 \setminus Z_7|} - 1) \cdot 4^{|X \setminus \bar{D}|} + (2^{|Z_7 \setminus Z_4|} - 1) \cdot (2^{|Z_4 \setminus Z_7|} - 1) \cdot 4^{|X \setminus \bar{D}|} \\
& + (2^{|Z_7 \setminus Z_5|} - 1) \cdot (2^{|Z_5 \setminus Z_7|} - 1) \cdot 4^{|X \setminus \bar{D}|} + (2^{|Z_8 \setminus Z_1|} - 1) \cdot (2^{|Z_1 \setminus Z_8|} - 1) \cdot 4^{|X \setminus \bar{D}|} \\
& + (2^{|Z_8 \setminus Z_2|} - 1) \cdot (2^{|Z_2 \setminus Z_8|} - 1) \cdot 4^{|X \setminus \bar{D}|} + (2^{|Z_4 \setminus Z_2|} - 1) \cdot (2^{|Z_2 \setminus Z_4|} - 1) \cdot 4^{|X \setminus \bar{D}|} \\
& + (2^{|Z_4 \setminus Z_3|} - 1) \cdot (2^{|Z_3 \setminus Z_4|} - 1) \cdot 4^{|X \setminus \bar{D}|} + (2^{|Z_5 \setminus Z_2|} - 1) \cdot (2^{|Z_2 \setminus Z_5|} - 1) \cdot 4^{|X \setminus \bar{D}|} \\
& + (2^{|Z_5 \setminus Z_3|} - 1) \cdot (2^{|Z_3 \setminus Z_5|} - 1) \cdot 4^{|X \setminus \bar{D}|} + (2^{|Z_7 \setminus Z_1|} - 1) \cdot (2^{|Z_1 \setminus Z_7|} - 1) \cdot 4^{|X \setminus \bar{D}|} \\
& + 2 \cdot (2^{|Z_2 \setminus Z_1|} - 1) \cdot (2^{|Z_1 \setminus Z_2|} - 1) \cdot 4^{|X \setminus \bar{D}|} + 2 \cdot (2^{|Z_3 \setminus Z_1|} - 1) \cdot (2^{|Z_1 \setminus Z_3|} - 1) \cdot 4^{|X \setminus \bar{D}|} \\
& + 2 \cdot (2^{|Z_3 \setminus Z_2|} - 1) \cdot (2^{|Z_2 \setminus Z_3|} - 1) \cdot 4^{|X \setminus \bar{D}|}.
\end{aligned}$$

Lemma 10. Let $D \in \Sigma_1(X, 10)$ and $Z_9 \neq \emptyset$. If X is a finite set, then the number $|I^*(Q_6)|$ may be calculated by formula

$$|I^*(Q_6)| = 1 + 3 + 3 + 3 + 3 + 1 = 14$$

Lemma 11. Let $D \in \Sigma_1(X, 10)$ and $Z_9 \neq \emptyset$. If X is a finite set, then the number $|I^*(Q_7)|$ may be calculated by formula

$$\begin{aligned}
|I^*(Q_7)| = & (2^{|Z_6 \setminus Z_9|} - 1) \cdot 2^{|(Z_2 \cap Z_1) \setminus Z_6|} \cdot (3^{|Z_2 \setminus Z_1|} - 2^{|Z_2 \setminus Z_1|}) \cdot (3^{|Z_1 \setminus Z_2|} - 2^{|Z_1 \setminus Z_2|}) \cdot 5^{|X \setminus \bar{D}|} \\
& + (2^{|Z_6 \setminus Z_9|} - 1) \cdot 2^{|(Z_3 \cap Z_1) \setminus Z_6|} \cdot (3^{|Z_3 \setminus Z_1|} - 2^{|Z_3 \setminus Z_1|}) \cdot (3^{|Z_1 \setminus Z_3|} - 2^{|Z_1 \setminus Z_3|}) \cdot 5^{|X \setminus \bar{D}|} \\
& + (2^{|Z_6 \setminus Z_9|} - 1) \cdot 2^{|(Z_3 \cap Z_2) \setminus Z_6|} \cdot (3^{|Z_3 \setminus Z_2|} - 2^{|Z_3 \setminus Z_2|}) \cdot (3^{|Z_2 \setminus Z_3|} - 2^{|Z_2 \setminus Z_3|}) \cdot 5^{|X \setminus \bar{D}|}.
\end{aligned}$$

Lemma 12. Let $D \in \Sigma_1(X, 10)$ and $Z_9 \neq \emptyset$. If X is a finite set, then the number $|I^*(Q_8)|$ may be calculated by formula

$$\begin{aligned}
|I^*(Q_8)| = & (2^{|Z_8 \setminus Z_2|} - 1) \cdot (2^{|Z_6 \setminus Z_8|} - 1) \cdot (3^{|Z_2 \setminus Z_3|} - 2^{|Z_2 \setminus Z_3|}) \cdot 6^{|X \setminus \bar{D}|} \\
& + (2^{|Z_8 \setminus Z_1|} - 1) \cdot (2^{|Z_6 \setminus Z_8|} - 1) \cdot (3^{|Z_1 \setminus Z_3|} - 2^{|Z_1 \setminus Z_3|}) \cdot 6^{|X \setminus \bar{D}|} \\
& + (2^{|Z_7 \setminus Z_2|} - 1) \cdot (2^{|Z_6 \setminus Z_7|} - 1) \cdot (3^{|Z_2 \setminus Z_3|} - 2^{|Z_2 \setminus Z_3|}) \cdot 6^{|X \setminus \bar{D}|} \\
& + (2^{|Z_7 \setminus Z_1|} - 1) \cdot (2^{|Z_6 \setminus Z_7|} - 1) \cdot (3^{|Z_1 \setminus Z_3|} - 2^{|Z_1 \setminus Z_3|}) \cdot 6^{|X \setminus \bar{D}|} \\
& + (2^{|Z_5 \setminus Z_3|} - 1) \cdot (2^{|Z_6 \setminus Z_5|} - 1) \cdot (3^{|Z_3 \setminus Z_1|} - 2^{|Z_3 \setminus Z_1|}) \cdot 6^{|X \setminus \bar{D}|} \\
& + (2^{|Z_5 \setminus Z_2|} - 1) \cdot (2^{|Z_6 \setminus Z_5|} - 1) \cdot (3^{|Z_2 \setminus Z_1|} - 2^{|Z_2 \setminus Z_1|}) \cdot 6^{|X \setminus \bar{D}|} \\
& + (2^{|Z_4 \setminus Z_3|} - 1) \cdot (2^{|Z_6 \setminus Z_4|} - 1) \cdot (3^{|Z_3 \setminus Z_1|} - 2^{|Z_3 \setminus Z_1|}) \cdot 6^{|X \setminus \bar{D}|} \\
& + (2^{|Z_4 \setminus Z_2|} - 1) \cdot (2^{|Z_6 \setminus Z_4|} - 1) \cdot (3^{|Z_2 \setminus Z_1|} - 2^{|Z_2 \setminus Z_1|}) \cdot 6^{|X \setminus \bar{D}|}.
\end{aligned}$$

Figure 6 shows all XI -subsemilattices with six elements.




Figure 6. Diagram of all subsemilattices which are isomorphic.

Theorem 11. Let $D \in \Sigma_1(X, 10)$, $Z_9 \neq \emptyset$. If X is a finite set and I_D is a set of all idempotent elements of the semigroup $B_X(D)$. Then $|I_D| = \sum_{i=1}^8 |I^*(Q_i)|$.

Example 12. Let $X = \{1, 2, 3, 4, 5, 6, 7, 8\}$,

$$P_0 = \{6\}, P_1 = \{1\}, P_2 = \{2\}, P_3 = \{3\}, P_4 = \{4\}, \\ P_5 = \{5\}, P_7 = \{7\}, P_8 = \{8\}, P_9 = P_6 = \emptyset.$$

Then $\bar{D} = \{1, 2, 3, 4, 5, 6, 7, 8\}$, $Z_1 = \{2, 3, 4, 5, 6, 7, 8\}$, $Z_2 = \{1, 3, 4, 5, 6, 7, 8\}$, $Z_3 = \{1, 2, 4, 5, 6, 7, 8\}$, $Z_4 = \{2, 3, 5, 6, 7, 8\}$, $Z_5 = \{2, 3, 4, 6, 7, 8\}$, $Z_6 = \{4, 5, 6, 7, 8\}$, $Z_7 = \{1, 2, 4, 5, 6, 8\}$, $Z_8 = \{1, 2, 4, 5, 6, 7\}$ and $Z_9 = \{6\}$.

$$D = \{\{1, 2, 3, 4, 5, 6, 7, 8\}, \{2, 3, 4, 5, 6, 7, 8\}, \{1, 3, 4, 5, 6, 7, 8\}, \{1, 2, 4, 5, 6, 7, 8\}, \{2, 3, 5, 6, 7, 8\}, \\ \{2, 3, 4, 6, 7, 8\}, \{4, 5, 6, 7, 8\}, \{1, 2, 4, 5, 6, 8\}, \{1, 2, 4, 5, 6, 7\}, \{6\}\}$$

We have $Z_9 \neq \emptyset$. Where $|I^*(Q_1)| = 10$, $|I^*(Q_2)| = 1169$, $|I^*(Q_3)| = 2154$, $|I^*(Q_4)| = 349$, $|I^*(Q_5)| = 122$, $|I^*(Q_6)| = 14$, $|I^*(Q_7)| = 90$, $|I(Q_8)| = 8$, $|I_D| = 3916$.

3. Results

Lemma 13. Let $D \in \Sigma_1(X, 10)$ and $Z_9 = \emptyset$. Then the following sets exhaust all subsemilattices of the semilattice $D = \{Z_9, Z_8, Z_7, Z_6, Z_5, Z_4, Z_3, Z_2, Z_1, \bar{D}\}$ which contains the empty set:

- 1) $\{\emptyset\}$
(see diagram 1 of the **Figure 2**);
- 2) $\{\emptyset, \bar{D}\}$, $\{\emptyset, Z_8\}$, $\{\emptyset, Z_7\}$, $\{\emptyset, Z_6\}$, $\{\emptyset, Z_5\}$, $\{\emptyset, Z_4\}$, $\{\emptyset, Z_3\}$, $\{\emptyset, Z_2\}$, $\{\emptyset, Z_1\}$
(see diagram 2 of the **Figure 2**);
- 3) $\{\emptyset, Z_8, \bar{D}\}$, $\{\emptyset, Z_7, \bar{D}\}$, $\{\emptyset, Z_6, \bar{D}\}$, $\{\emptyset, Z_5, \bar{D}\}$, $\{\emptyset, Z_4, \bar{D}\}$, $\{\emptyset, Z_3, \bar{D}\}$, $\{\emptyset, Z_2, \bar{D}\}$, $\{\emptyset, Z_1, \bar{D}\}$, $\{\emptyset, Z_8, Z_3\}$, $\{\emptyset, Z_7, Z_3\}$, $\{\emptyset, Z_6, Z_3\}$, $\{\emptyset, Z_6, Z_2\}$, $\{\emptyset, Z_6, Z_1\}$, $\{\emptyset, Z_5, Z_1\}$, $\{\emptyset, Z_4, Z_1\}$
(see diagram 3 of the **Figure 2**);

- 4) $\{\emptyset, Z_4, Z_1, \bar{D}\}, \{\emptyset, Z_5, Z_1, \bar{D}\}, \{\emptyset, Z_6, Z_1, \bar{D}\}, \{\emptyset, Z_6, Z_2, \bar{D}\}, \{\emptyset, Z_6, Z_3, \bar{D}\},$
 $\{\emptyset, Z_7, Z_3, \bar{D}\}, \{\emptyset, Z_8, Z_3, \bar{D}\}$
 (see diagram 4 of the **Figure 2**);
- 5) $\{\emptyset, Z_5, Z_4, Z_1\}, \{\emptyset, Z_6, Z_4, Z_1\}, \{\emptyset, Z_6, Z_5, Z_1\}, \{\emptyset, Z_7, Z_6, Z_3\}, \{\emptyset, Z_8, Z_6, Z_3\}, \{\emptyset, Z_8, Z_7, Z_3\},$
 $\{\emptyset, Z_8, Z_4, \bar{D}\}, \{\emptyset, Z_8, Z_5, \bar{D}\}, \{\emptyset, Z_7, Z_2, \bar{D}\}, \{\emptyset, Z_7, Z_4, \bar{D}\}, \{\emptyset, Z_7, Z_5, \bar{D}\}, \{\emptyset, Z_8, Z_1, \bar{D}\},$
 $\{\emptyset, Z_8, Z_2, \bar{D}\}, \{\emptyset, Z_4, Z_2, \bar{D}\}, \{\emptyset, Z_4, Z_3, \bar{D}\}, \{\emptyset, Z_5, Z_2, \bar{D}\}, \{\emptyset, Z_5, Z_3, \bar{D}\}, \{\emptyset, Z_7, Z_1, \bar{D}\},$
 $\{\emptyset, Z_2, Z_1, \bar{D}\}, \{\emptyset, Z_3, Z_1, \bar{D}\}, \{\emptyset, Z_3, Z_2, \bar{D}\}$
 (see diagram 5 of the **Figure 2**);
- 6) $\{\emptyset, Z_5, Z_4, Z_1, \bar{D}\}, \{\emptyset, Z_6, Z_4, Z_1, \bar{D}\}, \{\emptyset, Z_6, Z_5, Z_1, \bar{D}\}, \{\emptyset, Z_7, Z_6, Z_3, \bar{D}\},$
 $\{\emptyset, Z_8, Z_6, Z_3, \bar{D}\}, \{\emptyset, Z_8, Z_7, Z_3, \bar{D}\}$
 (see diagram 6 of the **Figure 2**);
- 7) $\{\emptyset, Z_6, Z_2, Z_1, \bar{D}\}, \{\emptyset, Z_6, Z_3, Z_1, \bar{D}\}, \{\emptyset, Z_6, Z_3, Z_2, \bar{D}\}$
 (see diagram 7 of the **Figure 2**);
- 8) $\{\emptyset, Z_8, Z_6, Z_3, Z_2, \bar{D}\}, \{\emptyset, Z_8, Z_6, Z_3, Z_1, \bar{D}\}, \{\emptyset, Z_7, Z_6, Z_3, Z_2, \bar{D}\}, \{\emptyset, Z_7, Z_6, Z_3, Z_1, \bar{D}\},$
 $\{\emptyset, Z_6, Z_5, Z_3, Z_1, \bar{D}\}, \{\emptyset, Z_6, Z_5, Z_2, Z_1, \bar{D}\}, \{\emptyset, Z_6, Z_4, Z_3, Z_1, \bar{D}\}, \{\emptyset, Z_6, Z_4, Z_2, Z_1, \bar{D}\}$
 (see diagram 8 of the **Figure 2**);

Theorem 13. Let $D \in \Sigma_1(X, 10)$, $Z_9 = \emptyset$ and $\alpha \in B_X(D)$. Binary relation α is an idempotent relation of the semigroup $B_X(D)$ iff binary relation α satisfies only one conditions of the following conditions:

- a) $\alpha = \emptyset$;
- b) $\alpha = (Y_9^\alpha \times \emptyset) \cup (Y_T^\alpha \times T)$, where $T \in D$, $\emptyset \neq T$, $Y_T^\alpha \neq \emptyset$, and satisfies the conditions: $Y_T^\alpha \cap T \neq \emptyset$;
- c) $\alpha = (Y_9^\alpha \times \emptyset) \cup (Y_T^\alpha \times T) \cup (Y_{T'}^\alpha \times T')$, where $T, T' \in D$, $\emptyset \neq T \subset T'$, $Y_T^\alpha, Y_{T'}^\alpha \notin \{\emptyset\}$, and satisfies the conditions: $Y_9^\alpha \cup Y_T^\alpha \supseteq T$, $Y_T^\alpha \cap T \neq \emptyset$, $Y_{T'}^\alpha \cap T' \neq \emptyset$;
- d) $\alpha = (Y_9^\alpha \times \emptyset) \cup (Y_T^\alpha \times T) \cup (Y_{T'}^\alpha \times T') \cup (Y_0^\alpha \times \bar{D})$, where $T, T' \in D$, $\emptyset \neq T \subset T' \subset \bar{D}$, $Y_T^\alpha, Y_{T'}^\alpha, Y_0^\alpha \notin \{\emptyset\}$, and satisfies the conditions: $Y_9^\alpha \cup Y_T^\alpha \supseteq T$, $Y_9^\alpha \cup Y_T^\alpha \cup Y_{T'}^\alpha \supseteq T'$, $Y_T^\alpha \cap T \neq \emptyset$, $Y_{T'}^\alpha \cap T' \neq \emptyset$, $Y_0^\alpha \cap \bar{D} \neq \emptyset$;
- e) $\alpha = (Y_9^\alpha \times \emptyset) \cup (Y_T^\alpha \times T) \cup (Y_{T'}^\alpha \times T') \cup (Y_{T \cup T'}^\alpha \times (T \cup T'))$, where $T, T' \in D$, $T \setminus T' \neq \emptyset$, $T' \setminus T \neq \emptyset$, $Y_T^\alpha, Y_{T'}^\alpha \notin \{\emptyset\}$ and satisfies the conditions: $Y_9^\alpha \cup Y_T^\alpha \supseteq T$, $Y_9^\alpha \cup Y_{T'}^\alpha \supseteq T'$, $Y_T^\alpha \cap T \neq \emptyset$, $Y_{T'}^\alpha \cap T' \neq \emptyset$;
- f) $\alpha = (Y_9^\alpha \times \emptyset) \cup (Y_T^\alpha \times T) \cup (Y_{T'}^\alpha \times T') \cup (Y_{T \cup T'}^\alpha \times (T \cup T')) \cup (Y_0^\alpha \times \bar{D})$, where $T \setminus T' \neq \emptyset$, $T' \setminus T \neq \emptyset$, $Y_T^\alpha, Y_{T'}^\alpha, Y_0^\alpha \notin \{\emptyset\}$ and satisfies the conditions: $Y_9^\alpha \cup Y_T^\alpha \supseteq T$, $Y_9^\alpha \cup Y_{T'}^\alpha \supseteq T'$, $Y_T^\alpha \cap T \neq \emptyset$, $Y_{T'}^\alpha \cap T' \neq \emptyset$, $Y_0^\alpha \cap \bar{D} \neq \emptyset$;
- g) $\alpha = (Y_9^\alpha \times \emptyset) \cup (Y_6^\alpha \times Z_6) \cup (Y_T^\alpha \times T) \cup (Y_{T'}^\alpha \times T') \cup (Y_0^\alpha \times \bar{D})$, where $T, T' \in D$, $Z_6 \subset T$, $Z_6 \subset T'$, $T \setminus T' \neq \emptyset$, $T' \setminus T \neq \emptyset$, $T \cup T' = \bar{D}$, $Y_6^\alpha, Y_T^\alpha, Y_{T'}^\alpha \notin \{\emptyset\}$ and satisfies the conditions: $Y_9^\alpha \cup Y_6^\alpha \supseteq Z_6$, $Y_9^\alpha \cup Y_6^\alpha \cup Y_T^\alpha \supseteq T$, $Y_9^\alpha \cup Y_6^\alpha \cup Y_{T'}^\alpha \supseteq T'$, $Y_6^\alpha \cap Z_6 \neq \emptyset$, $Y_T^\alpha \cap T \neq \emptyset$, $Y_{T'}^\alpha \cap T' \neq \emptyset$;
- h) $\alpha = (Y_9^\alpha \times \emptyset) \cup (Y_T^\alpha \times T) \cup (Y_{T'}^\alpha \times T') \cup (Y_{T \cup T'}^\alpha \times (T \cup T')) \cup (Y_Z^\alpha \times Z) \cup (Y_0^\alpha \times \bar{D})$, where $T' \subset Z$, $T \setminus T' \neq \emptyset$, $T' \setminus T \neq \emptyset$, $(T \cup T') \setminus Z \neq \emptyset$, $Z \setminus (T \cup T') \neq \emptyset$, $Y_T^\alpha, Y_{T'}^\alpha, Y_Z^\alpha \notin \{\emptyset\}$ and satisfies the conditions: $Y_9^\alpha \cup Y_T^\alpha \supseteq T$, $Y_9^\alpha \cup Y_{T'}^\alpha \supseteq T'$, $Y_9^\alpha \cup Y_{T'}^\alpha \cup Y_Z^\alpha \supseteq Z$, $Y_T^\alpha \cap T \neq \emptyset$, $Y_{T'}^\alpha \cap T' \neq \emptyset$, $Y_Z^\alpha \cap Z \neq \emptyset$;

Lemma 14. Let $D \in \Sigma_1(X, 10)$ and $Z_9 = \emptyset$. If X is a finite set, then $|I^*(Q_1)| = 1$.

Lemma 15. Let $D \in \Sigma_1(X, 10)$ and $Z_9 = \emptyset$. If X is a finite set, then the number $|I^*(Q_2)|$ may be calcu-

lated by formula

$$\begin{aligned} |I^*(Q_2)| = & \left(2^{|\bar{D}|} - 1\right) \cdot 2^{|X \setminus \bar{D}|} + \left(2^{|\bar{Z}_8|} - 1\right) \cdot 2^{|X \setminus Z_8|} + \left(2^{|\bar{Z}_7|} - 1\right) \cdot 2^{|X \setminus Z_7|} + \left(2^{|\bar{Z}_6|} - 1\right) \cdot 2^{|X \setminus Z_6|} \\ & + \left(2^{|\bar{Z}_5|} - 1\right) \cdot 2^{|X \setminus Z_5|} + \left(2^{|\bar{Z}_4|} - 1\right) \cdot 2^{|X \setminus Z_4|} + \left(2^{|\bar{Z}_3|} - 1\right) \cdot 2^{|X \setminus Z_3|} \\ & + \left(2^{|\bar{Z}_2|} - 1\right) \cdot 2^{|X \setminus Z_2|} + \left(2^{|\bar{Z}_1|} - 1\right) \cdot 2^{|X \setminus Z_1|}. \end{aligned}$$

Lemma 16. Let $D \in \Sigma_1(X, 10)$ and $Z_9 = \emptyset$. If X is a finite set, then the number $|I^*(Q_3)|$ may be calculated by formula

$$\begin{aligned} |I^*(Q_3)| = & \left(2^{|\bar{Z}_8|} - 1\right) \cdot \left(3^{|\bar{D} \setminus Z_8|} - 2^{|\bar{D} \setminus Z_8|}\right) \cdot 3^{|X \setminus \bar{D}|} + \left(2^{|\bar{Z}_7|} - 1\right) \cdot \left(3^{|\bar{D} \setminus Z_7|} - 2^{|\bar{D} \setminus Z_7|}\right) \cdot 3^{|X \setminus \bar{D}|} \\ & + \left(2^{|\bar{Z}_6|} - 1\right) \cdot \left(3^{|\bar{D} \setminus Z_6|} - 2^{|\bar{D} \setminus Z_6|}\right) \cdot 3^{|X \setminus \bar{D}|} + \left(2^{|\bar{Z}_5|} - 1\right) \cdot \left(3^{|\bar{D} \setminus Z_5|} - 2^{|\bar{D} \setminus Z_5|}\right) \cdot 3^{|X \setminus \bar{D}|} \\ & + \left(2^{|\bar{Z}_4|} - 1\right) \cdot \left(3^{|\bar{D} \setminus Z_4|} - 2^{|\bar{D} \setminus Z_4|}\right) \cdot 3^{|X \setminus \bar{D}|} + \left(2^{|\bar{Z}_3|} - 1\right) \cdot \left(3^{|\bar{D} \setminus Z_3|} - 2^{|\bar{D} \setminus Z_3|}\right) \cdot 3^{|X \setminus \bar{D}|} \\ & + \left(2^{|\bar{Z}_2|} - 1\right) \cdot \left(3^{|\bar{D} \setminus Z_2|} - 2^{|\bar{D} \setminus Z_2|}\right) \cdot 3^{|X \setminus \bar{D}|} + \left(2^{|\bar{Z}_1|} - 1\right) \cdot \left(3^{|\bar{D} \setminus Z_1|} - 2^{|\bar{D} \setminus Z_1|}\right) \cdot 3^{|X \setminus \bar{D}|} \\ & + \left(2^{|\bar{Z}_8|} - 1\right) \cdot \left(3^{|\bar{Z}_3 \setminus Z_8|} - 2^{|\bar{Z}_3 \setminus Z_8|}\right) \cdot 3^{|X \setminus Z_3|} + \left(2^{|\bar{Z}_7|} - 1\right) \cdot \left(3^{|\bar{Z}_3 \setminus Z_7|} - 2^{|\bar{Z}_3 \setminus Z_7|}\right) \cdot 3^{|X \setminus Z_3|} \\ & + \left(2^{|\bar{Z}_6|} - 1\right) \cdot \left(3^{|\bar{Z}_3 \setminus Z_6|} - 2^{|\bar{Z}_3 \setminus Z_6|}\right) \cdot 3^{|X \setminus Z_3|} + \left(2^{|\bar{Z}_6|} - 1\right) \cdot \left(3^{|\bar{Z}_2 \setminus Z_6|} - 2^{|\bar{Z}_2 \setminus Z_6|}\right) \cdot 3^{|X \setminus Z_2|} \\ & + \left(2^{|\bar{Z}_6|} - 1\right) \cdot \left(3^{|\bar{Z}_1 \setminus Z_6|} - 2^{|\bar{Z}_1 \setminus Z_6|}\right) \cdot 3^{|X \setminus Z_1|} + \left(2^{|\bar{Z}_5|} - 1\right) \cdot \left(3^{|\bar{Z}_1 \setminus Z_5|} - 2^{|\bar{Z}_1 \setminus Z_5|}\right) \cdot 3^{|X \setminus Z_1|} \\ & + \left(2^{|\bar{Z}_4|} - 1\right) \cdot \left(3^{|\bar{Z}_1 \setminus Z_4|} - 2^{|\bar{Z}_1 \setminus Z_4|}\right) \cdot 3^{|X \setminus Z_1|}. \end{aligned}$$

Lemma 17. Let $D \in \Sigma_1(X, 10)$ and $Z_9 = \emptyset$. If X is a finite set, then the number $|I^*(Q_4)|$ may be calculated by formula

$$\begin{aligned} |I^*(Q_4)| = & \left(2^{|\bar{Z}_4|} - 1\right) \cdot \left(3^{|\bar{Z}_1 \setminus Z_4|} - 2^{|\bar{Z}_1 \setminus Z_4|}\right) \cdot \left(4^{|\bar{D} \setminus Z_1|} - 3^{|\bar{D} \setminus Z_1|}\right) \cdot 4^{|X \setminus \bar{D}|} \\ & + \left(2^{|\bar{Z}_5|} - 1\right) \cdot \left(3^{|\bar{Z}_1 \setminus Z_5|} - 2^{|\bar{Z}_1 \setminus Z_5|}\right) \cdot \left(4^{|\bar{D} \setminus Z_1|} - 3^{|\bar{D} \setminus Z_1|}\right) \cdot 4^{|X \setminus \bar{D}|} \\ & + \left(2^{|\bar{Z}_6|} - 1\right) \cdot \left(3^{|\bar{Z}_1 \setminus Z_6|} - 2^{|\bar{Z}_1 \setminus Z_6|}\right) \cdot \left(4^{|\bar{D} \setminus Z_1|} - 3^{|\bar{D} \setminus Z_1|}\right) \cdot 4^{|X \setminus \bar{D}|} \\ & + \left(2^{|\bar{Z}_6|} - 1\right) \cdot \left(3^{|\bar{Z}_2 \setminus Z_6|} - 2^{|\bar{Z}_2 \setminus Z_6|}\right) \cdot \left(4^{|\bar{D} \setminus Z_2|} - 3^{|\bar{D} \setminus Z_2|}\right) \cdot 4^{|X \setminus \bar{D}|} \\ & + \left(2^{|\bar{Z}_6|} - 1\right) \cdot \left(3^{|\bar{Z}_3 \setminus Z_6|} - 2^{|\bar{Z}_3 \setminus Z_6|}\right) \cdot \left(4^{|\bar{D} \setminus Z_3|} - 3^{|\bar{D} \setminus Z_3|}\right) \cdot 4^{|X \setminus \bar{D}|} \\ & + \left(2^{|\bar{Z}_7|} - 1\right) \cdot \left(3^{|\bar{Z}_3 \setminus Z_7|} - 2^{|\bar{Z}_3 \setminus Z_7|}\right) \cdot \left(4^{|\bar{D} \setminus Z_3|} - 3^{|\bar{D} \setminus Z_3|}\right) \cdot 4^{|X \setminus \bar{D}|} \\ & + \left(2^{|\bar{Z}_8|} - 1\right) \cdot \left(3^{|\bar{Z}_3 \setminus Z_8|} - 2^{|\bar{Z}_3 \setminus Z_8|}\right) \cdot \left(4^{|\bar{D} \setminus Z_3|} - 3^{|\bar{D} \setminus Z_3|}\right) \cdot 4^{|X \setminus \bar{D}|}. \end{aligned}$$

Lemma 18. Let $D \in \Sigma_1(X, 10)$ and $Z_9 = \emptyset$. If X is a finite set, then the number $|I^*(Q_5)|$ may be calculated by formula

lated by formula

$$\begin{aligned}
|I^*(Q_5)| = & (2^{|Z_5 \setminus Z_4|} - 1) \cdot (2^{|Z_4 \setminus Z_5|} - 1) \cdot 4^{|X \setminus Z_1|} + (2^{|Z_6 \setminus Z_4|} - 1) \cdot (2^{|Z_4 \setminus Z_6|} - 1) \cdot 4^{|X \setminus Z_1|} \\
& + (2^{|Z_6 \setminus Z_5|} - 1) \cdot (2^{|Z_5 \setminus Z_6|} - 1) \cdot 4^{|X \setminus Z_1|} + (2^{|Z_7 \setminus Z_6|} - 1) \cdot (2^{|Z_6 \setminus Z_7|} - 1) \cdot 4^{|X \setminus Z_3|} \\
& + (2^{|Z_8 \setminus Z_6|} - 1) \cdot (2^{|Z_6 \setminus Z_8|} - 1) \cdot 4^{|X \setminus Z_3|} + (2^{|Z_8 \setminus Z_7|} - 1) \cdot (2^{|Z_7 \setminus Z_8|} - 1) \cdot 4^{|X \setminus Z_3|} \\
& + (2^{|Z_8 \setminus Z_4|} - 1) \cdot (2^{|Z_4 \setminus Z_8|} - 1) \cdot 4^{|X \setminus \bar{D}|} + (2^{|Z_8 \setminus Z_5|} - 1) \cdot (2^{|Z_5 \setminus Z_8|} - 1) \cdot 4^{|X \setminus \bar{D}|} \\
& + (2^{|Z_7 \setminus Z_2|} - 1) \cdot (2^{|Z_2 \setminus Z_7|} - 1) \cdot 4^{|X \setminus \bar{D}|} + (2^{|Z_7 \setminus Z_4|} - 1) \cdot (2^{|Z_4 \setminus Z_7|} - 1) \cdot 4^{|X \setminus \bar{D}|} \\
& + (2^{|Z_7 \setminus Z_5|} - 1) \cdot (2^{|Z_5 \setminus Z_7|} - 1) \cdot 4^{|X \setminus \bar{D}|} + (2^{|Z_8 \setminus Z_1|} - 1) \cdot (2^{|Z_1 \setminus Z_8|} - 1) \cdot 4^{|X \setminus \bar{D}|} \\
& + (2^{|Z_8 \setminus Z_2|} - 1) \cdot (2^{|Z_2 \setminus Z_8|} - 1) \cdot 4^{|X \setminus \bar{D}|} + (2^{|Z_4 \setminus Z_2|} - 1) \cdot (2^{|Z_2 \setminus Z_4|} - 1) \cdot 4^{|X \setminus \bar{D}|} \\
& + (2^{|Z_4 \setminus Z_3|} - 1) \cdot (2^{|Z_3 \setminus Z_4|} - 1) \cdot 4^{|X \setminus \bar{D}|} + (2^{|Z_5 \setminus Z_2|} - 1) \cdot (2^{|Z_2 \setminus Z_5|} - 1) \cdot 4^{|X \setminus \bar{D}|} \\
& + (2^{|Z_5 \setminus Z_3|} - 1) \cdot (2^{|Z_3 \setminus Z_5|} - 1) \cdot 4^{|X \setminus \bar{D}|} + (2^{|Z_7 \setminus Z_1|} - 1) \cdot (2^{|Z_1 \setminus Z_7|} - 1) \cdot 4^{|X \setminus \bar{D}|} \\
& + (2^{|Z_2 \setminus Z_1|} - 1) \cdot (2^{|Z_1 \setminus Z_2|} - 1) \cdot 4^{|X \setminus \bar{D}|} + (2^{|Z_3 \setminus Z_1|} - 1) \cdot (2^{|Z_1 \setminus Z_3|} - 1) \cdot 4^{|X \setminus \bar{D}|} \\
& + (2^{|Z_3 \setminus Z_2|} - 1) \cdot (2^{|Z_2 \setminus Z_3|} - 1) \cdot 4^{|X \setminus \bar{D}|}.
\end{aligned}$$

Lemma 19. Let $D \in \Sigma_1(X, 10)$ and $Z_9 = \emptyset$. If X is a finite set, then the number $|I^*(Q_6)|$ may be calculated by formula

$$\begin{aligned}
|I^*(Q_6)| = & (2^{|Z_5 \setminus Z_4|} - 1) \cdot (2^{|Z_4 \setminus Z_5|} - 1) \cdot (5^{|D \setminus Z_1|} - 4^{|D \setminus Z_1|}) \cdot 5^{|X \setminus \bar{D}|} \\
& + (2^{|Z_6 \setminus Z_4|} - 1) \cdot (2^{|Z_4 \setminus Z_6|} - 1) \cdot (5^{|D \setminus Z_1|} - 4^{|D \setminus Z_1|}) \cdot 5^{|X \setminus \bar{D}|} \\
& + (2^{|Z_6 \setminus Z_5|} - 1) \cdot (2^{|Z_5 \setminus Z_6|} - 1) \cdot (5^{|D \setminus Z_1|} - 4^{|D \setminus Z_1|}) \cdot 5^{|X \setminus \bar{D}|} \\
& + (2^{|Z_7 \setminus Z_6|} - 1) \cdot (2^{|Z_6 \setminus Z_7|} - 1) \cdot (5^{|D \setminus Z_3|} - 4^{|D \setminus Z_3|}) \cdot 5^{|X \setminus \bar{D}|} \\
& + (2^{|Z_8 \setminus Z_6|} - 1) \cdot (2^{|Z_6 \setminus Z_8|} - 1) \cdot (5^{|D \setminus Z_3|} - 4^{|D \setminus Z_3|}) \cdot 5^{|X \setminus \bar{D}|} \\
& + (2^{|Z_8 \setminus Z_7|} - 1) \cdot (2^{|Z_7 \setminus Z_8|} - 1) \cdot (5^{|D \setminus Z_3|} - 4^{|D \setminus Z_3|}) \cdot 5^{|X \setminus \bar{D}|}.
\end{aligned}$$

Lemma 20. Let $D \in \Sigma_1(X, 10)$ and $Z_9 = \emptyset$. If X is a finite set, then the number $|I^*(Q_7)|$ may be calculated by formula

$$\begin{aligned}
|I^*(Q_7)| = & (2^{|Z_6|} - 1) \cdot 2^{(|Z_2 \cap Z_1|) \setminus Z_6} \cdot (3^{|Z_2 \setminus Z_1|} - 2^{|Z_2 \setminus Z_1|}) \cdot (3^{|Z_1 \setminus Z_2|} - 2^{|Z_1 \setminus Z_2|}) \cdot 5^{|X \setminus \bar{D}|} \\
& + (2^{|Z_6|} - 1) \cdot 2^{(|Z_3 \cap Z_1|) \setminus Z_6} \cdot (3^{|Z_3 \setminus Z_1|} - 2^{|Z_3 \setminus Z_1|}) \cdot (3^{|Z_1 \setminus Z_3|} - 2^{|Z_1 \setminus Z_3|}) \cdot 5^{|X \setminus \bar{D}|} \\
& + (2^{|Z_6|} - 1) \cdot 2^{(|Z_3 \cap Z_2|) \setminus Z_6} \cdot (3^{|Z_3 \setminus Z_2|} - 2^{|Z_3 \setminus Z_2|}) \cdot (3^{|Z_2 \setminus Z_3|} - 2^{|Z_2 \setminus Z_3|}) \cdot 5^{|X \setminus \bar{D}|}.
\end{aligned}$$

Lemma 21. Let $D \in \Sigma_1(X, 7)$ and $Z_9 = \emptyset$. If X is a finite set, then the number $|I^*(Q_8)|$ may be calculated by formula

lated by formula

$$\begin{aligned}
 |I^*(Q_8)| = & (2^{|Z_8 \setminus Z_2|} - 1) \cdot (2^{|Z_6 \setminus Z_8|} - 1) \cdot (3^{|Z_2 \setminus Z_3|} - 2^{|Z_2 \setminus Z_3|}) \cdot 6^{|X \setminus \bar{D}|} \\
 & + (2^{|Z_8 \setminus Z_1|} - 1) \cdot (2^{|Z_6 \setminus Z_8|} - 1) \cdot (3^{|Z_1 \setminus Z_3|} - 2^{|Z_1 \setminus Z_3|}) \cdot 6^{|X \setminus \bar{D}|} \\
 & + (2^{|Z_7 \setminus Z_2|} - 1) \cdot (2^{|Z_6 \setminus Z_7|} - 1) \cdot (3^{|Z_2 \setminus Z_3|} - 2^{|Z_2 \setminus Z_3|}) \cdot 6^{|X \setminus \bar{D}|} \\
 & + (2^{|Z_7 \setminus Z_1|} - 1) \cdot (2^{|Z_6 \setminus Z_7|} - 1) \cdot (3^{|Z_1 \setminus Z_3|} - 2^{|Z_1 \setminus Z_3|}) \cdot 6^{|X \setminus \bar{D}|} \\
 & + (2^{|Z_5 \setminus Z_3|} - 1) \cdot (2^{|Z_6 \setminus Z_5|} - 1) \cdot (3^{|Z_3 \setminus Z_1|} - 2^{|Z_3 \setminus Z_1|}) \cdot 6^{|X \setminus \bar{D}|} \\
 & + (2^{|Z_5 \setminus Z_2|} - 1) \cdot (2^{|Z_6 \setminus Z_5|} - 1) \cdot (3^{|Z_2 \setminus Z_1|} - 2^{|Z_2 \setminus Z_1|}) \cdot 6^{|X \setminus \bar{D}|} \\
 & + (2^{|Z_4 \setminus Z_3|} - 1) \cdot (2^{|Z_6 \setminus Z_4|} - 1) \cdot (3^{|Z_3 \setminus Z_1|} - 2^{|Z_3 \setminus Z_1|}) \cdot 6^{|X \setminus \bar{D}|} \\
 & + (2^{|Z_4 \setminus Z_2|} - 1) \cdot (2^{|Z_6 \setminus Z_4|} - 1) \cdot (3^{|Z_2 \setminus Z_1|} - 2^{|Z_2 \setminus Z_1|}) \cdot 6^{|X \setminus \bar{D}|}.
 \end{aligned}$$

Theorem 14. Let $D \in \Sigma_1(X, 10)$, $Z_9 = \emptyset$. If X is a finite set and I_D is a set of all idempotent elements of the semigroup $B_X(D)$, then $|I_D| = \sum_{i=1}^8 |I^*(Q_i)|$.

Example 15. Let $X = \{1, 2, 3, 4, 5, 6, 7\}$,

$$P_1 = \{1\}, P_2 = \{2\}, P_3 = \{3\}, P_4 = \{4\}, P_5 = \{5\}, P_7 = \{6\}, P_8 = \{7\}, P_0 = P_6 = P_9 = \emptyset.$$

Then $\bar{D} = \{1, 2, 3, 4, 5, 6, 7\}$, $Z_1 = \{2, 3, 4, 5, 6, 7\}$, $Z_2 = \{1, 3, 4, 5, 6, 7\}$, $Z_3 = \{1, 2, 4, 5, 6, 7\}$,
 $Z_4 = \{2, 3, 5, 6, 7\}$, $Z_5 = \{2, 3, 4, 6, 7\}$, $Z_6 = \{4, 5, 6, 7\}$, $Z_7 = \{1, 2, 4, 5, 7\}$, $Z_8 = \{1, 2, 4, 5, 6\}$ and $Z_9 = \emptyset$.

$$\begin{aligned}
 D = & \{\{1, 2, 3, 4, 5, 6, 7\}, \{2, 3, 4, 5, 6, 7\}, \{1, 3, 4, 5, 6, 7\}, \{1, 2, 4, 5, 6, 7\}, \{2, 3, 5, 6, 7\}, \\
 & \{2, 3, 4, 6, 7\}, \{4, 5, 6, 7\}, \{1, 2, 4, 5, 7\}, \{1, 2, 4, 5, 6\}, \emptyset\}
 \end{aligned}$$

We have $Z_9 = \emptyset$. Where $|I^*(Q_1)| = 1$, $|I^*(Q_2)| = 1121$, $|I^*(Q_3)| = 2141$, $|I^*(Q_4)| = 349$, $|I^*(Q_5)| = 119$,
 $|I^*(Q_6)| = 14$, $|I^*(Q_7)| = 90$, $|I^*(Q_8)| = 8$, $|I_D| = 3843$.

It was seen in ([4], Theorem 2) that if α and β are regular elements of $B_X(D)$ then $V(D, \alpha \circ \beta)$ is an XI -subsemilattice of D . Therefore $\alpha \circ \beta$ is regular elements of $B_X(D)$. That is the set of all regular elements of $B_X(D)$ is a subsemigroup of $B_X(D)$.

References

- [1] Diasamidze, Ya. and Makharadze, Sh. (2013) Complete Semigroups of Binary Relations. Monograph. Kriter, Turkey, 620 p.
- [2] Diasamidze, Ya. and Makharadze, Sh. (2010) Complete Semigroups of Binary Relations. Monograph. M., Sputnik+, 657 p. (In Russian)
- [3] Diasamidze, Ya., Makharadze, Sh. and Diasamidze, Il. (2008) Idempotents and Regular Elements of Complete Semigroups of Binary Relations. *Journal of Mathematical Sciences, Plenum Publ. Cor., New York*, **153**, 481-499.
- [4] Diasamidze, Ya. and Bakuridze, Al. (to appear) On Some Properties of Regular Elements of Complete Semigroups Defined by Semilattices of the Class $\Sigma_4(X, 8)$.

Scientific Research Publishing (SCIRP) is one of the largest Open Access journal publishers. It is currently publishing more than 200 open access, online, peer-reviewed journals covering a wide range of academic disciplines. SCIRP serves the worldwide academic communities and contributes to the progress and application of science with its publication.

Other selected journals from SCIRP are listed as below. Submit your manuscript to us via either submit@scirp.org or **Online Submission Portal**.

