

# Assessment of the Heavy Metals and Natural Radioactivity in Phosphate Mines and Occupational Health Effects at Some Egyptian Regions

E. R. Atta, Kh. M. Zakaria, M. S. Ibrahim

Nuclear and Radiological Regulatory Authority, Cairo, Egypt

Email: drkhaledzakaria@gmail.com

**How to cite this paper:** RAtta, E.R., Zaka-  
ria, Kh.M. and Ibrahim, M.S. (2016) As-  
sessment of the Heavy Metals and Natural  
Radioactivity in Phosphate Mines and Oc-  
cupational Health Effects at Some Egyptian  
Regions. *Journal of Environmental Protec-  
tion*, 7, 1657-1669.

<http://dx.doi.org/10.4236/jep.2016.711135>

**Received:** August 8, 2016

**Accepted:** October 28, 2016

**Published:** October 31, 2016

Copyright © 2016 by authors and  
Scientific Research Publishing Inc.

This work is licensed under the Creative  
Commons Attribution International  
License (CC BY 4.0).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

---

## Abstract

In this work, the specific activities of natural radionuclides, namely  $^{238}\text{U}$  series,  $^{232}\text{Th}$  series and  $^{40}\text{K}$ , have been measured in collected sedimentary phosphate deposits samples from El-Hamraween, El-Quser and Safaga phosphate mines in Egypt. HPGe  $\gamma$ -spectrometry was used. This study was undertaken to estimate the radiation hazard indices in phosphate mining at the studied mines on their occupational workers, to establish correlation relationships between the some measured heavy metals such as As, Cd and Pb in blood workers and their concentration in phosphate rock ores and to determinate the biomarkers in the blood workers such as malondialdehyde (MDA) and superoxide dismutase (SOD). The phosphate mine sample of El-Hamrawein has the lowest activity concentration of  $^{226}\text{Ra}$ ,  $^{232}\text{Th}$  and  $^{40}\text{K}$  in comparison to the phosphate mine samples of El-Quseir and Safaga (El-Hamrawein < El-Quseir < Safaga). The activity concentrations for the radionuclide's considerably were fluctuated; for  $^{226}\text{Ra}$  activity concentrations varied from 222.4 to 255.8  $\text{Bq}\cdot\text{kg}^{-1}$ , 122.4 to 188.3  $\text{Bq}\cdot\text{kg}^{-1}$  and 115.4 to 165.8  $\text{Bq}\cdot\text{kg}^{-1}$  for Safaga, El-Quseir and El-Hamrawein, respectively. For  $^{232}\text{Th}$  activity concentrations varied from 135.6 to 212.3  $\text{Bq}\cdot\text{kg}^{-1}$ , 112.8 to 167.4  $\text{Bq}\cdot\text{kg}^{-1}$  and 132.8 to 188.6  $\text{Bq}\cdot\text{kg}^{-1}$  for Safaga, El-Quseir and El-Hamrawein, respectively. For  $^{40}\text{K}$  activity concentrations varied from 225.2 to 312.8  $\text{Bq}\cdot\text{kg}^{-1}$ , 168.7 to 268.9  $\text{Bq}\cdot\text{kg}^{-1}$  and 95.2 to 155.8  $\text{Bq}\cdot\text{kg}^{-1}$  for Safaga, El-Quseir and El-Hamrawein, respectively. The workers of old ages have higher concentration of the investigated heavy metals than those of young ages. There is a good relation between the concentration of the investigated metals in phosphate mine samples and their concentration in blood of the occupational workers in these mines.

---

## Keywords

Superoxide Dismutase, Malondialdehyde, Natural Radionuclides, Heavy Metals, Phosphate, Occupational Workers, Health Effects

---

## 1. Introduction

Studies of natural environmental radiation and radioactivity are of great importance and interest for environmental chemistry as well as many other disciplines. Environmental radiation originates from a number of naturally occurring and human-made sources. The estimation of exposure to ionizing radiation is an important goal of regulatory authorities and radiation protection scientists. Thus knowledge of the background radiation level is of paramount importance [1] [2]. Radioactive materials occur naturally everywhere in the environment (e.g., uranium series, thorium series and potassium-40). By far the largest proportion of human exposure to radiation comes from natural sources—from external sources of radiation, including cosmic and terrestrial radiation, and from inhalation or ingestion of radioactive materials. The global average annual human exposure from natural sources is 2.4 mSv/year. Some sources (e.g., uranium) can be concentrated during extraction by mining and other industrial activities [3]. The phosphate rock extractions are man-made sources of air- and water-borne radionuclide releases to the environment. Phosphate rocks contain relatively high concentrations of naturally occurring radioactive materials from uranium series and thorium series ( $^{238}\text{U}$  and  $^{232}\text{Th}$ ). Mining, milling, transportation of phosphate ores, manufacturing of phosphate fertilizers containing uranium are ways in which the workers, public and the environment are exposed to enhance natural radioactivity [4]. Man is exposed to ionizing radiation with or without his consent. Because of the lethal effects of ionizing radiation, the practice has been to monitor and assess the levels of exposure and keep one's exposure to ionizing radiation as low as reasonably achievable (also known as the ALARA principle). In Egypt, phosphate formations are established in different sites. Three locations of phosphate mines have been selected for this study and they are Safaga (33°57'E 26°44'N), El-Quseir (34°17'E 26°06'N) and El-Hamrawein (35°12'E 26°15'N). The terrestrial gamma radionuclides cause the major contribution to the annual average doses of radiation exposure of the world's population [3]. Natural environmental radioactivity series mainly from primordial radionuclides, such as the nuclides from both  $^{238}\text{U}$  and  $^{232}\text{Th}$  series and their decay products as well as  $^{40}\text{K}$  occur at trace levels in all ground formations [5]. Exposure of workers and the public to radiation from phosphate rock is therefore unlikely. The European Commission has issued a draft proposal for revision of the Basic Safety Standards for the protection of workers and the general public against the dangers of ionizing radiation [6]. Mined phosphate poses concerns due to levels of heavy metals that may present in phosphate rock. Trace heavy metals constitute the main oxidative stress factor in mammals by the production of free radicals which cause the cardiovascular diseases [7]. The heavy metals oxidize li-

pids and proteins to form deterioration products with high radical activity which produces Reactive Oxygen Species (ROS) [8]. So, oxidative stress is an imbalance for the production of ROS which leads to an increased oxidation of cellular components [9].

This work aims to measure the specific activities of natural radionuclides, namely  $^{226}\text{Ra}$  ( $^{238}\text{U}$ ) series,  $^{232}\text{Th}$  series and  $^{40}\text{K}$ , in collected sedimentary phosphate deposits samples from El-Hamraween, El-Quser and Safaga phosphate mines in Egypt and to establish correlation relationships between some heavy metals such as As, Cd and Pb in blood workers and their concentration in phosphate rock ores and to determinate the biomarkers in blood of occupational workers such as malondialdehyde (MDA) and superoxide dismutase (SOD) in addition to the hematological parameters (Hemoglobin concentration, red blood count, white blood count and platelets count). This work can be considered as a step toward investigating natural radioactivity in phosphate hills in Egypt which is considered as the main part of radiological baseline map in Egypt. The experimental results will be used as guides for decision makers in solving some natural environmental problems which may be found in those areas of phosphate mines. On the other hand, the radiation safety measures and radiation safety standers should be taken into account for workers and environmental ecosystems in phosphate mines areas

## 2. Material and Methods

### 2.1. Sampling and Sample Preparation

In Egypt, phosphate formations are established in different sites. Three locations of open-pit phosphate mines have been selected for this study. These samples were collected by services partner (Baltic Control Company). El-Hamrawein ( $35^{\circ}12'E$   $26^{\circ}15'N$ ), El-Quseir ( $34^{\circ}17'E$   $26^{\circ}06'N$ ) and Safaga ( $33^{\circ}57'E$   $26^{\circ}44'N$ ) locations. Phosphate sampling was carried out in the months of April, May and December 2014. Twelve samples were collected from each mine; four samples were collected from the subsurface phosphate horizon (subsurface samples), four samples were collected from the exposed lower phosphate horizon (surface samples lower bed) and four samples were collected from the exposed upper phosphate horizon (surface samples upper bed). The collected samples were transferred to labeled polyethylene bags, closed and transferred to the laboratory for preparation and measurements. The collected samples were dried at room temperature for a week. The quartering technique was used to get a representative sample for each horizon. So, there are three representative samples for each mine. The Phosphate samples were prepared and analyzed as reported in [10].

### 2.2. Radioactivity Measurements

The activity concentration of the natural  $^{238}\text{U}$  series,  $^{232}\text{Th}$  series and  $^{40}\text{K}$  in the investigated samples were determined using a high-resolution HPGe  $\gamma$ -spectrometry system with 30% counting efficiency. These investigations were carried out in the Laboratory of Egyptian Nuclear and Radiological Regulatory Authority. This was performed by taking 250  $\text{cm}^3$  counting vials filled up to a height of 7 cm, which correspond to 170  $\text{cm}^3$ . The measurement duration was up to 80,000 sec. The obtained spectra were ana-

lyzed. The determination of the presence radionuclides and calculation of their activities were based on the following gamma-ray transitions (in keV): the  $^{226}\text{Ra}$  activities (or  $^{238}\text{U}$  activities for samples assumed to be in radioactive equilibrium) were estimated from  $^{234}\text{Th}$  (92.38 keV, 5.6%), while  $\gamma$ -energies of  $^{214}\text{Pb}$  (351.9 keV, 35.8%) and  $^{214}\text{Bi}$  (609.3, 45%), 1764.5 keV, 17% ) and  $^{226}\text{Ra}$  (185.99 KeV, 3.5% ) were used to estimate the concentration of  $^{226}\text{Ra}$ . The Gamma- ray energies of  $^{212}\text{Pb}$  (238.6 keV, 45%), and  $^{228}\text{Ac}$  (338.4 keV, 12.3%), (911.07 keV, 29%), (968.90 keV, 17 %) were used to estimate the concentration of  $^{232}\text{Th}$ . The activity concentrations of  $^{40}\text{K}$  were measured directly by its own gamma rays (1460.8 keV, 10.7%). In order to determine the background distribution due to naturally occurring radionuclides in the environment around the detector, an empty polystyrene container was counted in the same manner as the samples. The activity concentrations were calculated after measurement and subtraction of the background. The activities were determined from measuring their respective decay daughters. The activity concentrations were calculated from the intensity of each line taking into account the mass of the sample, the branching ratios of the  $\gamma$ -decay, the time of counting and the efficiencies of the detector [11] [12].

### 2.3. Determination of Heavy Metals in Phosphate Samples

Atomic absorption Spectrophotometer (AAS) is a simple and well available technique for the determinations of heavy metals in the soil samples. Heavy metals in sediments were determined according to [12]. These investigations were carried out in National Research Center, Cairo, Egypt. The metal ions were determined by Atomic Absorption Spectrophotometer, Perkin Elmer model Analyst 100 which is manufactured in USA. The sediments were digested with 5:1 mixture of HF and  $\text{HClO}_4$  acids; 1g (dry weight) sample was digested by 2 ml  $\text{HClO}_4$  and 10 ml HF to near dryness, subsequently a second addition of 1 ml of  $\text{HClO}_4$  and 10 ml of HF and evaporated to near dryness. Finally, 1 ml of  $\text{HClO}_4$  alone was added and the sample was evaporated until the appearance of white fumes. The residue was dissolved in 12N HCl and diluted to 25 ml with de-ionized water, the results obtained were determined according to [13].

### 2.4. Measurements of Heavy Metals in Blood Workers

Whole blood (1 ml) was transferred into screw capped polypropylene tubes and 5 ml nitric acid solution was added. After 10 minutes; it diluted with 5 ml deionized water, digested was continued in a microwave oven. In this digestion procedure, the temperature was increased gradually from 85°C to 230°C through 15 minutes. The sample preparation was completed with addition of 5 ml deionized water and 3 ml HCl 70% conc. Inductively-Coupled Plasma-Mass Spectroscopy (ICP-MS) (Agilent 7700X, USA) at (National Research Center, Cairo, Egypt) was used for the analysis of metals in collected samples. The standards for ICP-MS were prepared from stock solutions of lead, cadmium, arsenic at 10 mg/L concentrations obtained from Sigma-Aldrich, Australia, and labeled as Fluka Trace Cert Ultra Plasma with purity was higher than 99.999%. Calibration was performed by aqueous multi-element standard solutions. The method was validated by analysis of certified reference materials (Serorm Trace Elements, Bil-

lingstad, Norway). Accurate results were obtained for all elements according to [14].

## 2.5. Hematological Measurements

Ninety phosphates mine workers were taken from the three studied sites (Safaga site, El-Hamrawein site and El-Quseir site) in addition to a control group worker for investigations. All workers were divided according their age into three groups (first group 20–30 years, second group 30 - 40, third group 40 - 50 years) and 10 people as a control group was selected from non-phosphate mine workers (mean age 30 years). Hematological assessment Blood samples (5 mL) were collected by venepuncture of the cubital vein in the antecubital fossa by using a 5 mL disposable syringe. A part of the blood sample was then transferred to sterile vacuum tubes containing an anticoagulant Ethylene Diamine Tetraacetic Acid (EDTA), for whole blood analysis. The remaining blood was collected in sterile vacuum tubes with no added anticoagulant and was kept at room temperature for 2 h, where it was allowed to clot, as this was designated for serum separation for superoxide dismutase (SOD) and Malondialdehyde (MDA). The tubes were transported to a laboratory (Biomedical Lab, National Research Center, Egypt ) for analysis by using hematological investigations consisting of the Hemoglobin (Hb) concentration, RBC count, WBS, platelets count, hematological investigations were estimated in an automated blood counting machine (SYSMEX XE-2100; Sysmexbiomedical Lab, National Research Center, Egypt). Accurate results were obtained for all elements according to [15].

## 2.6. Determination of SOD and MDA

Superoxide dismutase (SOD) activity determination and Malondialdehyde (MDA) determination were performed at Biomedical Lab, National research center, Egypt. SOD determined according to the method of [16], and MDA determined according to method adopted by [17]. The samples were analyzed by spectrophotometer (Milton Roy spectronic 3000 ARRAY double beam spectrometer, manufactured in Germany).

## 3. Results and Discussion

### 3.1. Radioactivity Measurements in Phosphate Samples

Activity concentrations of the natural radionuclides ( $^{226}\text{Ra}$ ,  $^{232}\text{Th}$  and  $^{40}\text{K}$ ) were measured for the investigated phosphate mines samples by HPGe gamma spectrometry. According to the International Atomic Energy Agency, one kilogram of soil typically contains the following amounts of the following three natural radioisotopes 370 Bq of  $^{40}\text{K}$  (typical range 100 - 700 Bq), 25 Bq of  $^{226}\text{Ra}$  (typical range 10 - 50 Bq), 25 Bq of  $^{238}\text{U}$  (typical range 10 - 50 Bq) and 25 Bq of  $^{232}\text{Th}$  (typical range 7 - 50 Bq) (IAEA, 2000). The activity concentrations of these radionuclides in different layers of the investigated samples are shown in **Table 1**. It is recognize from the table that there is a clear variation in activity concentrations of  $^{226}\text{Ra}$ ,  $^{232}\text{Th}$  and  $^{40}\text{K}$  in layer samples (Subsurface, Surface upper bed and Surface lower bed) in every investigated mine and also in the different mines. As can be seen, activity concentrations of all analyzed radionuclides con-

**Table 1.** Activity concentrations of the selected natural radionuclides in the investigated phosphate samples.

Samples	The selected natural radionuclide's in phosphate mine at Safaga (33° 57'E 26° 44'N) ( Bq·kg <sup>-1</sup> )			activity of the investigated radionuclide's at El-Quseir (34° 17'E 26° 06'N) (Bq·kg <sup>-1</sup> )			activity of the investigated radionuclide's at El-Hamrawein (35° 12'E 26° 15'N) ( Bq·kg <sup>-1</sup> )		
	<sup>226</sup> Ra	<sup>232</sup> Th	<sup>40</sup> K	<sup>226</sup> Ra	<sup>232</sup> Th	<sup>40</sup> K	<sup>226</sup> Ra	<sup>232</sup> Th	<sup>40</sup> K
Subsurface samples	222.4 ± 7.2	135.6 ± 4.7	225.2 ± 11.5	122.4 ± 5.8	112.8 ± 7.3	168.7 ± 6.5	115.4 ± 5.9	95.2 ± 4.8	132.8 ± 6.8
Surface samples upper bed	235 ± 7.8	198.2 ± 7.1	264.7 ± 12.3	154.6 ± 7.1	132.7 ± 6.2	212.5 ± 9.7	135.2 ± 6.2	122.7 ± 8.3	145.2 ± 7.8
Surface samples lower bed	255.8 ± 9.3	212.3 ± 9.2	312.8 ± 12.8	188.3 ± 7.6	167.4 ± 8.4	268.9 ± 10.2	165.8 ± 6.6	155.8 ± 9.4	188.6 ± 8.7

siderably fluctuated; for <sup>226</sup>Ra activity concentrations varied from 222.4 to 255.8 Bq·kg<sup>-1</sup>, 122.4 to 188.3 Bq·kg<sup>-1</sup> and 115.4 to 165.8 Bq·kg<sup>-1</sup> for Safaga, El-Quseir and El-Hamrawein, respectively and these concentrations are 5 to 10 times higher than that value recommended by IAEA 2000. For <sup>232</sup>Th activity concentrations varied from 135.6 to 212.3 Bq·kg<sup>-1</sup>, 112.8 to 167.4 Bq·kg<sup>-1</sup> and 132.8 to 188.6 Bq·kg<sup>-1</sup> for Safaga, El-Quseir and El-Hamrawein, respectively and these concentrations are 5 to 8 times higher than that value recommended by IAEA 2000. For <sup>40</sup>K activity concentrations varied from 225.2 to 312.8 Bq·kg<sup>-1</sup>, 168.7 to 268.9 Bq·kg<sup>-1</sup> and 95.2 to 155.8 Bq·kg<sup>-1</sup> for Safaga, El-Quseir and El-Hamrawein, respectively and these concentrations are lower than that value recommended by IAEA 2000. Also the obtained results indicate that the phosphate mine sample of El-Hamrawein has the lowest activity concentration of <sup>226</sup>Ra, <sup>232</sup>Th and <sup>40</sup>K in compared to the phosphate mine samples of El-Quseir and Safaga (El-Hamrawein < El-Quseir < Safaga). This can be discussed in the light of increasing concentration of P<sub>2</sub>O<sub>5</sub> as shown in **Table 2**. In all investigated mines, the remarkable highly activity concentrations are record in surface lower bed samples and this can be discussed as the deep site has more phosphate ore rocks which contains more highly percentage of P<sub>2</sub>O<sub>5</sub> conjugated with uranium, thorium and potassium isotopes. [10], were reported that phosphate mine in Safaga has relatively high levels of <sup>226</sup>Ra The average activity ratio (range) of <sup>226</sup>Ra/<sup>228</sup>Ra in the region of Safaga-Quseir-Marsa Alam city (sample codes from 510 to 549) was 2.25 (0.83 - 5.33). The increases of these ratios, exceeding unity, could be attributed to the phosphate mining in Safaga-Quseir region and/or their geo-chemical behavior in the environment [10].

### 3.2. Concentration of Heavy Metals in the Studied Phosphate Samples

In this study the concentration of some famous hazardous heavy metals reflected the impacts on their environmental sites. Phosphate mines rocks ores and phosphate industry phosphogypsum make them directly affected with these products into human, plants and ecosystem. **Table 3** shows the heavy metals concentration in Safaga, Al Quseir and EL-Hamrawein sites, where the Co was recorded (11.2, 17.8 and 15.2 ppm) respectively, also AS was recorded in Safaga, Al Quseir and EL-Hamrawein (24.9, 26.8 and 20.2 ppm) respectively. As well as the Cd was recorded (0.88, 0.98 and 1.2 ppm) respectively. Also Pb was recorded (28.2, 22.3 and 26.7 ppm) respectively. These data revealed some variations in heavy metals concentration according to phosphate mines site. The data recoded increased levels of Pb concentrations

**Table 2.** Mean percentage of metal oxides in the investigated phosphate samples.

Site	P <sub>2</sub> O <sub>5</sub>	SiO <sub>2</sub>	Al <sub>2</sub> O <sub>3</sub>	Fe <sub>2</sub> O <sub>3</sub>	CaO	MgO	Na <sub>2</sub> O	K <sub>2</sub> O
El-Hamrawein	28	14.5	0.74	4.2	45	0.9	0.8	0.9
El-Quseir	30	15.3	0.52	3.1	47	1.1	0.6	1.3
Safaga	33	11.2	0.33	2.3	41	0.8	1.1	1.2

**Table 3.** Concentration of some heavy metals in the investigated phosphate samples.

phosphate samples	Heavy metals (ppm)		
	As	Cd	Pb
Safaga	24.9	0.88	28.2
El-Quseir	26.8	0.98	22.3
El-Hamrawein	20.2	1.2	26.7

in Safaga site in compared to Al Quseir and EL-Hamrawein sites, as well as Cd in Al Quseir and Safaga sites in compared to Al Quseir site, these results were acceptable when compared to global ratio of heavy metals in phosphate rocks. The concentration of cobalt phosphate rocks widely, generally ranging from about 1 to 40 ppm [18]. On the other hand, soils near cobalt-containing mineral deposits, mining and smelting facilities, or industries manufacturing or using cobalt alloys or chemicals may contain much higher levels of cobalt. Available studies of the carcinogenic effects of cobalt in occupationally-exposed humans have reported mixed results, with both positive and negative results. Lifetime inhalation of cobalt sulfate resulted in increased tumor incidences in both rats and mice [19]. The most common problem causing *cationic* metals (metallic elements whose forms in soil are positively charged cations e.g., Pb<sup>2+</sup>) are mercury, cadmium, lead, nickel, copper, zinc, chromium, and manganese [20]. Nones-essential metals, such as As, Pb and Cd, are toxic even in trace amounts. Intake of cadmium above safe limit causes high blood pressure, liver disease and nerve or brain damage. The essential metals can also produce toxic effects at higher concentrations. They tend to bio-accumulate, cause toxicity to plants and contaminate the food chain [21]. Only a few metals of proven hazardous nature are to be completely excluded in food for human consumption. Thus, only three metals, namely lead, cadmium and mercury, have been included in the regulations of the European Union for hazardous metals [22].

### 3.3. Percentage of Metal Oxides in Phosphate Ores

The chemical analysis of phosphate samples were carried out for showing the correlation between oxides of heavy metals regarding phosphate mine and their hazardous on the occupational workers. As shown in **Table 2**; Safaga mine sample has high content of P<sub>2</sub>O<sub>5</sub> and Na<sub>2</sub>O while it has low content of SiO<sub>2</sub>, Al<sub>2</sub>O<sub>3</sub>, Fe<sub>2</sub>O<sub>3</sub>, CaO and MgO in compared to the other two investigated samples. El-Hamrawein has low content in P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O while it has higher content of Al<sub>2</sub>O<sub>3</sub> and Fe<sub>2</sub>O<sub>3</sub> than the other two investigated samples. Al-Quseir mine sample has moderate contents of the investigated metal oxides except SiO<sub>2</sub>, CaO, MgO and K<sub>2</sub>O are higher values.

The obtained results revealed increasing of siliceous ores which have impact health hazards in workers. The workers exposures to these particles may have many artificial and hematological diseases. Inhalation of silica causes risk of tuberculosis, lung cancer and some autoimmune diseases such as scleroderma and rheumatoid arthritis. Freshly fractured silica dust appears to be more reactive and more hazardous than old or stale dust. This may be a consequence of a relatively higher surface charge on freshly formed particles [23].

### 3.4. Concentration of Some Heavy Metals in Workers Blood

The obtained results of investigated heavy metals in blood of occupational workers at phosphate mines are shown in Table 4. From Table 2 and Table 4, it is clear that there is a good relation between the concentration of the investigated metals (As, Cd and Pb) in phosphate mine samples and their concentration in blood of the occupational workers in these mines. Generally, Table 5 shows that the concentrations of the heavy metals in all ages of the occupational workers are higher than that of the control sample. Also it shows that old ages have higher concentration of these metals than young ages and this agreement with [24], which reported that Cd and Pb concentrations in the blood of aging workers are more than youngest workers. Workers at El-Hamrawein mine have low concentration of Arsenic (As) than other workers at both of Safaga and El-Quseir mines. Workers at Safaga mine have low concentrations of Cd than other workers at both of El-Hamrawein and El-Quseir mines. Workers at El-Quseir mine have

**Table 4.** Concentration of some heavy metals in blood workers (Mean ± SD & n, 10).

Heavy metal (µg/dL)	Control Mean ± SD n, 10	Workers at Safaga			Workers at El-Hamrawein			Workers at El-Quseir		
		Age of 20 to 30 (year)	Age of 30 to 40 (year)	Age of 40 to 50 (year)	Age of 20 to 30 (year)	Age of 30 to 40 (year)	Age of 40 to 50 (year)	Age of 20 to 30 (year)	Age of 30 to 40 (year)	Age of 40 to 50 (year)
As	0.046 ± 0.011	0.076 ± 0.015	0.121 ± 0.019	0.128 ± 0.039	0.077 ± 0.007	0.116 ± 0.032	0.123 ± 0.040	0.078 ± 0.021	0.149 ± 0.016	0.152 ± 0.008
Cd	0.086 ± 0.021	0.164 ± 0.027	0.162 ± 0.023	0.192 ± 0.060	0.181 ± 0.011	0.204 ± 0.018	0.224 ± 0.054	0.171 ± 0.025	0.194 ± 0.027	0.214 ± 0.023
Pb	0.362 ± 0.068	0.994 ± 0.105	1.244 ± 0.027	1.662 ± 0.180	0.882 ± 0.161	1.218 ± 0.075	1.196 ± 0.227	0.868 ± 0.077	1.121 ± 0.088	1.068 ± 0.171

**Table 5.** Concentration of some blood characteristics for workers (Mean ± SD & n, 10).

Blood Parameters	control	Workers at Safaga			Workers at El-Quseir			Workers at El-Hamrawein		
		Age of 20 to 30 (year)	Age of 30 to 40 (year)	Age of 40 to 50 (year)	Age of 20 to 30 (year)	Age of 30 to 40 (year)	Age of 40 to 50 (year)	Age of 20 to 30 (year)	Age of 30 to 40 (year)	Age of 40 to 50 (year)
Hemoglobin mg/dl	12.55 ± 0.31	11.96 ± 0.21	11.02 ± 0.32	10.38 ± 3.3	12.46 ± 0.11	11.56 ± 0.33	10.4 ± 0.065	12.8 ± 0.14	11.27 ± 0.25	9.4 ± 0.13
Red cell count × 10 <sup>6</sup> /cmm <sup>3</sup> mil	5.21 ± 0.09	4.34 ± 0.05	4.04 ± 0.08	3.96 ± 0.06	4.3 ± 0.02	4.14 ± 0.04	3.72 ± 0.073	21.6 ± 0.23	3.71 ± 0.024	3.38 ± 0.042
Leucocytic count/cmm <sup>3</sup>	6450 ± 9.1	6744 ± 8.3	5830 ± 5.2	7300 ± 5.1	5819 ± 7.84	5770 ± 6.94	6360 ± 7.36	6970 ± 8.53	6750 ± 9.66	6300 ± 6.22
Platelet count × 10 <sup>3</sup> /cmm <sup>3</sup>	290 ± 3.8	185 ± 4.3	181.2 ± 2.3	152.2 ± 2.3	239 ± 2.48	187.6 ± 5.28	173.2 ± 4.22	237 ± 4.88	166 ± 2.85	115.4 ± 2.84

low concentrations of Pb than other workers at both of EL-Hamrawein and Safaga mines. The variation of the investigated heavy metals between the workers at the different sites agrees with their concentrations in the phosphate mines. The enhancement of As, Cd and Pb in blood workers could be due to their accumulation through the inhalation of phosphate dust which may be released into the environment during mining process.

### 3.5. Determination of Hematological Parameters

The prevalence of occupational health hazards for phosphat mine workers in Safaga, Al Quseir and EL-Hamrawein. The experimental study were designed according to three different age stage were start from (20 - 30 y), (30 - 40 y) and (40 - 50 y) male workers selected in three phosphate mine sites and were undergo to hematological studying and other biochemical studies. In this study the results in **Table 4** shows the hematological parameters in Safaga phosphate mine workers at age (20 - 30) the Hemoglobin, Red cell count, Leucocytic count/and Platelet count recoded 11.96 mg/dl,  $4.34 \times 10^6/\text{cmm}^3$  mil, 6744 cmm and  $185 \times 10^3/\text{cmm}$  respectively. On the hand phosphate mine workers in safaga at age (30 - 40 y) recorded in Hb, RBC, WBCs and PLTC 11.02 mg/dl,  $4.04 \times 10^6/\text{cmm}^3$  mil, 5830/cmm and  $181.2 \times 10^3/\text{cmm}^3$ . Also mine workers in safaga at age (40 - 50) recorded in Hb, RBCs, WBCs and PLTC 10.38 mg/dl,  $3.96 \times 10^6/\text{cmm}^3$  mil, 7300 cmm and  $152.2 \times 10^3/\text{cmm}^3$ . The interpretation of this result was concepts for the duration time and intervals periods for the phosphate mine workers. It was clear when the workers increased in intervals and progress in life age the hematological functions decreased. The phosphate mine workers were stress under the effects of inhalable particles of heavy metals, chemicals compounds and radioactive nuclides in phosphate rocks and ores, which leads to different types of occupational hazards diseases. This result was conformed with the results of [25], which observed on workers in phosphate mines were exposed to inhalable heavy metals and radionuclide for a long time lead to hematological disorders. The hematological disorders were take place in phosphate mine works undergoes the oxidative stress factors. Oxidative damage is one of the results of this imbalance, comprising oxidative modification of cellular macromolecules It has long been recognized that ROS are harmful for cells, because they injure heamoglobine and RBCs, which leads to structural and functional impairments [8] [9]. On the other hand, hematological parameters in Al Quseir phosphate mine workers shows in **Table 4**. Hb, and RBCs, WBCs and Platelet count in workers revealed that normal limits in Hb% and RBCs, WBCs and platelet count for workers at age 20 - 30 y. Also workers at age 30 - 40 y recorded moderate level for hematological parameters. On the other hand, workers at age from (40 - 50 y) recorded decrease in hematological parameters. The interpretation of the data was due to the interval period and long time for workers duration in phosphate mine, and the age and life spine of workers. It is notable in these data the oldest age workers do not have any resistance against oxidative stress and losses of antioxidant.

### 3.6. Determination of SOD and MDA

In this study it is evident that phosphate miners were exposed to oxidative stress emitted

**Table 6.** Concentration of SOD and MDA in blood workers (Mean  $\pm$  SD & n, 10).

Parameters	Control Mean $\pm$ SD n, 10	Workers at Safaga			Workers at El-Quseir			Workers at El-Hamrawein		
		Age of 20 to 30 (year)	Age of 30 to 40 (year)	Age of 40 to 50 (year)	Age of 20 to 30 (year)	Age of 30 to 40 (year)	Age of 40 to 50 (year)	Age of 20 to 30 (year)	Age of 30 to 40 (year)	Age of 40 to 50 (year)
SOD U/g Hb	4.22 $\pm$ 0.85	8.91 $\pm$ 2.63	10.44 $\pm$ 2.89	11.54 $\pm$ 3.1	12.48 $\pm$ 3.11	15.49 $\pm$ 3.22	18.22 $\pm$ 3.89	7.22 $\pm$ 0.46	8.24 $\pm$ 0.97	10.83 $\pm$ 2.33
MDA U/L	10.22 $\pm$ 0.92	28.43 $\pm$ 2.32	32.58 $\pm$ 2.78	35.66 $\pm$ 2.98	24.35 $\pm$ 0.57	32.58 $\pm$ 3.11	38.75 $\pm$ 2.13	40.66 $\pm$ 3.44	46.75 $\pm$ 3.22	42.83 $\pm$ 3.89

from the particles of heavy metals and radionuclides. The oxidative stress plays a role in damage of immune system and makes a hematological disorder for phosphate workers. These symptoms were very clear in oldest workers than youngest works. The results in **Table 6** show the concentration of superoxide dismutase (SOD) and Malonaldehyde in blood of phosphate mine workers. The SOD and MDA recorded in Safaga phosphate mine workers age (20 - 30 y)  $8.91 \pm 2.63$  U/g Hb and  $10.22 \pm 0.92$  U/L compared to control non workers  $4.22 \pm 0.85$  U/g Hb and  $10.22 \pm 0.92$  respectively. This results was confirmed of results [26], he recorded that increased level of SOD activities in phosphate miners. Reduction in SOD activity as observed by us may be due to an increased endogenous production of ROS as evidenced by increased SOD. This decrease in antioxidant enzyme may be related to the consumption of activated enzymes against oxidative stress.

**Table 6** shows the concentrations of SOD and MDA in Al Quseir and EL-Hamrawein phosphate mine workers. The data revealed highly increased SOD in Al Quseir phosphate mine workers at all age stage in compared to control group non workers. On the other hand data in **Table 6**, recorded slightly increased in SOD activity at EL-Hamrawein phosphate mine workers, also MDA in EL-Hamrawein phosphate mine workers recorded highly activity for all interval age (20 - 30), (30 - 40 y) and (40 - 50 y) where  $40.66 \pm 3.44$ ,  $46.75 \pm 3.22$  and  $42.83 \pm 3.89$  U/L respectively in compared to non workers control group  $10.22 \pm 0.92$ . These results was confirmed with [22], where recorded that MDA levels are increased in phosphate mine workers, due to exposed to particles of heavy metal and radionuclide. These toxic particles lead to oxidative stress. Increased oxidative stress generation free radicals and increased reactive oxygen species (ROS). However, direct evidence for oxidative stress is often obscure following long-term and environmentally-relevant for workers in phosphate mines.

#### 4. Conclusions

This study revealed that MDA and SOD were acting as smart biomarkers for determination types and category which had been related to the health impacts on phosphate miners workers during exposures to heavy metals and natural radioactivity. An increased serum MDA level in phosphate miners may indicate a general oxidant effect of phosphate heavy metals dust and natural radioactive nuclides. Also, decreased SOD activity in phosphate miner might be a marker of diminished antioxidant defense system which was caused by heavy metals and natural radioactivity.

The workers in phosphate mine plants are more susceptible to health problems.

MDA, SOD and hematological tests acting specific tests to confirm the effect of phosphate mines' pollutants can be carried out. This study also investigates occupational and heavy metals exposure in the phosphate mines. On the other hand, we need to implement radiation regulations and standards through improving the working conditions to reduce the occupational radiation exposure to the accepted levels recommended by ICRP-and IAEA-Safety standard.

Radiological and environmental safety should be considered in phosphate mines. Regulations should be issued and applied by the administration of these sites. Radiological follow-up should be a routine. Medical follow-up system should be applied. Occupational health monitoring should be take into account according to US EPA 2010 and WHO 2006 standard and regulation for health measurements in occupational mines.

The activity concentrations of the investigated radionuclides considerably fluctuated in the studied mines. The concentrations of  $^{226}\text{Ra}$  are 5 to 10 times higher than that value recommended by IAEA 2000. The concentrations of  $^{232}\text{Th}$  are 5 to 8 times higher than that value recommended by IAEA 2000. The  $^{40}\text{K}$  concentrations are lower than that value recommended by IAEA 2000. The concentrations of the investigated heavy metals (As, Cd and Pb) in old ages of the occupational workers are higher than those of young ages, while these metals are concentrating in all ages more than the control people sample.

## References

- [1] UNSCEAR-A (2000) Annex-A, Dose Assessment Methodologies. United Nations Scientific Committee on the Effects of Atomic Radiation.
- [2] Christie, D.H., Chu, M.C. and Carr, Z. (2010) Global Networking for Biodosimetry Laboratory Capacity in Radiation Emergencies. *Health Physics*, **92**, 168.
- [3] United Nations Scientific Committee on the Effects of Atomic Radiation Sources (UNSCEAR) (2000) Effects and Risks of Ionizing Radiation. Report to the General Assembly with Annex B. United Nations, New York.
- [4] International Atomic Energy Authority (2004) Extent of Environmental Contamination by Naturally Occurring Radioactive Materials (NORM) and Technological Options for Mitigation. Technical Report No. 419, Vienna.
- [5] Al-Trabulsy, H.A., Khater, A.E.M. and Habbani, F.I. (2011) Radioactivity Levels and Radiological Hazard Indices at the Saudi Coastline of the Gulf of Aqaba. *Radiation Physics and Chemistry*, **80**, 343-348. <http://dx.doi.org/10.1016/j.radphyschem.2010.09.002>
- [6] European Commission (1999) Reference Levels for Workplaces Processing Materials with Enhanced Levels of Naturally Occurring Radionuclides. Radiation Protection No. 95, European Commission.
- [7] Gnandi, K., Rezaie, B. and Edoth, A.P. (2009) The Geochemical Characterization of Mine Effluents from the Phosphorite Processing Plant of Kpémé (Southern Togo). *Mine Water and the Environment*, **28**, 65-73. <http://dx.doi.org/10.1007/s10230-008-0058-0>
- [8] Nordberg, J. and Arner, E.S.J. (2001) Reactive Oxygen Species, Antioxidants and the Mammalian Thiredoxin System. *Free Radical Biology and Medicine*, **31**, 1287-1312. [http://dx.doi.org/10.1016/S0891-5849\(01\)00724-9](http://dx.doi.org/10.1016/S0891-5849(01)00724-9)

- [9] Borm, P.J.A., Bast, A., Wouters, E.F.M., Slangen, J.J.M., Swaen, G.M.H. and Boorder de, Tj. (1986) Red Blood Cell Antioxidant Parameters in Silicosis. *International Archives of Occupational and Environmental Health*, **58**, 235-244. <http://dx.doi.org/10.1007/BF00432106>
- [10] El-Mamoney, M.H. and Khater, A.E.M. (2004) Environmental Characterization and Radio-Ecological Impacts of Non-Nuclear Industries on the Red Sea Coast. *Journal of Environmental Radioactivity*, **73**, 151-168. <http://dx.doi.org/10.1016/j.jenvrad.2003.08.008>
- [11] Papaefthymiou, H. and Psychoudaki, M. (2008) Natural Radioactivity Measurements in the City of Ptolemais (Northern Greece). *Journal of Environmental Radioactivity*, **99**, 1011-1017. <http://dx.doi.org/10.1016/j.jenvrad.2007.12.001>
- [12] Keyser, R.M. (1995) Characterization and Applicability of Low-Background Germanium Detectors. Technical Note, EG&G ORTEC, Oak Ridge.
- [13] Price, W.J. (1983) Spectrochemical Analysis by Atomic Absorption. Wiley Heyden Ltd., Chichester.
- [14] Gajek, R. (2011) Optimization of Rapid Heavy Metals Blood Analysis for Ultra Low Detection Limits by ICP-MS with ISIS. Mass Spectrometry Applications to the Clinical Lab (MSACL), San Diego.
- [15] Dacie, S.T. and Lewis, S.M. (1991) Practical Haematology. 7th Edition, Churchill Livingstone Medical Division of Long Man Group U. K. Ltd., 26-36.
- [16] Misra and Fridovich. (1972) The Role of Superoxide Anion in the Autoxidation of Epinephrine and a Simple Assay for Superoxide Dismutase. *The Journal of Biological Chemistry*, **247**, 3170-3175.
- [17] Draper, H.H. and Hardley, M. (1990) Malondialdehyde Determination as Index of Lipid Peroxidation. *Methods in Enzymology*, **186**, 421-431. [http://dx.doi.org/10.1016/0076-6879\(90\)86135-I](http://dx.doi.org/10.1016/0076-6879(90)86135-I)
- [18] Conceicao, E.T. and Bonotio, D.M. (2006) Radionuclides, Heavy Metals and Fluorine Incidence at Tapira, Phosphate Rocks. Brazil and Their Industrial By-Products. *Environmental Pollution*, **139**, 232-243. <http://dx.doi.org/10.1016/j.envpol.2005.05.014>
- [19] Ming-Ho, Y. (2005) Environmental Toxicology: Biological and Health Effects of Pollutants, Chap. 12, 2nd Edition, CRC Press LLC, Boca Raton.
- [20] U.S. Environmental Protection Agency, Office of Solid Waste (1994) Other Mining Sectors: Phosphate and Molybdenite—Extraction and Beneficiation of Ores and Minerals, Volume 7. EPA/530-R-94-034, NTIS/PB94-201 001, Washington DC.
- [21] OSHA (2001) Construction Industry, Cobalt Metal, Dust, and Fume. Occupational Safety and Health Administration. [http://www.osha.gov/OshStd\\_data/1926\\_0055.html](http://www.osha.gov/OshStd_data/1926_0055.html)
- [22] Godt, J., Scheidig, F., Grosse-Siestrup, C., Esche, V., Brandenburg, P. and Reich, A. (2006) The Toxicity of Cadmium and Resulting Hazards for Human Health. *Journal of Occupational Medicine and Toxicology*, **1**, 22-33. <http://dx.doi.org/10.1186/1745-6673-1-22>
- [23] Ibrahim, M.S., Atta, E.R. and Zakaria, K.M. (2013) Assessment of Natural Radioactivity of Some Quarries Raw Materials in El-Minya Governorate, Egypt. *Arab Journal of Nuclear Science and Applications*, **47**, 208-216.
- [24] Ogwuegbu, M.O. and Ijioma, M.A. (2003) Effects of Certain Heavy Metals on the Population due to Mineral Exploitation. *International Conference on Scientific and Environmental Issues in the Population, Environment and Sustainable Development in Nigeria*, University of Ado Ekiti, Ekiti State, 10 October 2003, 8-10.
- [25] Agency for Toxic Substances and Disease Registry (ATSDR) (2004) Toxicological Profile for Copper. US Department of Health and Human Services, Public Health Service, Centers

for Diseases Control, Atlanta.

- [26] Codorean, E., Raducanu, A., Albulescu, L., Tanase, C., Meghea, A. and Albulescu, R. (2011) Multiplex Cytokine Profiling in Whole Blood from Individuals Occupationally Exposed to Particulate Coal Species. *Romanian Biotechnological Letters*, **16**, 6748-6759.



Scientific Research Publishing

**Submit or recommend next manuscript to SCIRP and we will provide best service for you:**

Accepting pre-submission inquiries through Email, Facebook, LinkedIn, Twitter, etc.

A wide selection of journals (inclusive of 9 subjects, more than 200 journals)

Providing 24-hour high-quality service

User-friendly online submission system

Fair and swift peer-review system

Efficient typesetting and proofreading procedure

Display of the result of downloads and visits, as well as the number of cited articles

Maximum dissemination of your research work

Submit your manuscript at: <http://papersubmission.scirp.org/>

Or contact [jep@scirp.org](mailto:jep@scirp.org)