

# Chromosomal Location of a Recessive Red-Eye Mutant Gene in the Brown Planthopper *Nilaparvata lugens* (Stål) (Insecta: Hemiptera)

Jirapong Jairin<sup>1\*</sup>, Phikul Leelagud<sup>1</sup>, Adisak Pongmee<sup>2</sup>, Kanuengnij Srivilai<sup>3</sup>

<sup>1</sup>Ubon Ratchathani Rice Research Center, Ubon Ratchathani, Thailand

<sup>2</sup>Roi Et Agricultural Research and Development Center, Roi Et, Thailand

<sup>3</sup>Sakon Nakhon Rice Research Center, Sakon Nakhon, Thailand

Email: \*jirapong.j@rice.mail.go.th

**How to cite this paper:** Jairin, J., Leelagud, P., Pongmee, A. and Srivilai, K. (2017) Chromosomal Location of a Recessive Red-Eye Mutant Gene in the Brown Planthopper *Nilaparvata lugens* (Stål) (Insecta: Hemiptera). *Advances in Entomology*, 5, 33-39. <http://dx.doi.org/10.4236/ae.2017.51003>

**Received:** January 2, 2017

**Accepted:** January 16, 2017

**Published:** January 19, 2017

Copyright © 2017 by authors and Scientific Research Publishing Inc.

This work is licensed under the Creative Commons Attribution International License (CC BY 4.0).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

## Abstract

The color of compound eyes is an important biological characteristic of insects. A red eye color mutation is commonly found in the brown planthopper (BPH), *Nilaparvata lugens* (Stål) (Hemiptera: Delphacidae), a serious insect pest of rice in tropical and temperate Asia. The genetic inheritance and physiological effect of the eye color mutation in the BPH have been studied, but the location of a *red* gene controlling the red eye mutant phenotype on a chromosome has not been elucidated. In this study, simple sequence repeats (SSRs), together with bulked segregant analysis (BSA), was performed to identify and map the location of the *red* gene. A total of 387 SSR markers distributed throughout the BPH autosome were used to survey two bulked DNA samples. Samples were generated from 29 brown-eyed and 29 red-eyed individuals derived from an F<sub>2</sub> generation of a cross between brown-eyed wild type and red-eyed mutant colonies. The SSR marker BM20 was shown to be associated with the red eye mutant phenotype. Ninety-five offspring of the F<sub>2</sub> generation were then used to map the gene. The present study constitutes the discovery of the location of the *red* gene, which may lead to the acquisition of the genetic determinant of the compound eye color mutation in BPH.

## Keywords

Brown Planthopper, Bulk Segregant Analysis, Linkage Map, Red-Eyed Mutant, Simple Sequence Repeat

## 1. Introduction

The brown planthopper (BPH) *Nilaparvata lugens* (Stål) (Arthropoda: Insecta), the most destructive phloem-sap-sucking insect pest of rice in Asia, belongs to the order Hemiptera, family Delphacidae. The BPH has a diploid chromosome number of 30 (28

autosomes combined with the sex determination system XY and XX in males and females, respectively) [1]. Among the few visible mutations in the BPH, eye color is one of the mutations that are regularly observed in laboratories [2] [3]. In Delphacidae, red-eyed mutants were reported in the small brown planthopper *Laodelphax striatellus* (Fallén) [4] and BPH [5]. Red-or orange-eyed mutant laboratory colonies of BPH have been established. Inheritance studies revealed that a single recessive gene on an autosome controlled the eye color mutations.

Compound eyes are the visual organs for the majority of insects. It is an important biological characteristic. The biological and biochemical characterizations of eye color in BPH mutants have been described. The mating behavior, egg production, developmental duration and mortality of nymphal stages were not significantly different between the normal-eyed and colored-eye mutant colonies [2] [3]. However, some phenotypes (*i.e.*, egg lethal effect, poor fecundity and survival ability) might be related to the eye color mutations [3] [5]. Genes having a stronger compensation ability for reproduction may also closely link to the red eye allele [6].


The color of BPH eyes is determined by ommochrome pigments. Xanthommatin (the pigment causing brown eye color) and pteridine (the pigment causing red eye color) are two primary forms of the pigments. Reduction of pigment granules and pigment concentrations is the biochemical basis of the red eye phenotype in the BPH [2]. It is believed that mutations in genes common to both pigments caused a reduction of both pigments [7] [8]. The causal mutation may occur in xanthommatin or pteridine biosynthesis, transportation of eye pigments, or in pigment granule formation [8].

Recent advances in BPH genomics facilitate the study of various phenotypes of the BPH. Numerous DNA markers were developed [9] [10] that made it possible to identify and map some genes in the BPH through linkage to existing DNA markers [10] [11]. In the present study, bulked segregant analysis (BSA) [12] with SSR markers was used to detect and locate the chromosomal location of the *red* gene controlling the red eye mutation in the BPH. The linked markers from this study will facilitate map-based cloning to identify the *red* gene in the BPH.

## 2. Materials and Methods

### 2.1. Insect Populations and Crosses

Two BPH populations collected in Thailand were used in this study: (1) Kalasin-2013 (KLS13) was collected in Kalasin province, northeastern Thailand in 2013 and (2) Krabi-2013 (KBI13) was collected in Krabi province, southern Thailand in 2013 (Figure 1). Both populations have been maintained in laboratory conditions for approximately 40 generations. A red-eyed mutant colony was established from a red-eyed female adult found in the KLS13 population. KBI13 was selected as a wild-type colony to cross with the red-eyed mutant colony to generate an F<sub>2</sub> mapping population. All BPH strains were reared on rice seedlings of cv. "TN1" and maintained in the laboratory at 26°C ± 2°C under a 14 L:10 D photoperiod. To generate the mapping population, a pair of insects from two different colonies was selected and crossed using single-pair mating. An F<sub>2</sub> population was generated from a cross of KLS13 (♀) × KBI13 (♂). The crosses were performed by placing a virgin female and a male from each colony on "TN1" seedlings



**Figure 1.** Locations of the collection sites of two brown planthopper, *Nilaparvata lugens* (Stål) populations in Thailand.

in a glass tube. After mating and oviposition, the parental insects were collected and stored in 95% ethanol. Seedlings with eggs were kept until the eggs hatched. Newly hatched nymphs of an  $F_1$  progeny were transferred into a plastic 800 mL container containing 80 to 100 seedlings and reared to the fifth-instar nymph stage. An individual fifth-instar nymph of each sex was selected and reared in a separate glass tube to obtain unmated adults. The  $F_1$  progenies were then allowed to inbreed to generate the  $F_2$  generation. All offspring were then collected and stored in 95% ethanol.

## 2.2. Total DNA Extraction and Whole-Genome Amplification

Genomic DNA of the preserved samples was isolated individually using a modified version of the potassium acetate procedure of Dellaporta *et al.* [13]. To avoid viability problems that may occur at later stages, second- and third-instar nymph stages of  $F_2$  offspring were collected for DNA isolation. The whole-genome amplification technique [14] was used to produce large amounts of DNA from individual insects for large-scale genotyping.

## 2.3. Tagging and Mapping of the Red-Eye Mutant Gene

BSA with SSR markers was used to tag a red-eye mutant locus. Based on the presence of eye color of the BPH, we generated two groups of 29 brown-eyed (B) and 29 red-eyed (R) progenies from the  $F_2$  population. A total of 387 SSR markers from Jairin *et al.* [9] and Jing *et al.* [10] covering 14 chromosomes of the BPH were selected to identify the R

and B groups. The genetic linkage map was calculated by Join Map 4 [15] using genotype data from 95  $F_2$  individuals derived from crosses of KLS13/KBI13. Grouping of markers was performed using a minimum-independence LOD (logarithm of odds) threshold of 5.0. Groups were converted into linkage maps using the regression algorithm provided by Join Map with a recombination frequency of less than 0.45, and then Kosambi's mapping function [16] was used to calculate map distances. Linkage maps were drawn using the Map Chart software for Windows [17].


### 3. Results and Discussion

#### 3.1. Establishment of the Red-Eyed Colony

A red-eyed female adult BPH was isolated from a field population collected from a farmer's rice field in KLS to establish a red-eyed mutant colony. The colony was used as genetic material for mapping the *red* gene. The mutant individuals could be easily discriminated from the brown-eyed individuals (Figure 2). Individuals in the mutant colony had the red eye phenotype at all developmental stages. We observed the red eye phenotype in both macropterous and brachypterous adults, as well as adults with both brown and black body colors. Except for eye color, no other external differences were observed between the colonies (Figure 3).



**Figure 2.** Eye colors of the brown-eyed wild type (a) and red-eyed mutant (b) brown planthopper, *Nilaparvata lugens* (Stål) female adults.



**Figure 3.** Wild type brown-eyed and red-eyed mutants of brown planthopper, *Nilaparvata lugens* (Stål) at different developmental stages and external characteristics. (a), (b): first in star nymphs, (c), (d): adult females with different wing forms, e: brown and black body colors of adult brown-eyed and red-eyed offspring.

### 3.2. Crossing Experiment

Cross mating between the homozygous normal brown-eyed KBI13 (♀) and red-eyed mutant KLS13 (♂) produced F<sub>1</sub> offspring with wild-type brown eye color. The F<sub>1</sub> offspring from the reciprocal crosses also showed only normal brown eye color. Inbreeding between heterozygous F<sub>1</sub> progenies resulted in F<sub>2</sub> offspring with brown eye color and red eye color at the expected ratio of 3:1 for a single recessive gene inheritance. The sex ratio in the offspring produced from most crosses was equal based on a chi-square test analysis (Table 1). These results suggest that the inheritance of the red eye mutant phenotype is controlled by a single autosomal recessive gene same as the previous findings [2] [3].

### 3.3. Mapping of the Red-Eye Mutant Gene


To detect the map position of the *red* locus, BSA was employed to identify SSR markers linked to the red eye mutant phenotype. Two bulks from F<sub>2</sub> offspring were generated, each consisting of 29 individual brown-eyed and red-eyed second and third instar nymphs. A total of 387 SSR markers distributed throughout 14 chromosomes were used to screen the pair of bulks. Only SSR markers on sex chromosomes were not selected for the analysis. The primer BM20 generated amplification products for heterozygous alleles in the B bulk and homozygous alleles in the R bulk. The brown-eyed F<sub>2</sub> offspring consisted of nymphs that were homozygous or heterozygous for the mutant allele, while the red-eyed group consisted of only nymphs that were homozygous at the locus. This result indicated that the *red* gene was linked to BM20 in the F<sub>2</sub> population.

We employed an additional 20 SSR markers surrounding the BM20 locus on chromosome 9 [10] to assay 95 F<sub>2</sub> offspring. To obtain accurate estimates of marker positions on the map, we only selected clearly polymorphic markers and carefully scored them to reduce errors. Unfortunately, many markers could not be scored, mostly due to banding patterns that were too difficult to interpret, including multiple bands and bands that did not show clear polymorphism on the 6% polyacrylamide gel. Of 20 SSR markers tested, only three markers showed clearly distinguishable polymorphisms and were easy to score. A genetic linkage map was constructed based on the segregation data. Finally, the *red* gene was mapped to a location between SSR markers BM20 and NLGS182 (Figure 4) on chromosome 9 according to Jing *et al.*'s map [10] or to the linkage group 2 according to our previous map [9]. This is the first study to successfully locate the *red* gene in the genome of this important insect pest of rice using marker-based genetic mapping. We have demonstrated that BSA coupled with SSR markers can

**Table 1.** Eye colors in F<sub>2</sub> offspring of crosses between brown-eyed wild-type (B) and red-eyed mutant (R) brown planthopper, *Nilaparvata lugens* (Stål).

Cross (♀ × ♂)	B	R	Total	$\chi^2_{3:1}$	P	Female		Male		B ♀: ♂		R ♀: ♂	
						B	R	B	R	$\chi^2_{1:1}$	P	$\chi^2_{1:1}$	P
R × B	48	19	67	0.40	0.53	25	12	23	7	0.08	0.77	0.32	0.25
B × R	34	10	44	0.12	0.73	19	4	15	6	0.47	0.49	0.40	0.53
B × R*	66	29	95	1.55	0.21	-	-	-	-	-	-	-	-

\*F<sub>2</sub> offspring from this cross were used as a mapping population. All offspring were collected at the 2<sup>nd</sup>-3<sup>rd</sup> instar nymphal stages.



**Figure 4.** Genetic linkage maps indicating the positions of the red eye mutant gene (*red*) on chromosome 9 of brown planthopper, *Nilaparvata lugens* (Stål). Marker names are listed to the right of the chromosome. The distance between markers is measured in centi Morgans.

easily and rapidly identify DNA markers linked to the *red* gene. The results from genetic crosses and genetic inheritance from the present and previous studies [2] [3] revealed that the red eye color is inherited as a simple Mendelian autosomal recessive gene. The location of the *red* gene on BPH chromosome 9 confirms that the red eye mutation is located on an autosome.

The red-eyed mutant colony has the potential for utilization in BPH research as a visible genetic marker and can be applied in various studies of genetics and ecology at a molecular level (e.g., population dynamics and genetics, evolution of compound eyes, visual acuity, migration, mating behavior, transgenic and RNAi technologies [7]). A xanthommatin-related gene, karmoisin (*Nlka*), reported to be involved in eye mutations in some insect species [18], was expected to cause the red eye mutation phenotype in the BPH. Liu *et al.* [7] identified the karmoisin gene from the BPH and found that the red eye phenotype of the BPH has no close connection with this gene. To identify the gene related to the red eye mutation, map-based cloning is one of the most effective approaches for cloning the *red* gene. Our finding will facilitate fine-scale mapping experiments to isolate the genes responsible for the red-eye mutation in the BPH through genotyping-by-sequencing approaches.

## Acknowledgements

The authors greatly acknowledge the financial support from the Rice Department, Ministry of Agriculture and Cooperatives, Thailand (Project No. 231473).

## References

- [1] Noda, H. and Tatewaki, R. (1990) Re-Examination of Chromosomes of Three Species of Rice Planthoppers (Homoptera: Delphacidae). *Applied Entomology and Zoology*, **25**, 538-540.
- [2] Liu, S.H., Yao, J., Yao, H.W., Jiang, P.L., Yang, B.J. and Tang, J. (2014) Biological and Biochemical Characterization of a Red-Eye Mutant in *Nilaparvata lugens* (Hemiptera: Delphacidae). *Insect Science*, **21**, 469-476. <https://doi.org/10.1111/1744-7917.12049>
- [3] Seo, B.Y., Jin, K.J. and Yeongtae, K. (2011) An Orange-Eye Mutant of the Brown Planthop-

- per, *Nilaparvata lugens* (Hemiptera: Delphacidae). *Journal of Asia-Pacific Entomology*, **14**, 469-472. <https://doi.org/10.1016/j.aspen.2011.06.005>
- [4] Ishii, T. (1966) Some Hereditary and Ecological Observations on Small Brown Planthopper, *Laodelphax striatellus* (Fallén), Having Red-Colored Eye. *Japanese Journal of Applied Entomology and Zoology*, **10**, 64-68. <https://doi.org/10.1303/jjaez.10.64>
- [5] Mochida, O. (1970) A Red-Eyed form of the Brown Planthopper, *Nilaparvata lugens* (Stål) (Hom., Auchenorrhyncha). *Bulletin of the Kyushu Agricultural Experiment Station*, **15**, 141-273.
- [6] Liu, S.H., Yang, B.J., Luo, J., Tang, J. and Wu, J.C. (2015) A Comparative Study on the Population Fitness of Three Strains of *Nilaparvata lugens* (Hemiptera: Delphacidae) Differ in Eye Color-Related Genes. *Journal of Economic Entomology*, **108**, 1675-1682. <https://doi.org/10.1093/jee/tov154>
- [7] Liu, S., Tang, J., Luo, J., Yang, B., Wang, A. and Wu, J. (2016) Cloning and Characterization of Karmoisin Homologue Gene (*Nlka*) in Two Brown Planthopper Strains with Different Eye Colors. *Rice Science*, **23**, 104-110. <https://doi.org/10.1016/j.rsci.2016.02.005>
- [8] Rasgon, J.L. and Scott, T.W. (2004) *Crimson*: A Novel Sex-Linked Eye Color Mutant of *Culex pipiens* L. (Diptera: Culicidae). *Journal of Medical Entomology*, **41**, 385-391. <https://doi.org/10.1603/0022-2585-41.3.385>
- [9] Jairin, J., Kobayashi, T., Yamagata, Y., Sanada-Morimura, S., Mori, K., Tashiro, K., Kuhara, S., Kuwazaki, S., Urino, M., Suetsugu, Y., Yamamoto, K., Matsumura, M. and Yasui, H. (2013) A Simple Sequence Repeat- and Single-Nucleotide Polymorphism-Based Genetic Linkage Map of the Brown Planthopper, *Nilaparvata lugens*. *DNA Research*, **20**, 17-30. <https://doi.org/10.1093/dnares/dss030>
- [10] Jing, S., Zhang, L., Ma, Y., Liu, B., Zhao, Y., Yu, H., Zhou, X., Qin, R., Zhu, L. and He, G. (2014) Genome-Wide Mapping of Virulence in Brown Planthopper Identifies Loci That Break Down Host Plant Resistance. *PLoS One*, **9**, e98911. <https://doi.org/10.1371/journal.pone.0098911>
- [11] Kobayashi, T., Yamamoto, K., Suetsugu, Y., Kuwazaki, S., Hattori, M., Jairin, J., Sanada-Morimura, S. and Matsumura, M. (2014) Genetic Mapping of the Rice Resistance-Breaking Gene of the Brown Planthopper *Nilaparvata lugens*. *Proceedings of the Royal Society B*, **281**, 0726. <https://doi.org/10.1098/rspb.2014.0726>
- [12] Michelmore, R.W., Paran, I. and Kesseli, R.V. (1991) Identification of Markers Linked to Disease-Resistance Genes by Bulk Segregant Analysis: A Rapid Method to Detect Markers in Specific Genomic Regions by Using Segregating Populations. *Proceedings of the National Academy of Sciences of the United States of America*, **88**, 9828-9832. <https://doi.org/10.1073/pnas.88.21.9828>
- [13] Dellaporta, S.L., Wood, J. and Hicks, J.R. (1983) A Plant DNA Miniprep: Version II. *Plant Molecular Biology Reporter*, **1**, 19-21. <https://doi.org/10.1007/BF02712670>
- [14] Dean, F.B., Nelson, J.R., Giesler, T.L. and Lasken, R.S. (2001) Rapid Amplification of Plasmid and Phage DNA Using *Phi29* DNA Polymerase and Multiply-Primed Rolling Circle Amplification. *Genome Research*, **11**, 1095-1099. <https://doi.org/10.1101/gr.180501>
- [15] Van Ooijen, J.W. (2006) JoinMap<sup>®</sup>4, Software for the Calculation of Genetic Linkage Maps in Experimental Populations. Kyazma BV, Wageningen.
- [16] Kosambi, D.D. (1944) The Estimation of Map Distances from Recombination Values. *Annals of Eugenics*, **12**, 172-175. <https://doi.org/10.1111/j.1469-1809.1943.tb02321.x>
- [17] Voorrips, R.E. (2002) MapChart: Software for the Graphical Presentation of Linkage Maps and QTLs. *Journal of Heredity*, **93**, 77-78. <https://doi.org/10.1093/jhered/93.1.77>
- [18] Lloyd, V., Ramaswami, M. and Krämer, H. (1998) Not Just Pretty Eyes: *Drosophila* Eye-Color Mutations and Lysosomal Delivery. *Trends in Cell Biology*, **8**, 257-259. [https://doi.org/10.1016/S0962-8924\(98\)01270-7](https://doi.org/10.1016/S0962-8924(98)01270-7)

**Submit or recommend next manuscript to SCIRP and we will provide best service for you:**

Accepting pre-submission inquiries through Email, Facebook, LinkedIn, Twitter, etc.

A wide selection of journals (inclusive of 9 subjects, more than 200 journals)

Providing 24-hour high-quality service

User-friendly online submission system

Fair and swift peer-review system

Efficient typesetting and proofreading procedure

Display of the result of downloads and visits, as well as the number of cited articles

Maximum dissemination of your research work

Submit your manuscript at: <http://papersubmission.scirp.org/>

Or contact [ae@scirp.org](mailto:ae@scirp.org)