


ISSN: 2161-6728

Volume 10, Number 1, January 2020



Advances in Molecular Imaging



ISSN: 2161-6728



9 772161 672006



01

<https://www.scirp.org/journal/ami>

Journal Editorial Board

ISSN 2161-6728 (Print) ISSN 2161-6752 (Online)

<https://www.scirp.org/journal/ami>

Editor-in-Chief

Prof. Orhan Nalcioglu

University of California-Irvine, USA

Editorial Board

Prof. Gjumrakch Aliev

University of Atlanta, USA

Dr. Ying Bai

Heart Flow Inc., USA

Prof. Baowei Fei

Emory University, USA

Dr. Zhong-Ping Feng

University of Toronto, Canada

Prof. Richard Hans Gomer

Texas A&M University, USA

Prof. Gultekin Gulsen

University of California-Irvine, USA

Prof. Mohammad Mojammel Al Hakim

University of Southampton, UK

Dr. Maria Kempe

Lund University, Sweden

Dr. David Sigmund Liebeskind

University of California, USA

Dr. Stefan Lorkowski

Friedrich Schiller University, Germany

Prof. Kenneth Maiese

UMDNJ-New Jersey Medical School, USA

Prof. Adalberto Merighi

University of Torino, Italy

Prof. Jean-Pierre Raufman

University of Maryland, USA

Prof. Gianfranco Risuleo

Sapienza University of Rome, Italy

Prof. Steven Alan Rosenzweig

Medical University of South Carolina, USA

Prof. Phillip Ruiz

University of Miami School of Medicine, USA

Dr. Joy Sinha

ANDalyze, Inc., USA

Prof. Maurizio Sorice

Sapienza University of Rome, Italy

Prof. Lun-Quan Sun

Central South University, China

Prof. Bin Tean Teh

National Cancer Center, USA

Prof. Masakazu Toi

Kyoto University, Japan

Prof. Horst Christian Weber

Boston University, USA

Dr. Jia Lin Yang

University of New South Wales, Australia

Table of Contents

Volume 10 Number 1

January 2020

Added Value of SPECT-CT Imaging in the Diagnosis of Unilateral Active Mandibular Hypercondylia in Adult: A Case Report and Review

F. Fokoue, S. El Mselmi, N. Abaouz, N. Ismaili Alaoui.....1

Advances in Molecular Imaging (AMI)

Journal Information

SUBSCRIPTIONS

The *Advances in Molecular Imaging* (Online at Scientific Research Publishing, <https://www.scirp.org/>) is published quarterly by Scientific Research Publishing, Inc., USA.

Subscription rates:

Print: \$59 per issue.

To subscribe, please contact Journals Subscriptions Department, E-mail: sub@scirp.org

SERVICES

Advertisements

Advertisement Sales Department, E-mail: service@scirp.org

Reprints (minimum quantity 100 copies)

Reprints Co-ordinator, Scientific Research Publishing, Inc., USA.

E-mail: sub@scirp.org

COPYRIGHT

Copyright and reuse rights for the front matter of the journal:

Copyright © 2020 by Scientific Research Publishing Inc.

This work is licensed under the Creative Commons Attribution International License (CC BY).

<http://creativecommons.org/licenses/by/4.0/>

Copyright for individual papers of the journal:

Copyright © 2020 by author(s) and Scientific Research Publishing Inc.

Reuse rights for individual papers:

Note: At SCIRP authors can choose between CC BY and CC BY-NC. Please consult each paper for its reuse rights.

Disclaimer of liability

Statements and opinions expressed in the articles and communications are those of the individual contributors and not the statements and opinion of Scientific Research Publishing, Inc. We assume no responsibility or liability for any damage or injury to persons or property arising out of the use of any materials, instructions, methods or ideas contained herein. We expressly disclaim any implied warranties of merchantability or fitness for a particular purpose. If expert assistance is required, the services of a competent professional person should be sought.

PRODUCTION INFORMATION

For manuscripts that have been accepted for publication, please contact:

E-mail: ami@scirp.org

Added Value of SPECT-CT Imaging in the Diagnosis of Unilateral Active Mandibular Hypercondylia in Adult: A Case Report and Review

F. Fokoue*, S. El Mselmi, N. Abaouz, N. Ismaili Alaoui

Faculty of Medicine and pharmacy, University Sidi Ben Abdellah, Fez, Morocco

Email: *fabricefokoue2017@gmail.com

How to cite this paper: Fokoue, F., El Mselmi, S., Abaouz, N. and Ismaili Alaoui, N. (2020) Added Value of SPECT-CT Imaging in the Diagnosis of Unilateral Active Mandibular Hypercondylia in Adult: A Case Report and Review. *Advances in Molecular Imaging*, 10, 1-5.

<https://doi.org/10.4236/ami.2020.101001>

Received: January 12, 2020

Accepted: January 26, 2020

Published: January 29, 2020

Copyright © 2020 by author(s) and Scientific Research Publishing Inc.

This work is licensed under the Creative Commons Attribution International License (CC BY 4.0).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

Abstract

Planar bone Scintigraphy has been frequently used to assess the active or inactive nature of mandibular hypercondylia. We report here the case of a patient whose diagnosis of active mandibular hypercondylia was retained after a complement with SPECT/CT imaging, the planar bone scan being negative. This case confirms the use of SPECT/CT as the examination of choice for full assessment of mandibular hypercondylia in adults. Using bone Scintigraphy SPECT/CT improves both sensitivity and specificity of diagnosing this condition.

Keywords

Hypercondylia, Mandibular Condylar Hyperplasia, Bone Scan, SPECT CT

1. Introduction

Mandibular hypercondylia is a rare and autonomous bone disease due to the excessive growth of the condylar cartilage induced seemingly by a disruption of the feedback loops between the different histological layers leading to a persistence of the multiplication of prechondroblastic cells. Mandibular hypercondylia affects the overall development of the mandible and this condition is characterized clinically by atypical facial asymmetry, with a pronounced mandible anatomy, a disturbed occlusion, and a tilted occlusal plane [1]. Planar bone scintigraphy has been frequently used to assess the active or inactive nature of this situation [2] [3].

2. Patient and Method

The patient is a 43-year-old female, with no significant pathological history, who has had mandibular asymmetry for 10 years. She is directed to the nuclear medicine department for a scintigraphic evaluation of condylar activity following the diagnosis of left mandibular hypercondylia, which was revealed after completing dental panoramic examination supplemented by a craniofacial diagnostic CT scan.

Bone scintigraphy was completed in two stages: Acquisition of early images, followed by late static images focusing on the skull along with the anterior and posterior face. A tomoscintigraphy scan then focused on the skull and was coupled with a low dose CT scan (SPECT/CT) between 0 - 10 min, then 2 hours after IV injection of 740 MBq (20 mCi) of metastable Technetium 99-labeled Methylene Diphosphonate (MDP-99mTC).

The acquisition of images was performed with a Siemens gamma camera model Symbia T6 2010. The analysis was done using a visual and quantitative approach. A relative bone fixation of 55% (or more) was considered pathological.

3. Results

The planar bone scintigraphy did not show any asymmetry with radio tracer uptake in the early times (vascular and tissue), nor any abnormality in bone fixation in the late stage between the two mandibular condyles (**Figure 1**).

The complement by SPECT CT revealed an asymmetry of visual and quantitative fixation depending on the left mandibular condyle with an enlargement of left condyle in transversal, coronal and sagittal reconstruction images respectively thus allowing to conclude to an active left unilateral mandibular hypercondylia (**Figure 2**).




Figure 1. Planar bone scan: Static images focused on the skull performed between 0 - 10 min, then 2 hours after IV injection of 740 MBq (20 mCi) of MDP-Technetium 99 m.




Figure 2. SPECT CT focused on the pathological area. The complement by SPECT CT revealed an asymmetry of visual and quantitative fixation depending on the left mandibular condyle with an enlargement of left condyle.

4. Discussion

Mandibular hypercondylia was first described by Robert Adams in 1836, while describing a case of rheumatoid arthritis [4]. It is a disease of the temporal-mandibular joint resulting in excessive mandibular growth, most often unilateral. It affects the overall development of the mandible and results clinically in an unsightly and functional problem.

Mandibular hypercondylia usually occurs between 10 and 30 years, with a female-dominated sex ratio of 69.6% according to Vernex-Boukerma. Cases between the ages of 9 and 80 have been reported in the literature, and this discrepancy could be explained by delays in diagnosis [1] [4].

The exact pathophysiology of mandibular hypercondylia is unknown, but several etiological factors have been reported, including hyper remodelling of post traumatic mandibular condyle, joint infection, hormonal disorders, and hypervascularization of the mandibular condyle, intrauterine changes and genetic factors [5].

Although the diagnosis can be formally established on histopathological exam, the radiological assessment following clinical examination remains essential to the diagnostic orientation. Information provided from a dental panorama and teleradiographic assessment, or diagnostic CT scan mostly allow the diagnosis of mandibular hypercondylia to be established.

The place of bone scintigraphy has a different interest from other imaging methods because it allows assessing the osteoblastic activity of the condylar cartilage [2] [3]. In a context of clinical and radiological diagnosis of hypercondylia, a positive bone scan shows the active nature of the disease and allows therapy to be adapted.

The molecular imaging techniques commonly used are planar bone scintigraphy and single photon emission computerized tomography (SPECT) which can be analysed quantitatively and qualitatively [5].

For most authors, SPECT can guide surgical management [6] and for others SPECT/CT could be used to measure residual activity after a condylectomy [7]. Condylectomy consists of a surgical excision of the cartilaginous cap, seat of the prechondroblastic cells, resulting in a definitive cessation of the pathological process. It is a key stage in the management of active hypercondylia [8].

In our study, all the modalities of bone scintigraphy were used. Planar imagery was negative at early and late stages but the complement by SPECT/CT made it possible due to fusion images to correlate the hyperfixation area of the tracer with the area of hyperplasia on the CT scans, increasing thus the sensitivity of the bone scintigraphy [9] [10] allowing the retained diagnosis of active unilateral mandibular hypercondylia.

5. Conclusion

This case confirms the fact that the SPECT/CT should be the examination of choice for the assessment of mandibular hypercondylia in adult; it improves both the sensitivity and the specificity of bone scintigraphy with a low irradiation rate and good value for money.

Acknowledgements

To the staff of the nuclear medicine service of Teaching Hospital Hassan 2 of Fez, Morocco.

Ethics Approval and Consent

Obtained by the ethics committee of the Teaching Hospital Hassan II and the Faculty of Medicine and Pharmacy of Fez.

Consent for Publication

Obtained from the patient.

Availability of Data and Material

Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

Funding

No funding received.

Author's Contributions

FF conceived of the study, participated in its design and wrote the manuscript. SM and NA participated in collecting personal data of the patient. NIA supervised the writing of this article. All the authors read and approved the final manuscript.


Conflicts of Interest

The authors declare that they have no competing interest regarding this article.

References

- [1] Gordeff, A., Mercier, J.M. and Delaire, J. (1988) L'hypercondylie mandibulaire. Ses différents aspects cliniques et son traitement. *Acta Stomatologica Belgica*, **85**, 259-276.
- [2] Regaieg, B.F.M., Charfi, H., Sfar, R., Melki, S., *et al.* (2015) Hypercondylie unilatérale: Valeur de la scintigraphie osseuse et de ses différents protocoles d'acquisition. *Médecine Nucléaire*, **39**, 32. <https://doi.org/10.1016/j.mednuc.2015.03.094>
- [3] Choukry, S., Benouhoud, J., Zouine, S., Nadi, S. and Guensi, A. (2019) Apport de la scintigraphie osseuse dans les hypercondylies. *Médecine Nucléaire*, **43**, 209. <https://doi.org/10.1016/j.mednuc.2019.01.102>
- [4] Salagnac, J.M. (2002) Les hypercondylies mandibulaires: Éléments de diagnostic différentiel. *Revue d'Orthopédie Dento-Faciale*, **36**, 75-83. <https://doi.org/10.1051/odf/2002002>
- [5] Nicot, R., Raoul, G. and Ferri, J. (2019) Hypercondylies. EMC Chirurgie orale et maxillo-faciale. Vol. 14.
- [6] Wen, B., Shen, Y. and Wang, C.Y. (2014) Clinical Values of ^{99m}Tc-MDP SPECT Bone Scintigraphy in the Diagnosis of Unilateral Condylar Hyperplasia. *Scientific World Journal*, **2014**, Article ID: 256256. <https://doi.org/10.1155/2014/256256>
- [7] Pripatnanont, P., Vittayakittipong, P., Marmanee, U., *et al.* (2005) The Use of SPECT to Evaluate Growth Cessation of the Mandible in Unilateral Condylar Hyperplasia. *International Journal of Oral and Maxillofacial Surgery*, **34**, 364-368. <https://doi.org/10.1016/j.ijom.2004.11.002>
- [8] Delaire, J. (1977) Le traitement des hypercondylies mandibulaires. Plaidoyer pour la condylectomie. *AOS*, **117**, 29-45.
- [9] Hodder, S.C., Ress, J.I., Olivier, T.B., *et al.* (2000) SPECT Bone Scintigraphy in the Diagnosis and Management of Mandibular Condylar Hyperplasia. *British Journal of Oral and Maxillofacial Surgery*, **38**, 87-93.
- [10] Lopez, B.D.F. and Corral, S.C.M. (2016) Comparison of Planar Bone Scintigraphy and Single Photon Emission Computed Tomography for Diagnosis of Active Condylar Hyperplasia. *Journal of Cranio-Maxillofacial Surgery*, **44**, 70-74. <https://doi.org/10.1016/j.jcms.2015.10.026>

Call for Papers



Advances in Molecular Imaging (AMI)

ISSN 2161-6728 (Print) ISSN 2161-6752 (Online)

<https://www.scirp.org/journal/ami>

Advances in Molecular Imaging is a peer-reviewed, open access journal that publishes original research articles, review articles, case reports, and clinical studies in all areas of molecular imaging:

Editor-in-Chief

Prof. Orhan Nalcioglu

University of California-Irvine, USA

Editorial Board

Prof. Gjumrakch Aliev
Dr. Ying Bai
Prof. Baowei Fei
Dr. Zhong-Ping Feng
Prof. Richard Hans Gomer
Prof. Gultekin Gulsen
Prof. Mohammad Mojammel Al Hakim
Dr. Maria Kempe
Dr. David Sigmund Liebeskind
Dr. Stefan Lorkowski
Prof. Kenneth Maiese
Prof. Adalberto Merighi

Prof. Jean-Pierre Raufman
Prof. Gianfranco Risuleo
Prof. Steven Alan Rosenzweig
Prof. Phillip Ruiz
Dr. Joy Sinha
Prof. Maurizio Sorice
Prof. Lun-Quan Sun
Prof. Bin Tean Teh
Prof. Masakazu Toi
Prof. Horst Christian Weber
Dr. Jia Lin Yang

Subject Coverage

Chemical Synthesis Platform Technologies
Computer Vision and Image Understanding
Contrast Media & Molecular Imaging
Imaging & Microscopy
Imaging Decisions MRI
Imaging Systems and Technology
ImmunoPET and ImmunoSPECT
In Vitro Molecular Diagnostics (IVMD)
Molecular Diagnostics in Cancer and Immune Disorders

Molecular Imaging and Biology
Molecular Imaging Using ImmunoPET
MRI/MDCT/PET Imaging of Various Organs
Neuroimaging
Nuclear Medicine and Molecular Imaging
Onco-Imaging
Preclinical Imaging Systems
Systems Biology
Tracer and Pharmacokinetic Modeling

Notes for Intending Authors

We are also interested in: 1) Short reports—2-5 page papers in which an author can either present an idea with a theoretical background but has not yet completed the research needed for a complete paper or preliminary data; 2) Book reviews—Comments and critiques.

Website and E-Mail

<https://www.scirp.org/journal/ami>

E-mail: ami@scirp.org

What is SCIRP?

Scientific Research Publishing (SCIRP) is one of the largest Open Access journal publishers. It is currently publishing more than 200 open access, online, peer-reviewed journals covering a wide range of academic disciplines. SCIRP serves the worldwide academic communities and contributes to the progress and application of science with its publication.

What is Open Access?

All original research papers published by SCIRP are made freely and permanently accessible online immediately upon publication. To be able to provide open access journals, SCIRP defrays operation costs from authors and subscription charges only for its printed version. Open access publishing allows an immediate, worldwide, barrier-free, open access to the full text of research papers, which is in the best interests of the scientific community.

- High visibility for maximum global exposure with open access publishing model
- Rigorous peer review of research papers
- Prompt faster publication with less cost
- Guaranteed targeted, multidisciplinary audience



**Scientific
Research
Publishing**

Website: <https://www.scirp.org>

Subscription: sub@scirp.org

Advertisement: service@scirp.org