

Effect of Modified Atmosphere Packaging (MAP) on the Stability of Anthocyanins and Degradation of Phenolic Compounds during Postharvest Storage of Pomegranate Fruit

Elham Lashkari

Advanced Academic Programs, Johns Hopkins University, Washington, D.C., USA Email: lashkari3lham@gmail.com

How to cite this paper: Lashkari, E. (2022) Effect of Modified Atmosphere Packaging (MAP) on the Stability of Anthocyanins and Degradation of Phenolic Compounds during Postharvest Storage of Pomegranate Fruit. *Food and Nutrition Sciences*, **13**, 316-335.

https://doi.org/10.4236/fns.2022.133024

Received: February 24, 2022 **Accepted:** March 25, 2022 **Published:** March 28, 2022

Copyright © 2022 by author(s) and Scientific Research Publishing Inc. This work is licensed under the Creative Commons Attribution International License (CC BY 4.0).

http://creativecommons.org/licenses/by/4.0/

Abstract

Effects of storage temperature, packaging material, elevated CO₂ and reduced O₂ contents in the package headspace, and their interaction effects on the total content of anthocyanins and degradation of phenolic compounds during long storage of "Malase Torsh Saveh" pomegranate were studied. The study findings showed that only storage temperature and its duration had significant effects on the degradation of phenolic compounds. The content of anthocyanins was affected by the single effect of storage time and its 3-way interaction effect with the storage temperature and the fruit's surrounding gas composition. Higher contents of both total anthocyanins and browning pigments were recorded at 2°C storage temperature. The Browning Index (BI) of the pomegranate extracts showed to change quadratically with the variations in the total anthocyanins content but linearly with the variations in the content of browning compounds. Thermal dipping treatment and its interaction with the storage time also had significant effect on the BI values of the extracts obtained from modified atmosphere packaged pomegranates stored at 6°C.

Keywords

Pomegranate, Anthocyanins, Phenolic Compounds, Postharvest Storage, Packaging

1. Introduction

The role of phenolic compounds in food quality, their health-promoting and therapeutic effects, along with their mechanism of actions on human health has been the topic of numerous researches in recent years. In addition to their health beneficial properties, phenolic compounds are connected with the quality of plant-derived foods such as appearance, color, taste and aroma. Pomegranate fruit (*Punica granatum L.*) is quite well-known for its nutritious nature and presenting a broad range of health beneficial activities, in particular antioxidative, anti-inflammatory, anti-carcinogenic, antimicrobial, neuro- and cardio-protective properties [1]-[6].

Most of these biological properties are attributed to the pomegranate's primary and secondary metabolites in specific phenolic compounds such as anthocyanins, flavanols, ellagi- and gallotannins, proanthocyanidins, and lignans [5] [7] [8] [9] [10]. These phenolic compounds are found in all parts of the fruit, including fruits peel (ellagitannins, phenolic acids, flavonoids), arils (flavonoids, hydrolysable and condensed tannins, and phenolic acids), seeds (phenolic acids, flavonoids, hydrolysable tannins), and membrane walls (mostly ellagitannins) [11] [12] [13] [14] [15]. Pomegranate's phenolic compounds possess very diverse chemical structures and appear in different quantities depending on the plant cultivar, geographical, agronomic, and other environmental factors including postharvest handling and storage conditions [16] [17] [18] [19].

Scientific findings have provided evidence that the type of technologies and treatments being applied during postharvest handling and storage have significant impact on the total contents of phenolic compounds, their biosynthesis, metabolism, biological activities and the involved enzymatic reactions [16] [20] [21] [22] [23]. These effects could be either favorable or unfavorable towards preserving the desired quality of the pomegranate fruit [16]. Therefore, current studies are also focused to find and develop technologies or a combination of them that allow maintaining the original quality of the fresh fruit, *i.e.*, color, flavor, texture and its nutritious compounds including vitamins, sugars, amino acids, antioxidants, and other health-promoting compounds, while avoiding weight loss, fungal decay, and common physiological disorders such as husk scald, pitting, and chilling injury during cold storage.

Various studies have shown that modified atmosphere packaging (MAP) technology by decreasing the oxygen content and/or increasing the carbon dioxide content of the atmosphere surrounding the fresh produce, can delay quality loss, tissue softening, enzymatic activities, and incidence of various physiological disorders and pathogenic developments during long storage of pomegranate fruit [20] [21] [23]-[31]. MAP can also prevent the fruit weight loss and shriveling by creating a higher relative humidity in the surrounding environment of the fruit. Therefore, MAP could be applied or combined with other storage treatments, for instance with various thermal treatments or refrigeration to inhibit potential development of any decay or physiological disorders [16] [32].

Nonetheless, identifying the applicable MAP that obtains the desired outcome in terms of maintaining the quality and nutritional aspects while avoiding decay and unwanted physiological disorders during long storage requires extensive research studies. The optimum MAP can vary for one pomegranate to another depending on the cultivar, packaging material, and other storage conditions in particular temperature [27]. Within this context, the aims of this study were 1) to determine the effects of MAP technology on the total contents of anthocyanins and degradation of phenolic compounds during cold storage of pomegranate fruit cv. "Malase Torsh Saveh", and 2) to investigate the effect of pre-storage thermal dipping (TD) as a complementary treatment along with MAP technology on the stability of anthocyanins and degradation of phenolic compounds during cold storage of pomegranates. This research is the first study that investigates the effects of MAP technology on the phenolic compounds of an Iranian pomegranate (*Cv. Malas e Torsh Saveh*) during long storage.

2. Materials and Methods

2.1. Materials

2.1.1. Plant Material

For this study, "Malas e Torsh Saveh" pomegranates, which is a popular commercial pomegranate cultivar in Iran, were harvested from Saveh Pomegranate Research Station on October 23^{rd} . Harvested fruits were transported to the University of Tehran's College of Agriculture the same day and stored overnight in one of the research cold rooms at 4°C - 5°C. Pomegranates with uniform size, shape, appearance, and sound outer skin were sorted for the study the day after.

2.1.2. Chemicals and Other Materials

Unperforated low-density polyethylene (LDPE) and polypropylene (PP) packaging bags were purchased from a local provider. Citric acid, sodium citrate, and sodium hydroxide were procured from Merck KGaA (Darmstadt, Germany).

2.2. Methods

2.2.1. Sample Preparation

For the purpose of effect screening, selected pomegranates were divided into different groups with two storage temperatures of $2^{\circ}C \& 6^{\circ}C$, two packaging materials of unperforated PP and LDPE, and two modified headspace gas compositions coded as GC1 and GC2 with elevated contents of CO₂ and reduced O₂ compare to the ambient atmosphere. The two control groups consist of 60 coded pomegranate fruits with no polymeric packaging were evenly placed inside wooden trays and stored at the ambient atmosphere of $2^{\circ}C \& 6^{\circ}C$ cold rooms with 85% - 90% RH throughout the study (Figure 1).

The one-chamber table-top packaging machine (Henkelman 200A, Denmark) was used to flush and insert the programmed gas mixtures inside the unperforated LDPE and PP bags. The two gas mixture compositions that applied prior the storage phase coded as GC1 with 5% $CO_2 + 5\% O_2$ and GC_2 with 10% $CO_2 + 5\% O_2$ gas composition respectively. Nitrogen gas made the remaining component of both mixtures. All sealed packages that each contains a pair of pomegranates



Figure 1. Control and modified atmosphere (MA) packaged pomegranates remained at one of the research cold rooms by the end of study.

were checked, and then transferred and stored in the 2°C and 6°C cold rooms for the long storage study.

2.2.2. Thermal Dipping Treatment

To investigate the potential effects of thermal dipping treatment on the total anthocyanins and phenolic compound contents of fruits packaged with MA during long storage, 60 pomegranates were selected and divided into two groups. Half of these fruits were dipped in the 46°C water bath (Tattnauer, NY, USA) for 2 minutes and left to completely air-dry prior to packaging. After they completely dried, they were coded and placed inside the unperforated LDPE bags in pairs (2 pomegranates in each package). The package headspace of both sampling groups was replaced with the gas composition of 5% CO₂ + 5% O₂ (GC1) and got sealed immediately. Both groups were stored at the cold room of 6°C for a duration of 12 weeks.

2.2.3. Quantification of Total Anthocyanins and Phenolic Compounds' Degradation Products

To measure the total anthocyanin contents and phenolic compound degradation products (or the browning compounds) during storage, first three packages from each sampling group were randomly selected and one pomegranate from each of these packages separated for the study. Also, three pomegranates were randomly selected from each tray of control groups. To obtain the juice, the fruits were half-cut with a clean knife and the juice from the arils was extracted by using a manual stainless-steel pomegranate squeezer. Next the extracted juices were centrifuged (Andreas Hettich centrifuge, Germany) for 20 min at 3000 rpm and the supernatant of the samples were used for the spectrophotometry measurements. For maximum stability of anthocyanins during measurements, the juice supernatants were diluted with citric acid buffer (pH = 3.4) at 1:3 (v/v) ratio.

Later, the total anthocyanin contents of the extracts were recorded by measuring the absorbance at 510 nm with the UV/Vis spectrophotometer instrument (Perkin Elmer Lambda EZ 201, MA, USA) and the recorded values were reported as the Absorbance Unit (AU), which is also expressed as the optical density (OD) per unit length (Equation (1)); the cuvette thickness used in this study was 1 cm:

$$OD_{\lambda} = \frac{A_{\lambda}}{l} \tag{1}$$

where:

l = the distance that light travels through the sample (the sample thickness) measured in cm;

 A_{λ} = absorbance at wavelength λ .

Absorbance at wavelength of 446 nm was recorded by the same spectrophotometer as an indication of polymerization and degradation of phenolic compounds or production of browning compound. Citric acid buffer (pH = 3.4) was used as the reference solution for the absorbance measurements at both 510 and 446 nm wavelengths [20] [21] [25] [26]. All measurements were performed in triplicate.

2.2.4. Browning Index (BI) Calculation

Browning Index of the extracted juice, which reflects the ratio of the total anthocyanin to the development of browning compound or phenolic degradation products was calculated for all sampling groups from Equation (2) [33].

$$BI = \frac{Absorbance at 446 \text{ nm}}{Absorbance at 510 \text{ nm}}$$
(2)

2.2.5. Determination of pH, Titratable Acidity (TA), & Total Soluble Solids (TSS)

4 grams of the extracted juice was diluted with 20 mL of distilled water, and values of pH were measured by the pH meter (744 HannaTM porable pH meter, Portugal) according to Hess-Pierce & Kader, 2003 [26]. Titratable acidity was determined by titrating the diluted juice with 0.1 N NaOH to pH = 8.1 - 8.3 and results were reported as (%) or gr Citric acid per 100 gr of juice. The TSS of the extracted juice was measured by using a refractometer (Bellingham + Stanley, U.K.) at 20°C.

2.2.6. Statistical Analysis

Storage temperature with 2 levels (2°C and 6°C), Packaging type with 3 levels (LDPE, PP, and no packaging or None), head space gas composition with 3 levels (air or ambient atmosphere, GC1, & GC2), and storage time with 4 levels (3 weeks intervals) were applied as the experimental design factors. The effect of hot-water dipping pretreatment on the MA packaged pomegranate fruits stored at 6°C was studied on 2 groups for 12 weeks and analyzed separately. All measurements were conducted in 3 replicates and the mean value was reported. JMP Pro software was used to perform the statistical analysis and data visualization.

Analysis of variance (ANOVA) at p < 0.05 and Student's t Test (for mean comparison) were applied when appropriate.

3. Results

3.1. Changes in the Total Content of Anthocyanins

Among the multiple factors being studied, only the single effect of storage time was found to have a significant effect on the total content of anthocyanins at p < p0.01 during this study. The results showed a decline from the average absorbance value of 0.917 \pm 0.054 (Mean \pm SE) at the beginning of the study through the 7th week of the storage followed by a slight increase towards the end of the study (0.49 ± 0.054) . The total content of anthocyanins was also affected by the 3-way interaction effects of the storage duration, storage temperature, and the surrounding gas composition. Overall, samples with MA packaging displayed lower values of total anthocyanins around the seventh week of their storage compared to the control samples with no packaging (Figure 2). Among the MA packaged treatments, storage temperature at 2°C, LDPE packaging, and the initial gas mixture of 5% CO_2 + 5% O_2 (GC1) were associated with higher amounts of total anthocyanin contents by the end of study. As illustrated in Figure 3, the interaction profile also indicates an interaction between the storage temperature and the type of packaging although it was not found to be statistically significant. Besides, higher amounts of anthocyanins were observed on average in the extracts of samples that were stored at 2°C storage temperature compared to samples that were stored at 6°C during long storage regardless of their packaging type and their surrounding atmospheric composition.

3.2. Changes in the Degradation of Phenolic Compounds

The results of this study showed that the single effects of storage time and temperature have had significant effects on the degradation of phenolic compounds and formation of browning pigments at p < 0.05. On average higher amount of phenolic compounds degradation products was observed by the end of study (0.332 ± 0.024) compared to the beginning of the study (0.278 ± 0.024) throughout the 10-week storage time. Likewise, degradation of phenolic compounds was enhanced at 2°C storage temperature with the average absorbance value of 0.323 \pm 0.017 at 446 nm compared to the corresponding values of 0.264 \pm 0.017 when samples were stored at 6°C temperature (Figure 4). None of the 2 & 3-way interaction effects was found to be significant. Yet, the interaction profile (Figure 5) displayed an interaction between the storage temperature and the type of packaging that may not be statistically significant. As shown in Figure 5, storage temperature of 2°C, UPP packaging, and the initial gas mixture of 5% CO_2 + 5% O_2 (GC1) induced higher amounts of degradation of phenolic compounds and subsequent formation of browning pigments among pomegranate fruits with MA packaging. Extracts of pomegranates that were stored with no packaging and surrounded by the air or ambient atmosphere of the cold room



Figure 2. Effects of packaging type (top) and gas compositions (bottom) on the total anthocyanins content during the 10 weeks of storage. Each error bar is constructed using 1 standard error from the mean. Packaging type: PP, polypropylene; LDPE, low-density polyethylene; NONE, no packaging is used. Gas composition: NRM, air or ambient atmosphere; GC1, initial MAP with 5% CO₂ and 5% O₂; GC2, initial MAP with 10% CO₂ and 5% O₂.

showed the least amounts of browning pigments on average at the end of study (Figure 4, bottom).

3.3. Changes in the Browning Index (BI)

The results of the data analysis indicate that the Browning Index was affected by the single effect of the storage time, its 2-way interaction effects with the type of packaging, and its 3-way interaction effects with the storage temperature and the surrounding atmospheric composition at p < 0.01. The average BI value was increased from 0.31 ± 0.02 at the beginning of the study to 0.69 ± 0.04 by the end



Figure 3. Interaction profile of temperature, packaging type, and gas composition effects on the total anthocyanins content of the MA packaged pomegranates. Packaging type: PP, polypropylene; LDPE, low-density polyethylene. Gas composition: GC1, initial MAP with 5% CO₂ and 5% O₂; GC2, initial MAP with 10% CO₂ and 5% O₂.

of study. The increase occurred during the first 4 weeks of the storage and remained around the same value during the rest of the 10-week study. Extracts of pomegranates with no packaging that were surrounded by the ambient atmosphere during storage displayed the lowest values of BI (0.53 ± 0.06) at the end of the storage compared to the treatments with MA packaging. Additionally, the extracts of pomegranates with PP packaging showed the highest BI value ($0.63 \pm$ 0.06) at the end of study (**Figure 6**, top). Overall higher BI values were associated with storage temperature at 2°C. Control samples with no packaging that were surrounded by the ambient atmosphere showed higher BI values when stored at 6°C. In contrast to air-stored samples, the extracts of MA packaged pomegranates displayed higher BI values when stored at 2°C cold room (**Figure 6**, bottom).

3.4. Changes in pH, TA, & Total Soluble Solids (TSS)

Storage time showed to have significant effects (at p < 0.01) on the pH values of



Figure 4. Effects of Storage temperature (top) and type of packaging (bottom) on development of browning compounds over 10 weeks of storage. Each error bar is constructed using 1 standard error from the mean. Packaging type: PP, polypropylene; LDPE, low-density polyethylene; NONE, no packaging is used.

the treatments over the course of study. The pH values displayed an increase from 3.45 ± 0.05 at the harvest to 3.67 ± 0.17 during the first month of storage that remained around the pH value of 3.64 ± 0.19 till the end of 10 weeks of storage. pH was not influenced significantly by the storage temperature; however, it was affected by the 2- & 3-way interaction effects of storage time with packaging type & the surrounding gas composition (p < 0.01). Overall, the control samples that were surrounded by the ambient atmosphere and had no packaging showed pH values closer to the average pH value at the harvest (pH = 3.53 ± 0.03). Whereas extracts from pomegranates that had unperforated PP packaging and



Figure 5. Interaction profile of temperature, packaging type, gas composition effects on the development of phenolic compounds degradation products of MA packaged pomegranates. Packaging type: PP, polypropylene; LDPE, low-density polyethylene. Gas composition: GC1, initial MAP with 5% CO₂ and 5% O₂; GC2, initial MAP with 10% CO₂ and 5% O₂.

packaged with the initial gas mixture of 5% $CO_2 + 5\% O_2$ (GC1) were associated with higher values of pH at the end of 10 weeks of storage with the average pH values of 3.66 ± 0.03 and 3.63 ± 0.02 respectively (**Figure 7**, top).

TA values were significantly affected by the single effect of gas composition and its interaction with the storage duration (p < 0.01). The average TA values displayed slight decrease from the average TA value at the harvest (1.02% ± 0.032%) until the 7th week of storage (0.918% ± 0. 032%) and then started to slowly increase toward the end of the study (0.974% ± 0.032%). As shown in **Figure 7** (bottom), the control samples that were surrounded by the ambient atmosphere and had no packaging showed higher amounts of TA, while the samples with PP packaging and the initial gas mixture of 5% CO₂ + 5% O₂ (GC1) showed the lowest values for TA by the end of study. These results are in agreement with the study findings on pH changes throughout the storage. Also, pomegranates that were packaged with the initial gas mixture of 10% CO₂ + 5% O₂ (GC2) exhibited the lowest pH values (pH = 3.5) and the highest TA (TA =



Figure 6. Effects of (top) packaging type and storage temperature, and (bottom) storage temperature, time, and gas composition on the values of Browning Index (BI) during 10 weeks of storage. Each error bar is constructed using 1 standard error from the mean. Packaging type: PP, polypropylene; LDPE, low-density polyethylene; NONE, no packaging is used. Gas composition: NRM, air or ambient atmosphere; GC1, initial MAP with 5% CO₂ and 5% O₂; GC2, initial MAP with 10% CO₂ and 5% O₂.

1.19%) values by the end of study. These findings suggest the potential slow penetration of CO_2 gas within the pomegranate's tissues and arils during the long storage that might have led to the formation of carbonic acid and the subsequent observation of pH reduction and %TA increase of the pomegranate extracts. These changes were particularity noticeable from the seventh week of storage.

The values of the extracts' total soluble solids (TSS) were affected only by the storage duration at p < 0.05 and none of the other factors *i.e.*, storage temperature, surrounding gas composition, and type of packaging and their 2- and 3-way interactions was found to be significant. The average TSS values showed slight decrease from the average value of 18.17 ± 0.13 at the harvest to 17.57 ± 0.13 by the end of storage, which indicates minor loss of saugers due to the low



Figure 7. Effects of packaging type and surrounding gas composition on the (top) pH, and TA (bottom) of pomegranates during 10 weeks of storage. Each error bar is constructed using 1 standard error from the mean. Packaging type: PP, polypropylene; LDPE, low-density polyethylene; NONE, no packaging is used. Gas composition: NRM, air or ambient atmosphere; GC1, initial MAP with 5% CO₂ and 5% O₂; GC2, initial MAP with 10% CO₂ and 5% O₂.

respiratory activities of pomegranate fruits during the long storage.

3.5. Effects of Thermal Dipping (TD) Treatment

The results show that the single effect of thermal dipping (TD) pretreatment was not significant on the total content of anthocyanins, degradation of phenolic compounds, and Browning Index during the long storage. However, as shown in **Table 1**. The 2-way interaction effect of thermal dipping treatment with the storage time had significant effect at p < 0.01 on the values of the Browning Index. The extracted juice obtained from pomegranates that were subjected to the TD treatment prior to the storage had higher BI values (mean BI = 0.95) compared to the control group (mean BI = 0.66) after 12 weeks of study (**Figure 8**).

The values of pH, TSS, and TA also have not been affected by the single effect of TD treatment. Still, the 2-way interaction effect of thermal dipping treatment with storage time was found to have significant effects on the pH values at p < 0.01, TSS at p < 0.05, and TA at p < 0.05. By the end of 12 weeks of storage, the extracted juice of samples that received TD treatment showed higher pH & lower TA and TSS values compared to the control samples that have not received thermal dipping treatment prior to the storage.

	Thermal Dipping Treatment			Time	Thermal Dipping Treatment * Time			
	With	Without			With		Without	
Absorption at 510 nm (Indicator of the total anthocyanins content)	0.487 ± 0.067a	0.480 ± 0.073a	Wk. 1	0.917 ± 0.052a	Wk. 1	0.917 ± 0.083a	Wk. 1	0.917 ± 0.083a
			Wk. 3	$0.331\pm0.078\mathrm{b}$	Wk. 3	0.464 ± 0.11bc	Wk. 3	$0.197\pm0.017\mathrm{d}$
			Wk. 6	$0.367 \pm 0.046b$	Wk. 6	0.394 ± 0.084bd	Wk. 6	0.340 ± 0.053bd
			Wk. 9	$0.395 \pm 0.098b$	Wk. 9	0.263 ± 0.045cd	Wk. 9	$0.528 \pm 0.1681b$
			Wk. 12	$0.409\pm0.022b$	Wk. 12	0.399 ± 0.041bd	Wk. 12	$0.420 \pm 0.026 bd$
Absorption at		5a 0.244 ± 0.026a	Wk. 1	0.287 ± 0.038a	Wk. 1	0.287 ± 0.061ab	Wk. 1	0.287 0.061ab
446 nm (Indicator of	0.263 ± 0.025a		Wk. 3	$0.175 \pm 0.023b$	Wk. 3	0.196 ± 0.036bd	Wk. 3	0.155 ± 0.032cd
phenolic			Wk. 6	0.161 ± 0.018b	Wk. 6	0.170 ± 0.025bd	Wk. 6	0.151 ± 0.029d
compounds'			Wk. 9	0.318 ± 0.03a	Wk. 9	$0.283\pm0.023ab$	Wk. 9	0.352 ± 0.051a
degradation products)			Wk. 12	0.327 ± 0.034a	Wk. 12	0.379 ± 0.046a	Wk. 12	0.275 ± 0.03abc
Browning Index (BI) of the juice extract	0.651 ± 0.088a	0.595 ± 0.069a	Wk. 1	$0.310 \pm 0.035 d$	Wk. 1	$0.310 \pm 0.055e$	Wk. 1	$0.310 \pm 0.055e$
			Wk. 3	0.626 ± 0.126bc	Wk. 3	0.438 ± 0.034de	Wk. 3	0.814 ± 0.207bc
			Wk. 6	0.442 ± 0.016cd	Wk. 6	0.444 ± 0.03de	Wk. 6	0.440 ± 0.018de
			Wk. 9	0.930 ± 0.116a	Wk. 9	1.112 ± 0.101a	Wk. 9	0.748 ± 0.153bc
			Wk. 12	0.806 ± 0.079ab	Wk. 12	0.950 ± 0.05ab	Wk. 12	0.662 ± 0.088cd

Table 1. Effect of thermal dipping treatment, storage time and their two-way interaction on the contents of total anthocyanins, degradation of phenolic compounds, and browning index (BI) of the juice extracts of the stored MA packaged pomegranates.

Data are presented as the Mean \pm ST.DEV. Comparisons of Means and LSD between pairs were determined by Student's t test (α = 0.05). For each source of effect, levels or means that are not connected by the same letter are significantly different (p < 0.05).



Figure 8. Effects of thermal dipping treatment on the values of Browning Index (BI) during 12 weeks of storage. Each error bar is constructed using 1 standard error from the mean. Dark gray bars represent the sample group with no thermal dipping treatment, and bars in lighter gray represent the sample group that were subjected to the thermal dipping treatment.

4. Discussion

The findings of this study showed an overall decrease in the total contents of anthocyanins during 10 weeks of storage, in which multiple factors might have been involved. Some of the factors that affect anthocyanins stability or their susceptibility to degradation are temperature, increased sugar, pH, acidity level, metal chelation, and copigmentation with other flavonols and tannins [19] [34]. Likewise, regulation and activity of phenylalanine ammonia-lyase (PAL), which is the key enzyme in biosynthesis of flavonoids can be another influential factor [16]. Therefore, the overall decline in the contents of anthocyanins during the storage could be to some extent as the result of the effects of cold storage temperature on the enzymatic activities of PAL that inversely has affected the biosynthesis of anthocyanins. Similarly, the pomegranate extracts that obtained from MA packaged samples showed less absorbance at 510 nm compared to those obtained from air-stored pomegranates, which became even more evident by the seventh week of storage. These findings are consistent with the findings of previous studies on MA packaging of "Mollar de Elche" and "Wonderful" pomegranate cultivars, suggesting elevated amounts of CO₂ in the surrounding atmosphere of stored pomegranates might have a role in disrupting the PAL activity that subsequently can lead to the suppression of anthocyanins biosynthesis and pigmentation [25] [26] [34]. Additionally, pH reduction and TA increase as the result of gradual penetration of CO₂ within the tissues and arils of MA packaged pomegranates and potential formation of carbonic acid could be another contributing factor to the anthocyanins' stability and their noticeable content increase from week 7th of storage towards the end of the 10-week study. So, despite the lack of significant difference between the anthocyanin contents of control and MA packaged pomegranates by the end of study, yet different underlying factors might have been participated throughout the storage.

Nonetheless, a different pattern in the contents of anthocyanins during the cold storage of MA packaged "Hicaznar" and "Hicrannar" cultivars was reported. Selcuk & Erkan (2014) reported an increase in the total anthocyanin contents of "Hicrannar" pomegranates during the first 100 days of storage at 6°C that followed by a decrease over the last 20 days of the study in both MAP and control treatments through 120 days of storage [28]. Total anthocyanin contents in "Assaria" pomegranates that were subjected to several storage treatments also showed an increase during the first month of storage with continued decrease towards the end of 4 months of cold storage at 5°C [22]. "Ruby" pomegranates when air-stored at 5 and 7°C for 16 weeks displayed an increase in their anthocyanin concentration through 8 and 12 weeks of storage that followed by a slight decrease and no change in their content correspondingly till the end of cold storage [35]. Selcuk & Erkan (2015) and Çandir et al. (2018 & 2019) also reported an overall increase in the anthocyanin content of "Hicaznar" pomegranates during the first 4 months of cold storage and a gradual decrease in the anthocyanin content toward the end of storage in both air-stored & MA packaged pomegranates when stored at 6°C for 6 months [23] [30] [31]. Yet, Çandir *et al.* (2018) observed a delayed anthocyanins synthesis and lower anthocyanin concentrations in MA packaged "Hicaznar" pomegranates [30]. The difference in the observed anthocyanins contents of these cultivars during long storage can be attributed to the inherent differences among various cultivars given that the same trend has been recorded for both control and MA packaged pomegranates during long storage that indicates the potential continued postharvest metabolic activities including biosynthesis of anthocyanins in the fruit despite of being refrigerated and/or subjected to the MA packaging.

The current study results also showed higher contents of browning pigments due to the degradation of phenolic compounds when pomegranates stored at 2°C compared to 6°C storage temperature. These findings can be attributed to the enzymatic activities of polyphenol oxidases (PPO) and peroxidases (POD) that catalyzed enzymatic browning in fruit tissues as the results of environmental stresses such as low storage temperatures and mechanical injuries [16]. Likewise, elevated CO₂ contents in the surrounding atmosphere of MA packaged pomegranates can be another contributing factor that enhanced polymerization of phenolic compounds in the tissues and arils of MA packages pomegranates compared to the control (air-stored) samples. These observations are consistent with the findings reported for "Mollar de Elche" pomegranates when packaged with CO₂-enriched MA and stored at 5°C for 12 weeks [25]. The results of this study also recorded higher amounts of browning pigments over the cold storage that can be an indication of PPO and POD enzymatic activities as the results of minor incident of chilling injury in the tissues that often led to development of browning pigments in pomegranates when stored at low temperatures for a long time.

As shown in Figure 9, the variations in the browning index (BI) values during long storage is more influenced by the variation in the total contents of anthocyanins rather than degradation of other phenolic compounds, which is an interesting finding and implies that anthocyanins are the main phenolic compounds in the arils. Likewise, these results indicate minor development of browning pigments as the result of degradation of phenolic compounds other than anthocyanins during 10 weeks of storage that is even more noticeable at 2°C compared to 6°C storage temperature. Storage temperature of 2°C is associated with higher contents of both total anthocyanins and development of browning pigments that imply the potential effect of low temperatures on the gene regulations, expression and physiological activity of the enzymes that are involved in the biosynthesis of phenolic compound such as anthocyanins and those that are participating in the enzymatic browning as the results of the incidence of physiological disorders (e.g. chilling injury) due to environmental stresses such as long storage at low temperatures or elevated CO₂ contents in the surrounding atmosphere of the fruit.

As presented in **Figure 10**, the values of the BI of the extracts change quadratically with the variations in the contents of total anthocyanins but linearly



Figure 9. Profiles of changes in the (top) total anthocyanins, (middle) browning index, and (bottom) development of browning pigments during 10 weeks of storage at 2° C (solid line) and 6° C (dashed line).



Figure 10. (a) Response surface 3D showing changes in the Browning Index (BI) of the extracted juice in relation to the total anthocyanins content (absorbance at 510 nm) and phenolic compound degradation products (absorbance at 446 nm) during storage, and (b) prediction profiler for the anticipated Browning Index.

with the variations in the contents of the browning compounds. Consequently, the prediction profiles can be used to determine the optimum anthocyanin contents and storage conditions based on the desired BI value for the anticipated applications such as fresh consumption, processed food products for instance juices, jams, jellies, or as a value-added ingredient e.g., natural colorant or a bioactive compound.

The study results also revealed that thermal dipping (TD) treatment can affect the BI value of the pomegranate extracts by enhancing degradation of anthocyanins, their polymerization, and subsequent formation of browning compounds throughout the storage. The increase in the BI value of the samples that received TD treatment prior their packaging is more visible since the ninth week of the study. Similarly, the significant 2-way interaction effect of TD with the storage time on the pH, TA, and TSS changes during long storage of MA packaged pomegranates indicates the higher metabolisms of soluble solids, mainly sugars (such as glucose, fructose, arabinose) and organic acids (e.g., citric acid) by the pomegranates that have been subjected to the thermal treatment compared to the control group. Nonetheless, the TSS/TA ratio of pomegranates was not significantly affected (data not shown here) by TD treatment after 12-weeks of storage at 6°C.

5. Conclusion

In summary, the total contents of anthocyanins in "Malase Torsh Saveh" pomegranates declined through 10 weeks of storage in all treatments. Among samples with modified atmosphere (MA) packaging, pomegranates with LDPE packaging and the initial gas composition of 5% CO₂ + 5% O₂ in their package headspace maintained higher contents of total anthocyanins when stored at 2°C. Nonetheless, higher incidence of browning pigments was also recorded at 2°C, possibly as the results of occurrence of minor chilling injury throughout the cold storage. Lesser browning pigments were developed among samples with MA packaging through the first seven weeks of storage, however, their contents started to increase towards the end of study. The browning index (BI) values of the extracts showed an overall increase throughout the storage and they were more influenced by the variations in the content of total anthocyanins in the extracts rather than the content of browning pigments, which indicates minor development of browning pigments as the result of phenolic compounds degradation during long storage. Pre-storage thermal dipping (TD) as a complementary treatment along with MAP showed no significant effects on the total content of anthocyanins, degradation of phenolic compounds, or development of browning compounds during the cold storage. However, BI values of MA packaged pomegranates that had received TD prior to the storage started to significantly increase after 2 months of storage at 6°C.

Acknowledgements

This research was conducted at the Department of Food Science & Technology,

and it was funded by University of Tehran, College of Agriculture and Natural Resources, Karaj, Tehran. The author also thanks Dr. Banafshe Lashkari, PhD, from University of Waterloo for her assistance in article retrieval.

Ethics Statement

The author declares that this research (a) is the author's own original work, (b) has not been published previously, and (c) is not under consideration for publication elsewhere.

Conflicts of Interest

The author declares no known conflicts of interest regarding publication of this paper.

References

- Afaq, F., Saleem, M., Krueger, C.G., Reed, J.D. and Mukhtar, H. (2005) Anthocyanin- and Hydrolysable Tannin-rich Pomegranate Fruit Extract Modulates MAPK and NF-κB Pathways and Inhibits Skin Tumorigenesis in CD-1 Mice. *International Journal of Cancer*, 113, 423-433. <u>https://doi.org/10.1002/ijc.20587</u>
- [2] Reddy, M.K., Gupta, S.K., Jacob, M.R., Khan, S.I. and Ferreira, D. (2007) Antioxidant, Antimalarial and Antioxidant Activities of Tannin-Rich Fractions, Ellagitannins and Phenolic Acids from Punica granatum L. *Planta Medica*, **73**, 461-467. <u>https://doi.org/10.1055/s-2007-967167</u>
- [3] Haghayeghi, K., Shetty, K. and Labbé, R. (2013) Inhibition of Foodborne Pathogens by Pomegranate Juice. *Journal of Medicinal Food*, 16, 467-470. <u>https://doi.org/10.1089/jmf.2012.0233</u>
- [4] Howell, A.B. and D'Souza, D.H. (2013) The Pomegranate: Effects on Bacteria and Viruses That Influence Human Health. *Evidence-Based Complementary and Alternative Medicine*, 2013, Article ID: 606212. <u>https://doi.org/10.1155/2013/606212</u>
- [5] Pepe, G., Rapa, S.F., Salviati, E., Bertamino, A., Auriemma, G., Cascioferro, S., Autore, G., Quaroni, A., Campiglia, P. and Marzocco, S. (2020) Bioactive Polyphenols from Pomegranate Juice Reduce 5-Fluorouracil-Induced Intestinal Mucositis in Intestinal Epithelial Cells. *Antioxidants*, 9, Article No. 699. https://doi.org/10.3390/antiox9080699
- [6] Mukherjee, S., Ghosh, S., Choudhury, S., Gupta, P., Adhikary, A. and Chattopadhyay, S. (2021) Pomegranate Polyphenols Attenuate Inflammation and Hepatic Damage in Tumor-Bearing Mice: Crucial Role of nf-κb and The Nrf2/GSH Axis. *Journal of Nutritional Biochemistry*, **97**, Article ID: 108812. <u>https://doi.org/10.1016/j.jnutbio.2021.108812</u>
- [7] Viuda-Martos, M., Fernández-López, J. and Pérez-Álvarez, J. (2010) Pomegranate and Its Many Functional Components as Related to Human Health: A Review. *Comprehensive Reviews in Food Science and Food Safety*, 9, 635-654. <u>https://doi.org/10.1111/j.1541-4337.2010.00131.x</u>
- [8] Gómez-Caravaca, A.M., Verardo, V., Toselli, M., Segura-Carretero, A., Fernández Gutiérrez, A. and Caboni, M.F. (2013) Determination of The Major Phenolic Compounds in Pomegranate Juices by HPLC-DAD-ESI-MS. *Journal of Agricultural and Food Chemistry*, 61, 5328-5337. https://doi.org/10.1021/jf400684n
- [9] Ambigaipalan, P., de Camargo, A. and Shahidi, F. (2016) Phenolic Compounds of

Pomegranate Byproducts (Outer Skin, Mesocarp, Divider Membrane) and Their Antioxidant Activities. *Journal of Agricultural and Food Chemistry*, **64**, 6584-6604. https://doi.org/10.1021/acs.jafc.6b02950

- [10] Fahmy H., Hegazi, N., El-Shamy, S. and Farag, M.A. (2020) Pomegranate Juice as a Functional Food: A Comprehensive Review of Its Polyphenols, Therapeutic Merits, and Recent Patents. *Food & Function*, **11**, 5768-5781. <u>https://doi.org/10.1039/D0FO01251C</u>
- [11] Ambigaipalan, P., de Camargo, A. and Shahidi, F. (2017) Identification of Phenolic Antioxidants and Bioactives of Pomegranate Seeds Following Juice Extraction Using HPLC-DAD-ESI-MSⁿ. *Food Chemistry*, 221, 1883-1894. <u>https://doi.org/10.1016/j.foodchem.2016.10.058</u>
- [12] Bar-Ya'akov, I., Tian, L., Amir, R. and Holland, D. (2019) Primary Metabolites, Anthocyanins, and Hydrolyzable Tannins in the Pomegranate Fruit. *Frontiers in Plant Science*, **10**, Article No. 620. <u>https://doi.org/10.3389/fpls.2019.00620</u> <u>https://www.frontiersin.org/article/10.3389/fpls.2019.00620</u>
- [13] Díaz-Mula, H.M., Tomás-Barberán, F.A. and García-Villalba, R. (2019) Pomegranate Fruit and Juice (cv. Mollar), Rich in Ellagitannins and Anthocyanins, Also Provide a Significant Content of a Wide Range of Proanthocyanidins. *Journal of Agricultural and Food Chemistry*, **67**, 9160-9167. <u>https://doi.org/10.1021/acs.jafc.8b07155</u>
- [14] Fourati, M., Smaoui, S., Hlima, H.B., Elhadef, K., Ben Braiek, O., Ennouri, K., Mtibaa, A.C. and Mellouli, L. (2020) Bioactive Compounds and Pharmacological Potential of Pomegranate (*Punica granatum*) Seeds—A Review. *Plant Foods for Human Nutrition*, **75**, 477-486. <u>https://doi.org/10.1007/s11130-020-00863-7</u>
- [15] Hegazi, M.N., El-Shamy, S., Fahmy, H. and Farag, A.M. (2021) Pomegranate Juice as a Super-Food: A Comprehensive Review of Its Extraction, Analysis, and Quality Assessment Approaches. *Journal of Food Composition and Analysis*, **97**, Article ID: 103773. https://doi.org/10.1016/j.jfca.2020.103773
- [16] Tomás-barberán, F.A. and Espín, J.C. (2001) Phenolic Compounds and Related Enzymes as Determinants of Quality in Fruits and Vegetables. *Journal of Science of Food and Agriculture*, 81, 853-876. <u>https://doi.org/10.1002/jsfa.885</u>
- [17] Gözlekçi, S., Saraçoğlu, O., Onursal E. and Ozgen, M. (2011) Total Phenolic Distribution of Juice, Peel, and Seed Extracts of Four Pomegranate Cultivars. *Pharmacognosy Magazine*, 7, 161-164. <u>https://doi.org/10.4103/0973-1296.80681</u>
- [18] Li, X., Wasila, H., Liu, L., Yuan, T., Gao, Z., Zhao, B. and Ahmad, I. (2015) Physicochemical Characteristics, Polyphenol Compositions and Antioxidant Potential of Pomegranate Juices from 10 Chinese Cultivars and the Environmental Factors Analysis. *Food Chemistry*, **175**, 575-584. <u>https://doi.org/10.1016/j.foodchem.2014.12.003</u>
- [19] Lee, C.Y. (2018) Fruits and Vegetables. In: de Man, J.M., Finley, J.W., Hurst, W.J. and Lee C.Y., Eds., *Principles of Food Chemistry*, 4th Edition, Springer, Cham, 435-455. <u>https://doi.org/10.1007/978-3-319-63607-8 11</u>
- [20] Gill, M.I., Sánchez, R., Ginés Marín, J. and Artés, F. (1996) Quality Changes in Pomegranates during Ripening and Cold Storage. *Zeitschrift für Lebensmittel-Untersuchung* und Forschung, 202, 481-485. <u>https://doi.org/10.1007/BF01197269</u>
- [21] Holcroft, D.M., Gil, M.I. and Kader, A.A. (1998) Effect of Carbon Dioxide on Anthocyanins, Phenylalanine Ammonia Lyase and Glucosyltransferase in the Arils of Stored Pomegranates. *Journal of American Society of Horticultural Science*, 123, 136-140. <u>https://doi.org/10.21273/JASHS.123.1.136</u>

- [22] Miguel, G., Fontes, C., Antunes, D., Neves, A. and Martins, D. (2004) Anthocyanin Concentration of "Assaria" Pomegranate Fruits during Different Cold Storage Conditions. *BioMed Research International*, **2004**, Article ID: 387670. https://doi.org/10.1155/S1110724304403076
- [23] Selcuk, N. and Erkan, M. (2015) Changes in Phenolic Compounds and Antioxidant Activity of Sour-Sweet Pomegranates cv. 'Hicaznar' During Long-Term Storage Under Modified Atmosphere Packaging. *Postharvest Biology and Technology*. 109, 30-39. <u>https://doi.org/10.1016/j.postharvbio.2015.05.018</u>
- [24] Beaudry, R. (1999) Effect of O₂ and CO₂ Partial Pressure on Selected Phenomena Affecting Fruit and Vegetable Quality. *Postharvest Biology and Technology*, 15, 293-303. <u>https://doi.org/10.1016/S0925-5214(98)00092-1</u>
- [25] Artés, F., Villaescusa, R. and Tudela, J.A. (2000) Modified Atmosphere Packaging of Pomegranate. *Journal of Food Science*, 65, 1112-1116. <u>https://doi.org/10.1111/j.1365-2621.2000.tb10248.x</u>
- [26] Hess-Pierce, B. and Kader, A.A. (2003) Responses of "Wonderful" Pomegranate to Controlled Atmospheres. *Acta Horticulturae*, **600**, 751-757. <u>https://doi.org/10.17660/ActaHortic.2003.600.115</u>
- [27] Caleb, O.J., Opara, U.L. and Witthuhn, C.R. (2012) Modified Atmosphere Packaging of Pomegranate Fruit and Arils: A Review. *Food and Bioprocess Technology*, 5, 15-30. <u>https://doi.org/10.1007/s11947-011-0525-7</u>
- [28] Selcuk, N. and Erkan, M. (2014) Changes in Antioxidant Activity and Postharvest Quality of Sweet Pomegranates cv. *Hicrannar* under Modified Atmosphere Packaging. *Postharvest Biology and Technology*, **92**, 29-36. https://doi.org/10.1016/j.postharvbio.2014.01.007
- [29] Rao, D.V. and Shivashankara, K.S. (2018) Effect of Modified Atmosphere Packaging on the Extension of Storage Life and Quality Maintenance of Pomegranate (cv. "Bhagwa") at Ambient and Low Temperatures. *Journal of Food Science and Technology*, 55, 2103-2113. <u>https://doi.org/10.1007/s13197-018-3125-y</u>
- [30] Çandır, E., Özdemir, A. and Aksoy, M. (2018) Effects of Chitosan and Modified Atmosphere Packaging on Postharvest Quality and Bioactive Compounds of Pomegranate Fruit cv. "*Hicaznar*". *Scientia Horticulturae*, 235, 235-243. <u>https://doi.org/10.1016/j.scienta.2018.03.017</u>
- [31] Çandır, E., Özdemir, A. and Aksoy, M. (2019) Effects of Modified Atmosphere Packaging on the Storage and Shelf Life of Hicaznar Pomegranate Fruits. *Turkish Journal of Agriculture and Forestry*, 43, 241-253.
- [32] Artés, F. and Tomás-Barberán, F.A. (2000) Post-Harvest Technological Treatments of Pomegranate and Preparation of Derived Products. *Options Méditerranéennes. Série A: Séminaires Méditerranéens*, **42**, 199-204.
- [33] Muche, B.M., Speers, R.A. and Rupasinghe, H. (2005) Storage Temperature Impacts on Anthocyanins Degradation, Color Changes and Haze Development in Juice of "Merlot" and "Ruby" Grapes (*Vitis vinifera*). *Frontiers in Nutrition*, **5**, Article No. 100. <u>https://doi.org/10.3389/fnut.2018.00100</u>
- [34] Gill, M.I., Holcroft, D.M. and Kader, A.A. (1997) Changes in Strawberry Anthocyanins and Other Polyphenols in Response to Carbon Dioxide Treatment. *Journal of Agricultural and Food Chemistry*, 45, 1662-1667. <u>https://doi.org/10.1021/jf960675e</u>
- [35] Fawole, O.A. and Opara, U.L. (2013) Effects of Storage Temperature and Duration on Physiological Response of Pomegranate Fruit. *Industrial Crops and Products*, 47, 300-309. <u>https://doi.org/10.1016/j.indcrop.2013.03.028</u>