

Signed Tilings by Ribbon L n-Ominoes, n Odd, via **Gröbner Bases**

Viorel Nitica

Department of Mathematics, West Chester University, West Chester, PA, USA Email: vnitica@wcupa.edu

How to cite this paper: Nitica, V. (2016) Signed Tilings by Ribbon L n-Ominoes, n Odd, via Gröbner Bases. Open Journal of Discrete Mathematics, 6, 297-313. http://dx.doi.org/10.4236/ojdm.2016.64025

Received: July 9, 2016 Accepted: October 11, 2016 Published: October 14, 2016

 \odot

Copyright © 2016 by author and Scientific Research Publishing Inc. This work is licensed under the Creative Commons Attribution International License (CC BY 4.0).

http://creativecommons.org/licenses/by/4.0/ **Open Access**

Abstract

We show that a rectangle can be signed tiled by ribbon L *n*-ominoes, *n* odd, if and only if it has a side divisible by n. A consequence of our technique, based on the exhibition of an explicit Gröbner basis, is that any k-inflated copy of the skewed Ln-omino has a signed tiling by skewed L n-ominoes. We also discuss regular tilings by ribbon L n-ominoes, n odd, for rectangles and more general regions. We show that in this case obstructions appear that are not detected by signed tilings.

Keywords

Polyomino, Replicating Tile, L-Shaped Polyomino, Skewed L-Shaped Polyomino, Signed Tilings, Gröbner Basis, Coloring Invariants

1. Introduction

In this article, we study tiling problems for regions in a square lattice by certain symmetries of an L-shaped polyomino. Polyominoes were introduced by Golomb in [1] and the standard reference about this subject is the book *Polyominoes* [2]. The *L*-shaped polyomino we study is placed in a square lattice and is made out of $n, n \ge 3$, unit squares, or *cells*. See Figure 1(a). In a $a \times b$ rectangle, a is the height and b is the base. We consider translations (only!) of the tiles shown in Figure 1(b). They are ribbon L-shaped n-ominoes. A ribbon polyomino [3] is a simply connected polyomino with no two unit squares lying along a line parallel to the first bisector y = x. We denote the set of tiles by T_{u} .

Tilings by T_n , *n* even are studied in [4] and [5], with [4] covering the case n = 4. We recall that a replicating tile is one that can make larger copies of itself. The order of replication is the number of initial tiles that fit in the larger copy. Replicating tiles were introduced by Golomb in [6]. In [7], we study replication of higher orders for several



(b) The set of tiles \mathcal{T}_n .

Figure 1. An *L n*-omino and the tile set T_n .

replicating tiles introduced in [6]. In particular, it is suggested there that the skewed *L*-tetromino shown in **Figure 2(a)** is not replicating of order k^2 for any odd *k*. The question is equivalent to that of tiling a *k*-inflated copy of the straight *L*-tetromino using only four, out of eight possible, orientations of an *L*-tetromino, namely those that are ribbon. The question is solved in [4], where it is shown that the *L*-tetromino is not replicating of any odd order. This is a consequence of a stronger result: a tiling of the first quadrant by T_4 always follows the rectangular pattern, that is, the tiling reduces to a tiling by 4×2 and 2×4 rectangles, each tiled in turn by two tiles from T_4 .

The results in [4] are generalized in [5] to T_n , n even. The main result shows that any tiling of the first quadrant by T_n reduces to a tiling by $n \times 2$ and $2 \times n$ rectangles, with each rectangle covered by two ribbon *L*-shaped *n*-ominoes. An application is the characterization of all rectangles that can be tiled by T_n , n even: a rectangle can be tiled if and only if both sides are even and at least one side is divisible by n. Another application is the existence of the local move property for an infinite family of sets of tiles: T_n , n even, has the local move property for the class of rectangular regions with respect to the local moves that interchange a tiling of an $n \times n$ square by n/2 vertical rectangles, with a tiling by n/2 horizontal rectangles, each vertical/horizontal rectangle being covered by two ribbon *L*-shaped *n*-ominoes. One shows that neither of these results is valid for any odd n. The rectangular pattern of a tiling of the first quadrant persists if one adds an extra 2×2 tile to T_n , n even. A rectangle can be tiled by the larger set of tiles if and only if it has both sides even. It is also shown in the paper that the main result implies that a skewed *L*-shaped *n*-omino, n even, (see Figure 2(b)) is not a replicating tile of order k^2 for any odd k.

We investigate in this paper tiling properties of T_n , n odd. Parallel results with [8] are not possible due to the fact, already observed in [5], that there are rectangles that have tilings by T_n , n odd, which do not follow the rectangular pattern. See **Figure 3**. Instead of regular tilings, one can study signed tilings. These are finite placements of tiles on a plane, with weights +1 or -1 assigned to each of the tiles. We say that they tile a region R if the sum of the weights of the tiles is 1 for every cell inside R and 0 for every cell elsewhere.

A useful tool in the study of signed tilings is a Gröbner basis associated to the polynomial ideal generated by the tiling set. If the coordinates of the lower left corner of a cell are (α, β) , one associates to the cell the monomial $x^{\alpha}y^{\beta}$. This correspondence



Figure 2. Skewed polyominoes.



Figure 3. A tiling of a (3n, 3n+1) rectangle by T_n .

associates to any bounded tile placed in the square lattice a Laurent polynomial with all coefficients 1. The polynomial associated to a tile *P* is denoted by f_P . The polynomial associated to a tile translated by an integer vector (γ, δ) is the initial polynomial multiplied by the monomial $x^{\gamma}y^{\delta}$. If the region we want to tile is bounded and if the tile set consists of bounded tiles, then the whole problem can be translated in the first quadrant via a translation by an integer vector, and one can work only with regular polynomials in $\mathbb{Z}[X,Y]$. See Theorem 13 below.

Our main result is the following:

Theorem 1. A rectangle can be signed tiled by T_n , $n \ge 5$ odd, if and only if it has a side divisible by n.

Theorem 1 is proved in Section 4 using a Gröbner basis for the tiling set computed in Section 3.

For completeness, we briefly discuss regular tilings by $T_n, n \ge 5$ odd.

Theorem 1 gives for regular tilings by $T_n, n \ge 5$ odd, a corollary already known (see Lemma 2 in [5]):

Theorem 2. If $n \ge 5$ odd, a rectangle with neither side divisible by n cannot be tiled by T_n .

If one of the sides of the rectangle is divisible by *n*, we recall first the following result of Herman Chau, mentioned in [5], which is based on a deep result of Pak [3].

Theorem 3. A rectangle with both sides odd cannot be tiled by T_n , $n \ge 5$ odd. If one of the sides of the rectangle is even, one has the following result.

Theorem 4. Let $n \ge 5$ odd and assume that a rectangle has a side divisible by n and a side of even length.

1) If one side is divisible by n and the other side is of even length, then the rectangle can be tiled by T_n .

2) If the side divisible by n is of length at least 3n + 1 and even, and the other side is of length at least 3n and odd, then the rectangle can be tiled by T_n .

Proof. 1) The rectangle can be tiled by $2 \times n$ or $n \times 2$ rectangles, which can be tiled by two tiles from T_n .

2) We use the tiling shown in **Figure 4**. The (3n, 3n+1) rectangle is tiled as in **Figure 3**, and the other two rectangles can be tiled by $2 \times n$ or $n \times 2$ rectangles, which in turn can be tiled by two tiles from T_n .

A consequence of the technique used in the proof of Theorem 1 is:

Proposition 5. If $n \ge 5$ odd and $k \ge 1$, then a k-inflated copy of the L n-omino has a signed tiling by ribbon L n-ominoes.

Proposition 5 is proved in Section 5.

As any $2n \times 2n$ square can be tiled by T_n , it follows that if k is divisible by 2n then the skewed *Ln*-omnino is replicating of order k^2 . Information about other orders of replication can be found by using Pak's invariant [3].

Proposition 6. Let $n \ge 5$ odd.

1) If $k \ge 1$ is odd and divisible by n, then the skewed L n-omino is not replicating of order k^2 .

2) If $k \ge 1$ is even and not divisible by n, then the skewed L n-omino is not replicating of order k^2 .

Proposition 6 is proved in Section 6. Proposition 6 leaves open the question of replication of the skewed *L n*-omino if *k* is odd and not divisible by *n*. Some cases can be solved by using Pak's higher invariants f_2, \dots, f_m [3], which are all zero for tiles in T_n . For example, if n = 5, a 3-inflated copy of the *L* pentomino has $f_2 = -1$, showing the impossibility of tiling. A general result for regular tilings is out of reach due to the fact that for *k* odd and congruent to 1 modulo *n*, the leftover region that appears (see the proof of Proposition 6) is just an *L n*-omino that has all higher invariants f_2, \dots, f_m equal to zero. This is in contrast to the case of regular tilings by T_n, n even, discussed in [5], which is very well understood.



Figure 4. A tiling of an (odd, even) rectangle by T_n .



For completeness, we also consider the tile set T_n^+ , consisting of T_n , *n* odd, and an extra 2×2 square. For *n* even, the tile set T_n^+ is studied in [5]. It is shown there that there is a similarity between the regular tiling properties of T_n^+ and T_n .

Theorem 7. If $n \ge 5$ odd, any region in a square lattice can be signed tiled by T_n^+ **Theorem 7** is proved in Section 7.

Barnes developed in [9] [10] a general method for solving signed tiling problems with complex weights. In Section 8, we review the method of Barnes and offer an alternative proof of Theorem 1 based on this method. Having available a Gröbner basis for the tiling set helps even if Barnes method is used.

Theorem 8. If complex or rational weights are allowed to replace the integral weights, a rectangle can be signed tiled by $T_n, n \ge 5$ odd, if and only if it has a side divisible by n.

Signed tilings by T_n , *n* even, are more complicated than in the odd case. They are discussed in [11].

We make a final comment about the paper. While the methods that we use are well known, and algorithmic when applied to a particular tiling problem, here we apply them to solve simultaneously an infinite collection of tiling problems.

2. Summary of Gröbner Basis Theory

An introduction to signed tilings can be found in the paper of Conway and Lagarias [12]. One investigates there signed tilings by the 3-bone, a tile consisting of three adjacent regular hexagons. The Gröbner basis approach to signed polyomino tilings was proposed by Bodini and Nouvel [13]. In [8] one uses this approach to study signed tilings by the n-bone, a tile consisting of n collinear adjacent regular hexagons.

Let $\overline{R}[X] = R[X_1, \dots, X_k]$ be the ring of polynomials with coefficients in a principal ideal domain (PID) *R*. The only (PID) of interest in this paper is \mathbb{Z} , the ring of integers. A *term* in the variables x_1, \dots, x_k is a power product $x_1^{\alpha_1} x_2^{\alpha_2} \cdots x_{\ell}^{\alpha_{\ell}}$ with $\alpha_i \in \mathbb{N}, 1 \le i \le \ell$; in particular $1 = x_1^0 \cdots x_{\ell}^0$ is a term. A term with an associated coefficient from *R* is called *monomial*. We endow the set of terms with the *total degree-lexicographical order*, in which we first compare the degrees of the monomials and then break the ties by means of lexicographic order for the order $x_1 > x_2 > \cdots > x_{\ell}$ on the variables. If the variables are only x, y and x > y, this gives the total order:

$$1 < y < x < y^{2} < xy < x^{2} < y^{3} < xy^{2} < x^{2}y < x^{3} < y^{4} < \cdots.$$
(1)

For $P \in \overline{R}[X]$ we denote by HT(P) the leading term in P with respect to the above order and by HM(P) the monomial of HT(P). We denote by HC(P) the coefficient of the leading monomial in P. We denote by T(P) the set of terms appearing in P, which we assume to be in simplest form. We denote by M(P) the set of monomials in P. For a given ideal $I \subset \overline{R}[X]$ an associated Gröbner basis may be introduced for example as in Chapters 5, 10 [14]. Our summary follows the approach there. If $G \subset \overline{R}[X]$ is a finite set, we denote by I(G) the ideal generated by G in $\overline{R}[X]$.

Definition 1. Let $f, g, p \in \overline{R}[X]$. We say that fD-reduces to g modulo p and write $f \rightarrow_p g$ if there exists $m \in M(f)$ with HM(p)|m, say m = m' HM(p), and g = f - m'p. For a finite set $G \subset \overline{R}[X]$, we denote by \rightarrow_G the reflexive-transitive closure of \rightarrow_p , $p \in G$. We say that g is a normal form for f with respect to G if $f \rightarrow_G g$ and no further D-reduction is possible. We say that f is D-reducible modulo G if $f \rightarrow_G 0$.

It is clear that if $f \rightarrow_G 0$, then f belongs to the ideal generated by G in $\overline{R}[X]$. The converse is also true if G is a Gröbner basis.

Definition 2. A *D*-Gröbner basis is a finite set *G* in $\overline{R}[X]$ with the property that all *D*-normal forms modulo *G* of elements of I(G) equal zero. If $I \subset \overline{R}[X]$ is an ideal, then a *D*-Gröbner basis of *I* is a *D*-Gröbner basis that generates the ideal *I*.

Proposition 9. Let G be a finite set of $\overline{R}[X]$. Then the following statements are equivalent:

1) G is a Gröbner basis.

2) Every $f \neq 0, f \in I(G)$, is *D*-reducible modulo *G*.

Note that if R is only a (PID), the normal form of the division of f by G is not unique. We introduce now the notions of S-polynomial and G-polynomial.

Definition 3. Let $0 \neq g_i \in \overline{R}[X]$, i = 1, 2, with $HC(g_i) = a_i$, $HT(g_i) = t_i$. Let $a = b_i a_i = \operatorname{lcm}(a_1, a_2)$ with $b_i \in R$, and $t = s_i t_i = \operatorname{lcm}(t_1, t_2)$ with $s_i \in T$. Let $c_1, c_2 \in R$ such that $\operatorname{gcd}(a_1, a_2) = c_1 a_1 + c_2 a_2$. Then:

$$S(g_1, g_2) = b_1 s_1 g_1 - b_2 s_2 g_2,$$

$$G(g_1, g_2) = c_1 s_1 g_1 + c_2 s_2 g_2.$$
(2)

Remark. If $HC(g_1) = HC(g_2)$, then $G(g_1, g_2)$ can be chosen to be g_1 .

Theorem 10. Let G be a finite set of $\overline{R}[X]$. Assume that for all $g_1, g_2 \in G$, $S(g_1, g_2) \rightarrow_G 0$ and $G(g_1, g_2)$ is top-D-reducible modulo G. Then G is a Gröbner basis.

Assume now that *R* is a Euclidean domain with unique remainders (see page 463 [14]). This is the case for the ring of integers \mathbb{Z} if we specify remainders upon division by $0 \neq m$ to be in the interval [0, m).

Definition 4. Let $f, g, p \in \overline{R}[X]$. We say that *f E*-reduces to *g* modulo *p* and write $f \rightarrow_{p,E} g$ if there exists $m = at \in M(f)$ with HM(p)|t, say t = sHT(p), and g = f - qsp where $0 \neq q \in R$ is the quotient of *a* upon division with unique remainder by HC(p).

Proposition 11. *E-reduction extends D-reduction, i.e., every D-reduction step in an E-reduction step.*

Theorem 12. Let *R* be an Euclidean domain with unique remainders, and assume *G* is a finite set of $\overline{R}[X]$ and a *D*-Gröbner basis. Then the following hold:

1) $f \to_{G,E} 0$ for all $f \in I(G)$, where $\to_{G,E}$ denotes the *E*-reduction modulo *G*.

2) E-reduction modulo G has unique normal forms.

The following result connects signed tilings and Gröbner bases. See [13] and [8] for a proof.



Theorem 13. A polyomino P admits a signed tiling by translates of prototiles P_1, P_2, \dots, P_k if and only if for some (test) monomial $x^{\alpha}y^{\beta}$ the polynomial $x^{\alpha}y^{\beta}f_p$ is in the ideal generated in $\mathbb{Z}[X,Y]$ by the polynomials f_{P_1}, \dots, f_{P_k} . Moreover, the set of test monomials $\{x^{\alpha}y^{\beta}\}$ can be indexed by any set of multi-indices which is cofinal in (\mathbb{N}^n, \leq) .

3. Gröbner Basis for T_n, n Odd

We write $n = 2k + 1, k \ge 2$. The polynomials (in condensed form) associated to the tiles in T_n are:

$$G_{1}(k) = \frac{y^{2k} - 1}{y - 1} + x,$$

$$G_{2}(k) = y^{2k - 1} + x \frac{y^{2k} - 1}{y - 1},$$

$$G_{3}(k) = \frac{x^{2k} - 1}{x - 1} + y,$$

$$G_{4}(k) = x^{2k - 1} + y \frac{x^{2k} - 1}{x - 1}.$$
(3)

We show in the rest of this section that a Gröbner basis for the ideal generated in $\mathbb{Z}[X,Y]$ by $G_1(k), G_2(k), G_3(k), G_4(k)$, is given by the polynomials (written in condensed from):

$$B_{1}(k) = \frac{y^{k+2}-1}{y-1} + x \frac{x^{k-1}-1}{x-1},$$

$$B_{2}(k) = \frac{y^{k+1}-1}{y-1} + x \frac{x^{k}-1}{x-1},$$

$$B_{3}(k) = xy-1.$$
(4)

It is convenient to look at the elements of the basis geometrically, as signed tiles, see **Figure 5**. The presence of $B_3(k)$ in the basis allows reducing the algebraic proofs to combinatorial considerations. Indeed, using addition by a multiple of $B_3(k)$, one can translate, along a vector parallel to the first bisector y = x, cells labeled by +1 from one



Figure 5. The Gröbner basis $\{B_1(k), B_2(k), B_3(k)\}$.

position in the square lattice to another. See **Figure 6**. We will use this property repeatedly to check certain algebraic identities.

Proposition 14. $G_1(k), G_2(k), G_3(k), G_4(k)$ are in the ideal generated by $B_1(k), B_2(k), B_3(k)$.

Proof. The geometric proofs appear in **Figure 7**. First we translate one of the tiles $B_i(k), i = 1, 2$, multiplying by a power of x or a power of y, and then rearrange the cells from $B_i(k)$ using diagonal translations given by multiples of $B_3(k)$. The initial tiles $B_i(k), i = 1, 2$, have the cells marked by a +, and the final tiles $G_i(k), i = 1, 2, 3, 4$, are colored in light gray.

Proposition 15. $B_1(k), B_2(k), B_3(k)$ are in the ideal generated by $G_1(k), G_2(k), G_3(k), G_4(k)$.

Proof. We first show that $B_3(k)$ belongs to the ideal generated by $G_1(k), G_2(k), G_3(k), G_4(k)$. One has:

$$B_{3}(k) = -G_{1}(k) + G_{2}(k) + \left(-xy\frac{y^{2(k-1)} - 1}{y - 1} + y\frac{y^{2(k-1)} - 1}{y^{2} - 1}\right)G_{3}(k) + xy\frac{y^{2(k-1)} - 1}{y^{2} - 1}G_{4}(k).$$
(5)



Figure 6. Tiles arithmetic.



Figure 7. Generating $\{G_1(k), G_2(k), G_3(k), G_4(k)\}$ from $\{B_1(k), B_2(k), B_3(k)\}$.

Using (3), the RHS of Equation (5) becomes:

$$\begin{aligned} &\frac{1}{(y^2-1)(x-1)} \Big[-(y^{2k}-1)(y+1)(x-1) - x(y^2-1)(x-1) + y^{2k-1}(y^2-1)(x-1) \\ &+ x(y^{2k}-1)(y+1)(x-1) + \Big[-xy(y+1)(y^{2(k-1)}-1) + y(y^{2(k-1)}-1) \Big] \\ &\times \Big[y(x-1) + x^{2k}-1 \Big] + xy(y^{2(k-1)}-1) \Big[y(x^{2k}-1) + x^{2k-1}(x-1) \Big] \Big] \\ &= \frac{1}{(y^2-1)(x-1)} \Big[\Big(-y^{2k}+1 \Big) (xy-y+x-1) - x(y^2x-y^2-x+1) + y^{2k-1}(y^2x-y^2-x+1) \\ &+ (xy^{2k}-x)(xy-y+x-1) + (-xy^{2k}+xy^2-xy^{2k-1}+xy+y^{2k-1}-y) \\ &\times (xy-y+x^{2k}-1) + (xy^{2k-1}-xy)(yx^{2k}-y+x^{2k}-x^{2k-1}) \Big] \\ &= \frac{1}{(y^2-1)(x-1)} \Big[-y^{2k+1}x + y^{2k-1} - y^{2k}x + y^{2k} + xy-y+x-1 - x^2y^2 + xy^2 + x^2 - x \\ &+ xy^{2k+1} - y^{2k+1} - xy^{2k-1} + y^{2k-1} + x^2y^{2k+1} - xy^{2k+1}y^2 - xy^{2k} - x^2y + xy - x^2 \\ &+ xy^{2k+1} - y^{2k+1} - xy^{2k-1} + y^{2k-1} + x^2y^3 - xy^3 + x^{2k+1}y^2 - xy^2 - x^2y^{2k} + xy^{2k} \\ &- x^{2k+1}y^{2k-1} + xy^{2k-1} + x^2y^2 - xy^2 + x^{2k+1}y - xy + xy^{2k} - y^{2k} + x^{2k}y^{2k-1} - y^{2k-1} - xy^2 \\ &+ y^2 - x^{2k}y + y + x^{2k+1}y^{2k} - x^{2k+1}y^2 - xy^{2k} + xy^2 + x^{2k-1}y^{2k-1} - x^{2k}y^{2k-1} - x^{2k}y^{2k-1} + x^{2k}y^2 \\ &= \frac{1}{(y^2-1)(x-1)} \Big[xy-1-x^2y+x^2y^3 - xy^3 - xy^2 + y^2 + x \Big] = xy-1 = B_3(k). \end{aligned}$$

After we obtain $B_3(k)$, polynomials $B_1(k)$, $B_2(k)$ can be obtained geometrically by reversing the processes in **Figure 7**. Reversing the process in **Figure 7(a)**, we first obtain a copy of $y^{k-2}B_1(k)$. This copy can be translated to the right using multiplication by x^{k-2} , and then can be pulled back with the corner in the origin using a translation by a vector parallel to y = x. Reversing the process in **Figure 7(c)**, we first obtain a copy of $x^{k-1}B_2(k)$. This copy can be translated up using multiplication by y^{k-1} , and then can be pulled back with the corner in the origin using a translation by a vector parallel to y = x.

A step by step geometric proof of formula (5) for n = 7 is shown in Figure 8. All cells in the square lattice without any label have weight zero. The proof can be easily generalized for any odd n.

Proposition 16. $B_1(k), B_2(k), B_3(k)$ and $G_1(k), G_2(k), G_3(k), G_4(k)$ generate the same ideal in $\mathbb{Z}[X,Y]$.

Proof. This follows from Propositions 14, 15.

Proposition 17. One has the following D-reductions

$$S(B_{1}(k), B_{2}(k)) = -y^{k}B_{1}(k) + x^{k-1}B_{2}(k) + \left(x^{k-1}\frac{y^{k}-1}{y-1} - y^{k}\frac{x^{k-1}-1}{x-1}\right)B_{3}(k)$$
$$S(B_{1}(k), B_{3}(k)) = B_{2}(k) + \frac{y^{k}-1}{y-1}B_{3}(k)$$
$$S(B_{2}(k), B_{3}(k)) = B_{1}(k) + \frac{x^{k-1}-1}{x-1}B_{3}(k).$$

Consequently, $B_1(k), B_2(k), B_3(k)$ is a Gröbner basis.



Figure 8. The polynomial $B_3(7)$ is generated by $\{G_1(k), G_2(k), G_3(k), G_4(k)\}$.

Proof. The leading monomial of $B_1(k)$ is y^{k+1} , the leading monomial of $B_2(k)$ is x^k and the leading monomial of $B_3(k)$ is xy. We reduce the S-polynomials related to the set $B_1(k), B_2(k), B_3(k)$:

$$\begin{split} S\left(B_{1}(k),B_{2}(k)\right) &= x^{k}B_{1}(k) - y^{k+1}B_{2}(k) \\ &= x^{k}\left(\frac{y^{k+2}-1}{y-1} + x\frac{x^{k-1}-1}{x-1}\right) - y^{k+1}\left(\frac{y^{k+1}-1}{y-1} + x\frac{x^{k}-1}{x-1}\right) \\ &= \frac{-xy^{2k+2}-x^{k}y^{k+2} + x^{k} + y^{2k+2} - y^{k+1} + x^{2k}y - x^{k+1}y + xy^{k+2} - x^{2k} + y^{k+1}x^{k+1}}{(x-1)(y-1)} \\ &= -y^{k}B_{1}(k) + x^{k-1}B_{2}(k) + \left(x^{k-1}\frac{y^{k}-1}{y-1} - y^{k}\frac{x^{k-1}-1}{x-1}\right)B_{3}(k). \\ S\left(B_{1}(k),B_{3}(k)\right) &= xB_{1}(k) - y^{k}B_{3}(k) \\ &= x\left(\frac{y^{k+2}-1}{y-1} + x\frac{x^{k-1}-1}{x-1}\right) - y^{k}(xy-1) \\ &= \frac{x + x^{k+1}y - x^{2}y - x^{k+1} + x^{2}y^{k+1} - xy^{k+1} + xy^{k+1} - xy^{k} - y^{k+1} + y^{k}}{(x-1)(y-1)} \\ &= B_{2}(k) + \frac{y^{k}-1}{y-1}B_{3}(k). \end{split}$$



$$S(B_{2}(k), B_{3}(k)) = yB_{2}(k) - x^{k-1}B_{3}(k) = y\left(\frac{y^{k+1}-1}{y-1} + x\frac{x^{k}-1}{x-1}\right) - x^{k-1}(xy-1)$$
$$= \frac{xy^{k+2} - y^{k+2} - x^{k} + x^{k}y^{2} - yx^{k-1} - xy^{2} + y + x^{k-1}}{(x-1)(y-1)}$$
$$= B_{1}(k) + \frac{x^{k-1}-1}{x-1}B_{2}(k).$$

We show now that all above reductions are *D*-reductions by looking at the elimination of the terms of highest degree in the *S*-polynomials.

The terms of highest degrees in $S(B_1(k), B_2(k))$, after the initial reduction

$$\begin{aligned} x^{k}B_{1}(k) - y^{k+1}B_{2}(k) &= x^{k}\left(y^{k+1} + y^{k} + y^{k-1} + \dots + x^{k-1} + x^{k-2} + x^{k-3} + \dots\right) \\ &- y^{k+1}\left(y^{k} + y^{k-1} + y^{k-2} + \dots + x^{k} + x^{k-1} + x^{k-2} + \dots\right), \end{aligned}$$

are (in this order)

$$-y^{2k+1} + x^k y^k - x^{k-1} y^{k+1} - y^{2k}.$$

The terms $-y^{2k+1} - y^{2k}$ are contained in

$$-y^{k}B_{1}(k) = -y^{k}(y^{k+1} + y^{k} + y^{k-1} + \dots + x^{k-1} + x^{k-2} + x^{k-3} + \dots),$$

which does not contains terms of higher degree then $x^k y^k - x^{k-1} y^{k+1}$. The remaining terms $x^k y^k - x^{k-1} y^{k+1}$ are contained in

$$\begin{pmatrix} x^{k-1} \frac{y^k - 1}{y - 1} - y^k \frac{x^{k-1} - 1}{x - 1} \end{pmatrix} B_3(k) = \begin{bmatrix} x^{k-1} (y^{k-1} + y^{k-2} + y^{k-3} + \cdots) - y^k (x^{k-2} + x^{k-3} + \cdots) \end{bmatrix} (xy - 1),$$

which also does not contain terms of higher degree then $x^k y^k - x^{k-1} y^{k+1}$.

The term of highest degrees in $S(B_1(k), B_3(k))$, after the initial reduction

$$xB_{1}(k) - y^{k}B_{3}(k) = x(y^{k+1} + y^{k} + y^{k-1} + \dots + x^{k-1} + x^{k-2} + x^{k-3} + \dots) - y^{k}(xy - 1)$$

is xy. This term is contained in

)

$$\frac{y^{k}-1}{y-1}B_{3}(k) = (y^{k-1}+y^{k-2}+\cdots)(xy-1),$$

which does not contain terms of higher degree then xy^k

The term of highest degrees in $S(B_2(k), B_3(k))$, after the initial reduction

$$yB_{2}(k) - x^{k-1}B_{3}(k) = y(y^{k} + y^{k-1} + y^{k-2} + \dots + x^{k} + x^{k-1} + x^{k-2} + \dots) - x^{k-1}(xy-1)$$

is y^{k+1} . This term is contained in B_1 , which does not contain terms of higher degree then y^{k+1} .

As all higher coefficients are equal to 1, we do not need to consider the G-polynomials.

4. Proof of Theorem 1

Consider a $q \times p, q \ge p \ge 1$, rectangle. Using the presence of $B_3(k)$ in the Gröbner basis, and Theorem 13, the existence of a signed tiling becomes equivalent to deciding

when the polynomial:

$$P_{p,q}(x) = 1 + 2x + 3x^{2} + \dots + px^{p-1} + px^{p} + \dots + px^{q-1} + (p-1)x^{q} + (p-2)x^{q+1} + \dots + 2x^{p+q-3} + x^{p+q-2}$$

is divisible by the polynomial:

$$Q(x) = 1 + x + x^{2} + \dots + x^{n-1}.$$

If p+q-1 < n, then $\deg Q > \deg P_{p,q}$, so divisibility does not hold. If $p+q-1 \ge n$ we look at $P_{p,q}(x)$ as a sum of p polynomials with all coefficients equal to 1:

$$\begin{split} P_{p,q}\left(x\right) &= 1 + x + x^{2} + x^{3} + \dots + x^{p-1} + x^{p} + \dots + x^{q-1} + x^{q} + x^{q+1} + \dots + x^{p+q-4} + x^{p+q-3} + x^{p+q-2} \\ &+ x + x^{2} + x^{3} + \dots + x^{p-1} + x^{p} + \dots + x^{q-1} + x^{q} + x^{q+1} + \dots + x^{p+q-4} + x^{p+q-3} \\ &+ x^{3} + \dots + x^{p-1} + x^{p} + \dots + x^{q-1} + x^{q} + x^{q+1} + \dots + x^{p+q-4} + \dots + x^{p} + \dots + x^{q-1}. \end{split}$$

Assume that $p+q-1=nm+r, 0 \le r < n$, and $p=ns+t, 0 \le t < n$. The remainder $R_{p,q}(x)$ of the division of $P_{p,q}(x)$ by Q(x) is the sum of the remainders of the division of the *p* polynomials above by Q(x).

If *r* is odd, one has the following sequence of remainders, each remainder written in a separate pair of parentheses:

$$\begin{aligned} R_{p,q}\left(x\right) &= \left(1 + x + x^{2} + \dots + x^{r-1}\right) + \left(x + x^{2} + \dots + x^{r-2}\right) + \left(x^{2} + \dots + x^{r-3}\right) + \dots \\ &+ \left(x^{\frac{r-1}{2}} + x^{\frac{r+1}{2}}\right) + \left(0\right) - \left(x^{\frac{r-1}{2}} + x^{\frac{r+1}{2}}\right) - \dots - \left(x + x^{2} + \dots + x^{r-2}\right) \\ &- \left(1 + x + x^{2} + \dots + x^{r-1}\right) + \left(x^{r+1} + x^{r+3} + \dots + x^{n-3} + x^{n-2}\right) \\ &+ \left(x^{r+2} + \dots + x^{n-3}\right) + \dots + \left(x^{\frac{r+n}{2}}\right) - \left(x^{\frac{r+n}{2}}\right) + \dots - \left(x^{r+2} + \dots + x^{n-3}\right) \\ &- \left(x^{r+1} + x^{r+3} + \dots + x^{n-3} + x^{n-2}\right) - \dots \end{aligned}$$

If $p \ge n$, the sequence of remainders above is periodic with period *n*, given by the part of the sequence shown above, and the sum of any subsequence of *n* consecutive remainders is 0. So if *p* is divisible by *n*, $P_{p,q}(x)$ is divisible by Q(x). If *p* is not divisible by *n*, then doing first the cancellation as above and then using the symmetry of the sequence of remainders about the remainder equal to 0, the sum of the sequence of remainders equals 0 only if r+1=t, that is, only if *q* is divisible by *n*.

If *r* is even, one has the following sequence of remainders, each remainder written in a separate pair of parentheses:

$$\begin{aligned} R_{p,q}\left(x\right) &= \left(1 + x + x^{2} + \dots + x^{r-1}\right) + \left(x + x^{2} + \dots + x^{r-2}\right) + \left(x^{2} + \dots + x^{r-3}\right) \\ &+ \dots + \left(x^{\frac{r}{2}}\right) - \left(x^{\frac{r}{2}}\right) - \dots - \left(x + x^{2} + \dots + x^{r-2}\right) \\ &- \left(1 + x + x^{2} + \dots + x^{r-1}\right) + \left(x^{r+1} + x^{r+2} + \dots + x^{n-3} + x^{n-2}\right) \\ &+ \left(x^{r+2} + \dots + x^{n-3}\right) + \dots + \left(x^{\frac{r+n-1}{2}} + x^{\frac{r+n+1}{2}}\right) + \left(0\right) - \left(x^{\frac{r+n-1}{2}} + x^{\frac{r+n+1}{2}}\right) \\ &- \dots - \left(x^{r+2} + \dots + x^{n-3}\right) - \left(x^{r+1} + x^{r+2} + \dots + x^{n-3} + x^{n-2}\right) - \dots \end{aligned}$$



If $p \ge n$, the sequence of remainders above is periodic with period *n*, given by the part of the sequence shown above, and the sum of any subsequence of *n* consecutive remainders is 0. So if *p* is divisible by *n*, $P_{p,q}(x)$ is divisible by Q(x). If *p* is not divisible by *n*, then doing first the cancellation as above and then using the symmetry of the sequence of remainders about the remainder equal to 0, the sum of the sequence of remainders equals 0 only if r+1=t, that is, only if *q* is divisible by *n*.

5. Proof of Proposition 5

Consider a k-inflated copy of the L n-omino. Using the presence of $B_3(k)$ in the Gröbner basis, and Theorem 13, the existence of a signed tiling of the copy becomes equivalent to deciding when a $k \times nk$ rectangle has a signed tiling by T_n . Theorem 1 implies that this is always the case.

6. Proof of Proposition 6

1) We employ a ribbon tiling invariant introduced by Pak [3]. Each ribbon tile of length *n* can be encoded uniquely as a binary string of length n-1, denoted $(\varepsilon_1, \dots, \varepsilon_{n-1})$, where a 1 represents a down movement and a 0 represents a right movement. The encoding of a $1 \times n$ bar is $(0, 0, \dots, 0)$, for a $n \times 1$ bar is $(1, 1, \dots, 1)$, and for the tiles in T_5 the encodings are shown in Figure 9. Pak showed that the function $f_1(\varepsilon_1, \dots, \varepsilon_{n-1}) = \varepsilon_1 - \varepsilon_{n-1}$ is an invariant of the set of ribbon tiles made of *n*-cells, which contains as a subset the tile set T_n . In particular, one has that

$$f_1(\varepsilon_1,\cdots,\varepsilon_{n-1})=\pm 1$$

for any tile in T_n . The area of a k-inflated copy of the L n-omino is an odd multiple of n and can be easily covered by $n \times 1$ and $1 \times n$ bars, each one having the invariant equal to zero. If we try to tile by T_n , then the invariant is zero only if we use an even number of tiles. But this is impossible because the area is odd.

2) Let $k = n\ell + r, 0 < r < n$. After cutting from a k-inflated copy a region that can be covered by $n \times 1$ and $1 \times n$ bars, and which has the f_1 invariant equal to zero, we are left with one of the regions shown in Figure 10. Case a) appears if 2r < n and case b) appears if 2r > n. Both of these regions can be tiled by r ribbon tiles of area n as in Figure 11. In the first case the sequence of r encodings of the ribbon tiles is:

 $\begin{array}{c} 1,1,1,\cdots,1,1,1,0,0,\cdots,0,0,0\\ 1,1,1,\cdots,1,1,0,0,0,\cdots,0,0,1\\ 1,1,1,\cdots,1,0,0,0,0,\cdots,0,1,1\\ \\ \end{array}$



Figure 9. The four L-shaped ribbon pentominoes and their encodings.



Figure 10. Leftover regions.



Figure 11. Tiling the leftover region by ribbon n tiles, cases n = 5; k = 4, and n = 17; k = 20.

where we start with n-r-1 ones and *r* zeros, and then shift the zeroes to the left by 1 at each step, completing the sequence at the end with ones. As $r \le n - r - 1$, the subsequence of zeroes does not reach the left side, so the f_1 invariant of the region is equal to 1.

In the second case, the sequence of r encodings of the ribbon tiles starts as above, but now the subsequence of zeroes reaches the left side. Then we have a jump of n-runits of the sequence of zeroes to the left, the appearance of an extra one at the right, and a completion of the sequence by zeroes to the right. Then the subsequence of ones that appears start shifting to the right till it reaches the right edge. The f_1 invariant of the region is equal to -1.

So in both cases the f_1 invariant is an odd number. Nevertheless, if the k-copy is tiled by T_n , one has to use an even number of tiles and the invariant is an even number. Therefore we have a contradiction.

7. Proof of Theorem 7

It is enough to generate the tile consisting of a single cell. We show the proof for n = 7in **Figure 12**. The proof can be easily generalized to any $n \ge 5$ odd. First we construct a domino with both cells having the same sign (as in Figure 12(c)), and then we use it to reduce the *L n*-omino until a single cell is left.

8. The Method of Barnes

In this section we give a proof of Theorem 8 following a method developed by Barnes. The reader of this section should be familiar with [9] [10]. We apply the method to the



Figure 12. Generating a single cell by T_n^+ .

infinite collection of tiling sets $T_n, n \ge 5$ odd.

Let $n \ge 5$ odd. Consider the polynomials (3) associated to the tiles in T_n and denote by *I* the ideal generated by $G_1(k), G_2(k), G_3(k), G_4(k)$. We show that the algebraic variety *V* defined by *I* is zero dimensional and consists only of the pairs of points

$$\left(\varepsilon, \frac{1-\varepsilon^n}{1-\varepsilon}\right),\tag{6}$$

where ε is an *n*-th root of identity different from 1.

Separate x from $G_1(k) = 0$ and replace in $G_2(k) = 0$ to have:

$$y^{2k-1} - \frac{y^{2k} - 1}{y - 1} \frac{y^{2k} - 1}{y - 1} = 0.$$

Eliminating the denominators gives:

$$y^{2k-1} - (y^{2k-1} + y^{2k-2} + \dots + y^2 + y + 1)^2 = 0,$$

which can be factored as:

 $(y^{2k} + y^{2k-1} + y^{2k-2} + \dots + y^2 + y + 1)(y^{2k-2} + y^{2k-3} + \dots + y^2 + y + 1) = 0.$

It is clear that all roots of the polynomial above, and of the corresponding polynomial in the variable x, are roots of unity of order 2k+1 and 2k-1. Using the system of equations that defines V, the roots of order 2k-1 can be eliminated. Moreover, the only solutions of the system are given by (6).

We show now that *I* is a radical ideal. For this we use an algorithm of Seidenberg which can be applied to find the radical ideal of a zero dimensional algebraic variety over an algebraically closed field. See Lemma 92 in [15]. Compare also with Theorem 7.1 in [9]. As *V* is zero dimensional, one can find $f_1(x)$ and $f_2(y)$ that belong to the radical ideal. We consider the square free polynomials:

$$f_1(x) = x^{2k} + x^{2k-1} + x^{2k-2} + \dots + x^2 + x + 1,$$

$$f_2(y) = y^{2k} + y^{2k-1} + y^{2k-2} + \dots + y^2 + y + 1.$$

If $f_1(x), f_2(y)$ are square free, then the ideal generated by *I* and $f_1(x), f_2(y)$ is radical. So, in order to show that *I* itself is radical, it is enough to show that $f_1(x), f_2(y)$ belong to *I*. It is easier to generate $f_i(x)$ using the Gröbner basis, so we will use this approach.

Proposition 18. The polynomials $f_1(x), f_2(y)$ belong to the ideal *I*. *Proof.* It is enough to generate $f_1(x)$. One has:

$$f_1(x) = xG_3(x) - B_3(k)$$

We can apply now the main result in Lemma 3.8, [9]: a region R is signed tiled by T_n if and only if the polynomial $f_R(x, y)$ associated to R evaluates to zero in any point of the variety V. If R is a rectangle of dimensions $p \times q$ in the square lattice, then

$$f_{R}(x, y) = \frac{x^{q} - 1}{x - 1} \frac{y^{p} - 1}{y - 1},$$

which clearly evaluates to zero in all points of V if and only if one of p,q is divisible by *n*.

The fact that Theorem 8 implies Theorem 1 follows the idea of Theorem 4.2 in [9]. Indeed, a set of generators for the regions that are signed tiled with rational numbers by T_n is given by the polynomials $f_1(x), f_2(y)$ above. Both of them can be generated by the Gröbner basis using only integer coefficients.

9. Conclusion

We show that a rectangle can be signed tiled by ribbon L n-ominoes, n odd, if and only if it has a side divisible by n. A consequence of our technique, based on the exhibition of an explicit Gröbner basis, is that any k-inflated copy of the skewed L n-omino has a signed tiling by skewed L n-ominoes.

Acknowledgements

V. Nitica was partially supported by Simons Foundation Grant 208729.

References

- [1] Golomb, S.W. (1954) Checker Boards and Polyominoes. American Mathematical Monthly, 61, 675-682. http://dx.doi.org/10.2307/2307321
- [2] Golomb, S.W. (1994) Polyominoes, Puzzles, Patterns, Problems, and Packings. Princeton University Press, Princeton.
- [3] Pak, I. (2000) Ribbon Tile Invariants. Transactions of the American Mathematical Society, 352, 5525-5561. http://dx.doi.org/10.1090/S0002-9947-00-02666-0
- [4] Chao, M., Levenstein, D., Nitica, V. and Sharp, R. (2013) A Coloring Invariant for Ribbon L-Tetrominoes. Discrete Mathematics, 313, 611-621. http://dx.doi.org/10.1016/j.disc.2012.12.007
- [5] Nitica, V. (2015) Every Tiling of the First Quadrant by Ribbon L n-Ominoes Follows the Rectangular Pattern. Open Journal of Discrete Mathematics, 5, 11-25. http://dx.doi.org/10.4236/ojdm.2015.52002
- [6] Golomb, S.W. (1964) Replicating Figures in the Plane. *Mathematical Gazette*, 48, 403-412. http://dx.doi.org/10.2307/3611700
- [7] Nitica, V. (2003) Rep-Tiles Revisited. In: Katok, S., Sossinsky, A. and Tabachnikov, S., Eds., MASS Selecta: Teaching and Learning Advanced Undergraduate Mathematics, American Mathematical Society, Providence, 205-217.

- [8] Dizdarevic, M.M., Timotijevic, M. and Zivaljevic, R.T. (2016) Signed Polyomino Tilings by n-in-Line Polyominoes and Gröbner Bases. *Publications de l'Institut Mathematique, Nou*velle série, 99, 31-42.
- Barnes, F.W. (1982) Algebraic Theory of Brick Packing I. *Discrete Mathematics*, 42, 7-26. http://dx.doi.org/10.1016/0012-365X(82)90049-8
- [10] Barnes, F.W. (1982) Algebraic Theory of Brick Packing II. Discrete Mathematics, 42, 129-144. <u>http://dx.doi.org/10.1016/0012-365X(82)90211-4</u>
- [11] Gill, K. and Nitica, V. (2016) Signed Tilings by Ribbon L n-Ominoes, n Odd, via Gröbner bases. Open Journal of Discrete Mathematics, 7, 185-206. http://dx.doi.org/10.4236/ojdm.2016.63017
- [12] Conway, J.H. and Lagarias, J.C. (1990) Tilings with Polyominoes and Combinatorial Group Theory. *Journal Combinatorial Theory, Series A*, 53, 183-208. http://dx.doi.org/10.1016/0097-3165(90)90057-4
- [13] Bodini, O. and Nouvel, B. (2004) Z-Tilings of Polyominoes and Standard Basis. In Combinatorial Image Analysis, Springer, Berlin, 137-150.
- [14] Becker, T. and Weispfenning, V. (in cooperation with Heinz Krendel) (1993) Gröbner Bases. Springer-Verlag, Berlin.
- [15] Seidenberg, A. (1974) Constructions in Algebra. *Transactions of the American Mathemati*cal Society, 197, 273-313.

👯 Scientific Research Publishing

Submit or recommend next manuscript to SCIRP and we will provide best service for you:

Accepting pre-submission inquiries through Email, Facebook, LinkedIn, Twitter, etc. A wide selection of journals (inclusive of 9 subjects, more than 200 journals) Providing 24-hour high-quality service User-friendly online submission system Fair and swift peer-review system Efficient typesetting and proofreading procedure Display of the result of downloads and visits, as well as the number of cited articles Maximum dissemination of your research work Submit your manuscript at: <u>http://papersubmission.scirp.org/</u>

Or contact ojdm@scirp.org