

Temperature Dependence of the Raman Intensity and the Bandwidth Close to the Order-Disorder Phase Transition in NaNO_3

Hamit Yurtseven*, Sukran Aslan

Department of Physics, Middle East Technical University, Ankara, Turkey

Email: *hamit@metu.edu.tr

Received June 5, 2012; revised July 15, 2012; accepted July 20, 2012

ABSTRACT

We analyze the temperature dependence of the Raman intensity using the experimental data according to a power-law formula close to the order-disorder transition ($T_c = 549$ K) in NaNO_3 . From this analysis, we extract the value of $\beta = 0.26$ as the critical exponent for the order parameter, which indicates a tricritical phase transition in this crystalline system. Using the temperature dependence of the order parameter, the Raman bandwidth is calculated at various temperatures in NaNO_3 . Our calculated bandwidths describe adequately the observed behavior of the order-disorder transition in this crystal.

Keywords: Raman Intensity; Bandwidth; NaNO_3

1. Introduction

NaNO_3 exhibits an order-disorder phase transition at $T_c = 548$ K [1]. Below T_c , the NO_3^- ions are oriented along one preferred direction which gives rise to the long range order parameter. Above T_c , the NO_3^- ions are equally positioned between the two stable states.

The experimental studies such as the thermal expansion [1-3], X-ray [4] and light scattering [5-7] have been given in the literature. A large λ anomaly in the heat capacity of NaNO_3 has been observed at T_c [8,9]. Besides, the molecular NO desorption from crystalline sodium nitrate [10], by the laser desorption the electronic structure [11] and the electron beam induced damage of NaNO_3 [12] have been studied experimentally.

In order to explain the mechanism of the phase transitions in NaNO_3 , some models have been proposed for its disordered structure. There are mainly free rotational [1], two-position disordered model [13], as described in some detail previously [14]. The free rotational model and two-position disordered models have been identified essentially with the XY and Ising models, respectively [14]. Also, the disordering due to the reorientation of the NO_3^- ions has been modeled for the disordered aragonite structure [15], disordered calcite structure [4] and the mixture of those two models [16], as pointed out in a previous study [6]. Additionally, the orientational disordering in NaNO_3 and CaCO_3 has been studied by molecular dy-

namics simulation using semiempirical potential models [17,18]. The temperature dependence of the specific heat C_p and of the lattice parameters in calcite (CaCO_3) and in nitratine (NaNO_3) have also been simulated [19]. In these two structures, an orientational ordering of the carbonate and nitrate groups occurs as the temperature decreases below the transition temperature. At high temperatures, the nitrate and carbonate groups are orientationally disordered about their three fold axes [14]. This is the orientational long-range disorder above T_c , whereas there exists a positional long range order below T_c [6]. At low temperatures, sodium nitrate crystallizes in the ordered calcite structure which has the D_{3d}^6 space group ($R3c$) with two formula units of NaNO_3 per unit cell. Above T_c , the crystal changes the high-temperature phase that has one NaNO_3 molecule with the crystal symmetry $R3m$.

It has been observed from the Raman spectra that the change from complete order to complete disorder for NaNO_3 is spread out over a wide temperature range of about 100°C [6]. It has been pointed out that the Raman spectrum of NaNO_3 at high temperatures is a combination of the ordinary first-order Raman spectrum and the disorder-induced spectrum [6]. Above T_c , a central peak in the low-frequency Raman spectra of sodium nitrate has been observed [7]. It has been obtained that the Raman intensity and the bandwidth of this central peak due to a motion of NO_3^- ions, exhibit an anomalous behaviour at T_c with increasing temperature [7]. Thus, the light scattering spectra of the order-disorder motion of NO_3^- ions

*Corresponding author.

is accompanied with the phase transition of NaNO_3 . So, the anomalous part of the temperature dependence of the Raman intensity of the central peak has been associated with the order parameter [7] which is an average value of the difference in number of NO_3^- ions in the two stable positions [4]. As a result of this, the anomalous part of the Raman intensity is due to rearrangement of NO_3^- ions around the z axis [7].

In this study we analyze the temperature dependence of the observed Raman intensity of the central peak [7] below T_c in NaNO_3 . This analysis of the Raman intensity associated with the order parameter is performed according to a power-law formula with the critical exponent β . Using the Raman intensity as the order parameter, the damping constant of the central peak is calculated at various temperatures for NaNO_3 . The expressions derived for the damping constant from the models of the pseudospin-phonon interaction and of the energy fluctuation [20,21] are used. The damping constant is fitted to the experimentally measured Raman bandwidths of the central peak in NaNO_3 .

Below, we give our analysis of the spectral intensity and our calculation of the damping constant of the central peak of NaNO_3 in Section 2. Section 3 gives our discussion of the results. Finally, conclusions are given in Section 4.

2. Analysis of the Spectral Intensity and Calculation of the Bandwidth

We analyzed the central-peak intensity measured [7] as a function of temperature near T_c in NaNO_3 according to a power-law expression given by

$$I = A_0 (1 - T/T_c)^{2\beta} \quad (1)$$

where β is the critical exponent for the order parameter and A_0 is the amplitude. Since the intensity (order parameter) increases as the temperature decreases below T_c , the observed Raman intensity can be analyzed according to Equation (1). The anomalous part of the intensity in the $x(\text{zz})y$ geometry was analyzed using the experimental data [7]. The fitted parameters β and A_0 were deduced within the range of the reduced temperature from our analysis. This analysis was performed by taking the logarithm of both sides of Equation (1) which gives

$$\log I = \log A_0 + 2\beta \log \varepsilon \quad (2)$$

where the reduced temperature is $\varepsilon = (T_c - T)/T_c$ with the transition temperature $T_c = 549.1$ K for NaNO_3 .

Table 1 gives the fitted parameters of the critical exponent β for the order parameter and the amplitude A_0 within the interval of the reduced temperature ε for NaNO_3 . **Figure 1** gives our plot of the spectral intensity as a function of temperature in a log-log scale according to Equation (2) for NaNO_3 . **Figure 1** represents the increasing

Table 1. Values of the critical exponent β for the order parameter and the amplitude A_0 from the analysis of the spectral intensity according to Equation (1) for NaNO_3 .

T_c (K)	β	A_0	I_0	$\varepsilon = (T_c - T)/T_c$
549.1	0.26	4.05	7.53	$5.8 \times 10^{-4} < \varepsilon < 3.1 \times 10^{-2}$




Figure 1. The anomalous part of the observed Raman intensity [7] as a function of temperature in a log-log scale below T_c according to Equation (2) in NaNO_3 .

intensity as the temperature decreases, which corresponds to the difference between the extrapolated (above T_c) and observed intensity [7] below T_c in the temperature region from 530 to 548 K. In a log-log scale, $\ln I$ decreases against $-\ln[(T_c - T)/T_c]$, as shown in **Figure 1**.

We also calculated the temperature dependence of the damping constant close to the order-disorder phase transition ($T_c = 548$ K) in NaNO_3 . This calculation was performed using the expressions,

$$\Gamma_{sp} = \Gamma_0 + A(1 - P^2) \ln \left\{ T_c / [T - T_c(1 - P^2)] \right\} \quad (3)$$

$$\Gamma_{sp} = \Gamma'_0 + A' \left\{ T(1 - P^2) / [T - T_c(1 - P^2)] \right\}^{1/2} \quad (4)$$

where Γ and Γ'_0 represent background damping constants and, A and A' are constants. In Equations (3) and (4), P defines the order parameter and it is simply proportional to the spectral intensity given by

$$P^2 \propto I/I_0 \quad (5)$$

where I_0 is the maximum intensity of a spectral band. Equations (3) and (4) have been derived as approximate relations in earlier studies [20,21]. Using the Matsushita's expression [22] for the damping constant Laulicht and Lucnar [20,21] have been able to obtain the temperature dependence of the linewidth as approximate relations (Equations (3) and (4)) and they have applied for their analysis of the Raman linewidths for the ferroelectric material, KH_2PO_4 . In this study, we also used Equations (3) and (4) to calculate the temperature dependence of

the Raman linewidths of the central peak, as measured experimentally [7] for NaNO_3 . Thus, using the temperature dependence of the Raman intensity as P^2 in Equations (3) and (4) below T_c , we obtained the temperature dependence of the Raman bandwidth in NaNO_3 . By fitting Equations (3) and (4) to the observed bandwidth of the central peak in the $x(\text{zz})y$ geometry [7], we determined the background bandwidths Γ_0 and Γ'_0 , and also the amplitudes A and A' , as given in **Table 2**. Since there is no ordering ($P = 0$) above T_c , Equations (3) and (4) were fitted to the observed bandwidths of the central peak in the $x(\text{zz})y$ geometry [7] at various temperatures.

We plot in **Figure 2** our calculated Raman bandwidths of the central peak as a function of temperature below and above the transition temperature ($T_c = 548$ K) in NaNO_3 . The observed Raman bandwidths of the central peak [7] are also plotted in **Figure 2**.

3. Discussion

The temperature dependence of the Raman intensity was analyzed here according to a power-law formula (Equation 1) with the exponent value of $\beta = 0.26$ for the order parameter P , as given in **Figure 1**. This analysis was performed using the intensity data [7]. Considering the temperature dependence of the order parameter as $P \propto$

Table 2. Values of the background bandwidths Γ_0 and Γ'_0 , and the amplitudes A and A' according to Equations (3) and (4) for the central peak in the $x(\text{zz})y$ geometry [7] in NaNO_3 .

$T_c = 548$ K	Γ_0 (cm^{-1})	A (cm^{-1})	Γ'_0 (cm^{-1})	A' (cm^{-1})
$T < T_c$	135.03	43.63	112.81	39.07
$T > T_c$	111.58	4.20	122.27	0.76




Figure 2. Temperature dependence of the bandwidth calculated from Equations (3) and (4) above and below the transition temperature ($T_c = 548$ K) in NaNO_3 . The observed bandwidths of the central peak in the $x(\text{zz})y$ geometry [7] are also given here.

$(T_c - T)^\beta$ and the relationship between the intensity and the order parameter as $I \propto P^2$, Equation (1) was used to analyze the temperature dependence of the intensity which increases with decreasing temperature below T_c . The β values for the order parameter which we deduced from the Raman intensity indicate that the order-disorder transition in NaNO_3 is the tricritical transition. This is due to the fact that the observed Raman intensity [7] considered as an order parameter exhibits largely a continuous change with the temperature. It has been pointed out previously [14] that this is the orientational order-disorder transition which occurs through continuous planar rotations of the nitrate group rather than discrete jumps.

It has been obtained [7] the value of $\beta = 1/2$ as the critical index of the order parameter by analyzing the anomalous part of their measured Raman intensity in the temperature range between about 450 K and T_c . In this study, we reanalyzed their observed central-peak intensity in a relatively narrow temperature range of 530 to 548 K (T_c), as stated above (**Figure 1**), where the anomalous behavior of the central peak is more pronounced [7]. When the temperature range was narrowed, as performed in this study, the β value decreased from $1/2$ to 0.26, which indicates a cross-over from a second order ($\beta = 0.5$) to a tricritical ($\beta = 0.25$) transition within the mean field theory in NaNO_3 . On the other hand, the birefringence data gives $\beta = 0.22$ for $500 \text{ K} < T < T_c$ and the transition is tricritical ($\beta = 0.25$) for $T < 500$ K in NaNO_3 [23,24]. It has been explained that this apparent change in the value of β from 0.25 to 0.22 in NaNO_3 is due to strong critical fluctuations at temperatures close to T_c in NaNO_3 , where the mean field theory does not hold [14]. This is not in agreement with our analysis which gives $\beta = 0.26$ closer to the tricritical value of 0.25 as predicted from the mean field theory in the interval of the reduced temperature $6 \times 10^{-4} < \varepsilon < 3 \times 10^{-2}$ (**Table 1**). Our exponent value is also closer to the theoretical value of $\beta = 0.228$ (2) for the two-dimensional XY model.

The temperature dependence of the Raman bandwidth which increases (**Figure 2**), is in accordance with decreasing intensity of the central peak as the temperature decreases, as observed experimentally [7]. Our damping constant calculated according to Equations (3) and (4) describes the observed behavior of the temperature dependence of the Raman bandwidth for the central peak above and below T_c in NaNO_3 (**Figure 2**). Below about 500 K, the observed bandwidth seems to be independent of temperature and its value of about 200 cm^{-1} remains the same at low temperatures [7]. However, our predicted values of the damping constant Γ increase rapidly below 500 K according to Equations (3) and (4). This indicates that the damping constant (Raman bandwidth) can be predicted in accordance with the experimental results down to about 500 K below which Equations (3) and (4) are no longer

valid for the order-disorder phase transition in NaNO_3 . This is also in accordance with our analysis of the Raman intensity for the central peak with the critical exponent value of $\beta = 0.26$ in the temperature interval of $530 \text{ K} < T < 548 \text{ K}$ (Table 1).

Increasing the Raman intensity and narrowing the bandwidth of the central peak with increasing temperature, as observed experimentally [7] indicate that a “central mode” exists due to rearrangements of NO_3^- ions close to the order-disorder phase transition in NaNO_3 , as suggested previously [7]. This also appears in the disorder-induced Raman spectrum at high temperatures, which becomes effective in addition to the ordinary first-order Raman spectrum for NaNO_3 . This central mode which is related to the order-disorder phase transition in NaNO_3 close to T_c , becomes very active far above T_c where the NO_3^- ions are no longer restricted to any fixed orientation, as suggested previously [7]. This leads to the melting phenomena in NaNO_3 .

4. Conclusion

The temperature dependence of the observed Raman intensity (order parameter) was analyzed using a power-law formula. From our analysis of the Raman intensity for the central peak, it is indicated that the exponent value of $\beta = 0.26$ describes the observed behavior as the tricritical transition in NaNO_3 in a given temperature interval. Also, the temperature dependence of the damping constant calculated here is in agreement with the observed Raman bandwidth in NaNO_3 . This analysis given here can be applied to the central peaks close to order disorder phase transitions in some other crystals.

REFERENCES

- [1] F. C. Kracek, E. Posnjak and S. B. Hendricks, “Gradual Transition in Sodium Nitrate II. The Structure at Various Temperatures and Its Bearing on Molecular Rotation,” *Journal of American Chemical Society*, Vol. 53, No. 9, 1931, pp. 3339-3348. [doi:10.1021/ja01360a016](https://doi.org/10.1021/ja01360a016)
- [2] R. J. Reeder, S. A. T. Redfern and E. Salje, “Spontaneous Strain at the Structural Phase Transition in NaNO_3 ,” *Physics and Chemistry of Minerals*, Vol. 15, No. 6, 1988, pp. 605-611. [doi:10.1007/BF00311033](https://doi.org/10.1007/BF00311033)
- [3] Y. Takeuchi and Y. Sasaki, “Elastic Properties and Thermal Expansion of NaNO_3 Single Crystal,” *Journal of Physical Society of Japan*, Vol. 61, 1992, pp. 587-595. [doi:10.1143/JPSJ.61.587](https://doi.org/10.1143/JPSJ.61.587)
- [4] Y. Shinnaka, “Anion Rotational Disorder in Sodium Nitrate,” *Journal of Physical Society of Japan*, Vol. 19, 1964, pp. 1281-1290. [doi:10.1143/JPSJ.19.1281](https://doi.org/10.1143/JPSJ.19.1281)
- [5] I. Nakagawa and J. L. Walter, “Optically Active Crystal Vibrations of the Alkali & #8208; Metal Nitrates,” *Journal of Chemical Physics*, Vol. 51, 1969, pp. 1389-1397.
- [6] T. Y. Shen, S. S. Mitra, H. Prask and S. F. Trevino, “Order-Disorder Phenomenon in Sodium Nitrate Studied by Low-Frequency Raman Scattering,” *Physical Review*, Vol. 12, No. 10, 1975, pp. 4530-4533. [doi:10.1103/PhysRevB.12.4530](https://doi.org/10.1103/PhysRevB.12.4530)
- [7] H. Yasaka, A. Sakai and T. Yagi, “A Central Peak in the Order-Disorder Phase Transition of Sodium Nitrate,” *Journal of Physical Society of Japan*, Vol. 54, 1985, pp. 3697-3700. [doi:10.1143/JPSJ.54.3697](https://doi.org/10.1143/JPSJ.54.3697)
- [8] V. C. Reinsborough and F. E. W. Wetmore, “Specific Heat of Sodium Nitrate and Silver Nitrate by Medium High Temperature Adiabatic Calorimetry,” *Australian Journal of Chemistry*, Vol. 20, No. 1, 1967, pp. 1-8. [doi:10.1071/CH9670001](https://doi.org/10.1071/CH9670001)
- [9] T. Jirri, J. Rogez, C. Bergman and J. C. Mathieu, “Thermodynamic Study of the Condensed Phases of NaNO_3 , KNO_3 and CsNO_3 and Their Transitions,” *Thermochimica Acta*, Vol. 266, 1995, pp. 147-161. [doi:10.1016/0040-6031\(95\)02337-2](https://doi.org/10.1016/0040-6031(95)02337-2)
- [10] R. A. Bradley, E. Lanzendorf, M. I. McCarthy, M. Orlando and W. P. Hess, “Molecular NO Desorption from Crystalline Sodium Nitrate by Resonant Excitation of the $\text{NO}_3^- \pi - \pi^*$ Transition,” *Journal of Physical Chemistry*, Vol. 99, No. 30, 1995, pp. 11715-11721. [doi:10.1021/j100030a015](https://doi.org/10.1021/j100030a015)
- [11] M. I. McCarthy, K. A. Peterson and W. P. Hess, “Electronic Structure of Sodium Nitrate: Investigations of Laser Desorption Mechanisms,” *Journal of Physical Chemistry*, Vol. 100, No. 16, 1996, pp. 6708-6714. [doi:10.1021/jp953108n](https://doi.org/10.1021/jp953108n)
- [12] N. G. Petrik, K. Knutsen, E. Paparazzo, S. Lea, D. M. Camaion and T. M. Orlando, “Electron Beam Induced Damage of NaNO_3 Single Crystals: An Energy, Temperature, and Quantum State Resolved Study,” *Journal of Physical Chemistry*, Vol. 104, No. 7, 2000, pp. 1563-1571. [doi:10.1021/jp993310g](https://doi.org/10.1021/jp993310g)
- [13] J. A. A. Ketelaar and B. Strijk, “The Atomic Arrangement in Solid Sodium Nitrate at High Temperatures,” *Recueil des Travaux Chimiques des Pays-Bas et de la Belgique*, Vol. 64, 1945, p. 174.
- [14] M. J. Harris, “A New Explanation for the Unusual Critical Behavior of Calcite and Sodium Nitrate, NaNO_3 ,” *American Mineral*, Vol. 84, 1999, pp. 1632-1640.
- [15] S. Siegel, “Effect of Neutron Bombardment on Order in the Alloy Cu_3Au ,” *Physical Review*, Vol. 75, No. 12, 1949, pp. 1823-1824. [doi:10.1103/PhysRev.75.1823](https://doi.org/10.1103/PhysRev.75.1823)
- [16] K. O. Strømme, “The Crystal Structure of Sodium Nitrate in the High-Temperature Phase,” *Acta Chemica Scandinavica*, Vol. 23, 1969, pp. 1616-1624. [doi:10.3891/acta.chem.scand.23-1616](https://doi.org/10.3891/acta.chem.scand.23-1616)
- [17] R. M. Lynden-Bell, M. Ferrario, I. R. McDonald and E. Salje, “A Molecular Dynamics Study of Orientational Disorder in Crystalline Sodium Nitrate,” *Journal of Physics Condensed Matter*, Vol. 1, No. 37, 1989, pp. 6523-6542. [doi:10.1088/0953-8984/1/37/002](https://doi.org/10.1088/0953-8984/1/37/002)
- [18] M. Ferrario, R. M. Lynden-Bell and R. M. McDonald, “Structural Fluctuations and the Order-Disorder Phase Transition in Calcite,” *Journal of Physics Condensed Matter*, Vol. 6, No. 7, 1994, pp. 1345-1358.

- [doi:10.1088/0953-8984/6/7/007](https://doi.org/10.1088/0953-8984/6/7/007)
- [19] J. Liu, C. G. Duan, M. M. Ossowski, W. N. Mei, R. W. Smith and I. R. Hardy, "Simulation of Structural Phase Transition in NaNO_3 and CaCO_3 ," *Physical Chemistry of Minerals*, Vol. 28, No. 8, 2001, pp. 586-590.
[doi:10.1007/s002690100191](https://doi.org/10.1007/s002690100191)
- [20] I. Laulicht and N. Lucnar, "Internal Mode Line-Broadening by Proton Jumps in KH_2PO_4 ," *Chemical Physics Letters*, Vol. 47, No. 2, 1977, pp. 237-240.
[doi:10.1016/0009-2614\(77\)80008-0](https://doi.org/10.1016/0009-2614(77)80008-0)
- [21] I. Laulicht, "On the Drastic Temperature Broadening of Hard Mode Raman Lines of Ferroelectric KDP Type Crystals near T_c ," *Journal of Physics and Chemistry of Solids*, Vol. 39, No. 8, 1978, pp. 901-906.
- [doi:10.1016/0022-3697\(78\)90153-1](https://doi.org/10.1016/0022-3697(78)90153-1)
- [22] M. Matsushita, "Anomalous Temperature Dependence of the Frequency and Damping Constant of Phonons near T_λ in Ammonium Halides," *Journal of Chemical Physics*, Vol. 65, No. 1, 1976, pp. 23-28. [doi:10.1063/1.432804](https://doi.org/10.1063/1.432804)
- [23] W. C. K. Poon and E. Salje, "The Excess Optical Birefringence and Phase Transition in Sodium Nitrate," *Journal of Physics C: Solid State Physics*, Vol. 21, No. 4, 1988, pp. 715-729. [doi:10.1088/0022-3719/21/4/009](https://doi.org/10.1088/0022-3719/21/4/009)
- [24] W. W. Schmahl and E. Salje, "X-Ray Diffraction Study of the Orientational Order Disorder Transition in NaNO_3 : Evidence for Order Parameter Coupling," *Physics and Chemistry of Minerals*, Vol. 16, No. 8, 1989, pp. 790-798.
[doi:10.1007/BF00209703](https://doi.org/10.1007/BF00209703)