

Cell Stress by Phosphate of Two Protozoa *Tetrahymena thermophila* and *Tetrahymena pyriformis*

Papa Daouda Mar¹, Bouchra El Khalfi¹, Jose Roman Perez-Castiñeira²,
Aurelio Serrano², Abdelaziz Soukri¹

¹Laboratory of Physiopathology, Genetics Molecular and Biotechnology (PGMB), Faculty of Sciences Ain Chock, University Hassan II of Casablanca, Casablanca, Morocco

²Instituto de Bioquímica Vegetal y Fotosíntesis, CSIC-Universidad de Sevilla, Centro de Investigaciones Científicas “Isla de la Cartuja”, Sevilla, Spain

Email: ab.soukri@gmail.com

How to cite this paper: Mar, P.D., El Khalfi, B., Perez-Castiñeira, J.R., Serrano, A. and Soukri, A. (2017) Cell Stress by Phosphate of Two Protozoa *Tetrahymena thermophila* and *Tetrahymena pyriformis*. *Advances in Bioscience and Biotechnology*, 8, 451-462.

<https://doi.org/10.4236/abb.2017.812033>

Received: September 1, 2017

Accepted: December 3, 2017

Published: December 6, 2017

Copyright © 2017 by authors and Scientific Research Publishing Inc. This work is licensed under the Creative Commons Attribution International License (CC BY 4.0).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

Abstract

Phosphorus is one of the bioelements most needed as a compound cell by living organisms. Phosphorus is involved in several pathologies: in human with bone and kidney diseases, in mammals with metabolism disorder (glucose, insulin...), in microorganisms whose phosphorus is involved in cell growth. Phosphorus has various forms including pyrophosphate, a by-product of multiple pathways of biosynthesis. Enzymes that hydrolyze pyrophosphate are called inorganic pyrophosphatases (PPases). Two major types of inorganic pyrophosphatases are distinguished: the soluble pyrophosphatases (sPPases) and the membrane pyrophosphatases (mPPases or H⁺/Na⁺-PPases). They play a key role in the control of intracellular inorganic pyrophosphate level and produce an important ions gradient (H⁺ or Na⁺) to the cells. In this work, we primarily focused on the physiological study in a phosphate-poor medium of two models *Tetrahymena thermophila* and *Tetrahymena pyriformis*, following the mobility, the growth and the morphology of cells. Secondly, we evaluated the enzymatic activity of soluble and membrane pyrophosphatases in both species grown in the same complex medium. A decrease of cell growth is correlated with unusual morphologies and different mobility in the stress medium. The measurement of soluble and membrane inorganic pyrophosphatases activities also shows a decrease which illustrates the lack of phosphate found in the stress medium. Deficiency of phosphate is a limiting factor for protozoan growth. These results indicate that *Tetrahymena* can be used as a model of cellular stress and consists of a target to study inorganic pyrophosphatases for a better understanding of phosphate cycle in higher organisms.

Keywords

Phosphorus, Pyrophosphatases, NBRIP, Tetrahymena

1. Introduction

Phosphorus is an essential element of cellular metabolism. It is an important component of cells structure, incorporating in the skeleton of molecules to form phosphate groups such as adenosine triphosphate (ATP), nucleic acids (DNA, RNA), phospholipids and phosphoproteins of animals and plants cells. Phosphorus is also a component of bones and teeth, its absorption *in vivo* is ensured by vitamin D and its serum level is regulated by renal reabsorption [1].

In Humans, inorganic phosphate is involved in various pathological disorders such as bone, kidney, and vascular calcification [2]. Indeed, a deficiency of phosphate is found in rickets disease from children, osteomalacia in adults and Toni-Fanconi syndrome which leads to bone loss [3] [4]. In mammals, it is shown that a low phosphate diet alters glucose metabolism [5] [6] and the expression of insulin-sensitive genes [7] [8]. In addition, it is shown that phosphate increases oxidative stress such as Klotho deficiency in mice, causing phosphate retention and impairment of cognition due to increased oxidative damage and apoptosis in Hippocampus neurons [9]. In microorganisms, a deficiency of phosphate in the culture medium induces a cell cycle arrest in *Saccharomyces cerevisiae* [10].

Phosphorus is present in several forms, in nature with its most oxidized form (PO_4^{2-}) and generally bound to metal ions which are components of various rocks and minerals; in living cells whose phosphorus is transported as orthophosphate anion ($\text{H}_2\text{PO}_4^{2-}$), where it is a part of phosphate groups such as pyrophosphate ($\text{PO}_3^{2-}\text{-O-PO}_3^{2-}$). Pyrophosphate is a by-product of numerous synthetic reactions (nucleic acids, proteins, polysaccharides) and plays an important role of cellular metabolism. Consisting of two orthophosphate groups united by a phosphoanhydride bond, pyrophosphate constitutes a chemical grouping, whose fundamental binding energy is stored for energy requirements of the cell. However, intracellular inorganic pyrophosphate is constantly regulated and the hydrolysis of this product indicates that the biosynthetic reactions proceed in the right direction [11] [12]. It is also shown that an intracellular pyrophosphate deregulation is associated with various diseases [13].

Inorganic pyrophosphatases are enzymes that hydrolyze inorganic pyrophosphate into orthophosphate. They are distinguished in two groups: soluble pyrophosphatases (sPPases) and membrane pyrophosphatases (mPPases or $\text{H}^+/\text{Na}^+\text{-PPases}$). Both enzymes are heterologous which have difference in structure and amino acid sequences, but they have similar patterns in catalytic site sequences [14]. Soluble pyrophosphatases are ubiquitous enzymes that play a key role in phosphorus metabolism by catalyzing the hydrolysis of metabolically

produced pyrophosphate. According to their omnipresence, sPPases are important in case of absence an accumulation of pyrophosphate causes metabolic dysfunctions [15] [16]. sPPases consist of three families: family I most frequent and found in most eukaryotic, archaea and bacteria organisms; family II sPPases found in some bacteria, archaea and primitive eukaryotes; the unexplored family III found in some bacterial species [17] [18]. On the one hand, family I sPPases are distinguished from family II sPPases by their activation cofactor: Mg^{2+} is the main cofactor of sPPases whereas Mn^{2+} is the main cofactor of sPPases II [19] [20]. On the other hand, the sequences of both proteins are not similar, nevertheless the active site of the two enzymes remains conserved [19] [21]. In addition Family I sPPases are more sensitive to fluoride than family II sPPases [22] [23].

Soluble pyrophosphatases are essential in metabolic reactions of bacteria and yeasts [16] [24]. They are involved in bacteria growth such as *Escherichia coli*. A dysfunction of sPPases leads to the cell cycle arrest and death of fermenting yeast [25]. Similarly, soluble pyrophosphatases are overexpressed in breast cancer cells [26]. The structure and the function of these enzymes are important in biomedical field because family II pyrophosphatases are specific to certain pathogens such as *Staphylococcus aureus*, *Streptococcus agalactiae*, *Streptococcus mutans* and *Bacillus anthracis*, which may be potential targets for discovery and development of new antibacterial products [27].

Membrane PPases are ionic pumps and found in plants, algae, some protozoa, bacteria, archaea. They produce a proton (H^+ -PPases) or sodium (Na^+ -PPases) gradient by coupling the hydrolysis energy of pyrophosphate [28]. H^+ -PPases are found in prokaryotes (vesicles, acidocalcisomes) [29] [30], protists and eukaryotes (vacuolar plant membranes, acidocalcisomal membranes of protists) [31] [32]. H^+ -PPases provide important energy reserves to organisms, especially under stress conditions and lack of energy [30] [33]. Na^+ -PPases are mainly encountered in prokaryotes.

Membrane pyrophosphatases provide an ion gradient under stress conditions, when ATP is almost absent [30] [34]. These enzymes do not appear to be present in mammals but could be potential targets for vaccines and drugs development against parasitic protists.

However, inorganic pyrophosphatases are rarely studied in protozoa, particularly in ciliates, which are the first used organisms to understand the genetic phenomenon of higher organisms. Ciliates are also characterized by a different evolution from commonly studied organisms in the opisthokont lineage [35]. The protozoan *Tetrahymena* is one of the most studied ciliated organisms and constitutes a eukaryotic cell model. *Tetrahymena thermophila* is a species widely used in cell physiology and molecular genetics studies which have led to the acquisition of much fundamental knowledge in biology [36] [37]. The species *Tetrahymena pyriformis* is also a very useful species in physiological and toxicological studies [38]. The protozoan *Tetrahymena* combines the biological com-

plexity of eukaryotes and the accessibility of unicellular organisms. It is introduced very early into the laboratory due to the easy culture in axenic media, the ideal length for studies under light or electronic microscope (30 - 50 μm).

In this work, we primarily focused on the physiological study in a phosphate-poor medium of two models *Tetrahymena thermophila* and *Tetrahymena pyriformis*, following the mobility, the growth and the morphology of cells. Secondly, we evaluated the enzymatic activity of soluble and membrane pyrophosphatases in both species grown in the same complex medium.

2. Materials and Methods

2.1. Strains and Culture Conditions

The strains *Tetrahymena thermophila* SB1969 and *Tetrahymena pyriformis* used in our study were cultured axenically in the standard medium (PPYE) containing proteose peptone (1.5%, w/v) and yeast extract (0.25%, w/v) [39]. The culture medium was inoculated with 1% (v/v) of *Tetrahymena thermophila* or *Tetrahymena pyriformis* preculture in the same medium. *Tetrahymena thermophila* cells were incubated at 32°C without shaking, while *Tetrahymena pyriformis* cells were incubated at 28°C without shaking. Medium without strains were also prepared in same condition to control eventual contamination.

2.2. NBRIP Medium

National Botanical Research Institute Phosphate medium (NBRIP) [40] was aseptically prepared and represent the stress-medium used in our study. NBRIP is deprived of phosphate and is composed of: $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$ (5 g/l); $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ (0.25 g/l); KCl (0.2 g/l); $(\text{NH}_4)_2\text{SO}_4$ (0.1 g/l); glucose (10 g/l); $\text{Ph} = 7 \pm 0.2$. The culture medium is inoculated with 1% (v/v) of *Tetrahymena thermophila* or *Tetrahymena pyriformis* preculture from PPYE medium and incubated at 32°C (*Tetrahymena thermophila*) or 28°C (*Tetrahymena pyriformis*) without shaking. Medium without strains were also prepared in same condition to control eventual contamination.

2.3. Microscopic Observation

The behavior of both species is analyzed into PPYE and NBRIP culture media under light microscope. This study is carried out by sterily sampling at 3H interval time. Samples are prepared and slide-based, and observed at objective $\times 10$ with Leica DM 500 and A.KRÜSS Optronic microscopes. Protozoan growth is also verified in both culture media by absorbance measurement at 600 nm using Hitachi U-1100 spectrophotometer. The optical zero corresponds to the control media without strain.

2.4. Proteins Extraction by Glass Beads

Cells were harvested at 6000 g for 10 minutes (Beckman Coulter Avanti J-25)

and suspended in Buffer A containing 10 mM Tris-HCl pH8, 10% glycerol (p/v), 1 mM EDTA pH8, 1 mM PMSF, 10 mM β -mercaptoethanol (at ratio of 1 g cells/5 ml Buffer A), in which 1 g of glass beads (0.4 - 0.6 mm of diameter) was added. Cells were broken through the beads by vortexing in 8 cycles (1 minute agitating-1 minute in ice). Then, beads and unbroken cells were removed by centrifugation at 700 g for 10 minutes (Beckman Coulter Allegra™ 21 Centrifuge). Soluble and membrane fractions were obtained by centrifuging the total extract at 120.000 g for 40 minutes (Beckman KL-80 Ultracentrifuge). The supernatant corresponds the soluble crude extract and the pellet was washed in two steps: with Buffer B (60 mM Tris-Hcl pH 8; 12% glycerol; 0.72 M KCl; 1 mM $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$; 1 mM PMSF) and centrifugation at 120.000 g/30 minutes—with Buffer A and centrifugation at 120.000 g/30 minutes. The final obtained pellet was suspended in Buffer A and corresponds to membrane crude extract.

2.5. Determination of Proteins Concentration

Proteins concentration was estimated according to Bradford technique [41]. Bovine Serum Albumin (0.1 mg/ml) was used as standard protein.

2.6. Pyrophosphatases Activity Assay

The determination of enzymatic activity is based on the method established by Rathbun and Betlach [42] which consist to measure released phosphate Pi after the hydrolysis of pyrophosphate PPI. The released phosphate form with molybdate ammonium a phosphomolybdate complex and the latter is reduced by stannous chloride giving a blue color. A unit of enzymatic activity corresponds to the amount of enzyme that hydrolyzes 1 μmol of PPI.

2.6.1. Soluble Pyrophosphatases Activity

Measurement of enzymatic activity was carried out in a reaction mixture (200 μl) containing 50 mM Tris-HCl pH 7.5; 2 mM MgCl_2 ; 0.5 mM Fluoride sodium (optional); adequate volume of soluble crude extract. The reaction was started with addition of 1mM Na_4PPi and incubation at 30°C for 10 minutes. Subsequently the reaction was stopped by adding the stop solution containing 37% Formaldehyde-3.5 M Acetate acid-30% Trichloroacetic acid). Then, ammonium molybdate (0.2% p/v) and stannous chloride (0.16% p/v) were added and absorbance was measured at 735 nm after 10 minutes of incubation.

2.6.2. Membrane Pyrophosphatases Activity

Membrane crude extract was used to realize the measurement of mPPases activity. Thus, reaction mixture (200 μl) was composed of: 50 mM Tris-HCl pH 7.5; 2 mM MgCl_2 ; 0.1 M KCl (optional); adequate volume of sample. The reaction was started with addition of 1 mM Na_4PPi and incubation at 30°C for 10 minutes. The reaction was stopped by adding 670 μl of stop solution, ammonium molybdate and stannous chloride were added and absorbance was measured 10 minutes later at 735 nm.

3. Results and Discussion

3.1. Analysis on Morphology, Mobility and Growth of Protozoa

The behavior of the two species *Tetrahymena thermophila* (TT) and *Tetrahymena pyriformis* (TP) is followed in PPYE and NBRIP culture media as described in material and methods. The cells are inoculated at 1% (v/v) and incubated at 32°C (TT) or 28°C (TP). Samples of each culture medium are taken sterile and three parameters are studied: density, mobility and morphology. The **Table 1** summarizes the results obtained from various observations under light microscope.

There is a high cell density the normal PPYE medium while a low density of protozoa is observed in NBRIP medium. This difference is accompanied by a change in mobility: protozoa move quickly in the PPYE medium and slowly in the NBRIP medium. As regards the cellular structure, the cells observed in the PPYE medium present a pear-shaped characterizing the normal cells of *Tetrahymena* and move very quickly in anarchic direction (**Figure 1**). In NBRIP medium, except of some pear-shaped protozoa that move slowly (**Figure 2(a)**), several cells are found in atypical form and are immobile (**Figure 2(b)**). These parameters measured in both culture media do not show a significant difference between both species TT and TP.




Figure 1. Microscopic image of *Tetrahymena thermophila* in PPYE medium (×10). CV = Contractile Vacuole.

Table 1. Parameters illustrating the behavior of both species *T. thermophila* and *T. pyriformis* according to culture media after 72 hours of incubation.

	<i>Tetrahymena thermophila</i>			<i>Tetrahymena pyriformis</i>		
	Density	Morphology	Mobility	Density	Morphology	Mobility
PPYE	+++	Pear-Shaped	Rapid moving	+++	Pear-Shaped	Rapid moving
NBRIP	+	Atypical Form >>> Pear-Shaped	Slow moving	+	Atypical Form >>> Pear-Shaped	Slow moving

(+: presence; >: higher than).




Figure 2. Microscopic images of *Tetrahymena thermophila* in NBRIP medium ($\times 10$). (a) Pear shaped; (b) Broken cell; (c) Degenerate cell.

Structurally, there is a difference in the architecture of cells vacuoles from PPYE and NBRIP media. The vacuoles are well defined and visible in the normal medium (**Figure 1**) while they are narrower in NBRIP medium (**Figure 2(a)**). Also the contractile vacuole is clearly visible in the PPYE cells and absent in the cells of the NBRIP medium. The contractile vacuole is an osmoregulatory organelle for *Tetrahymena* [43] and his absence may illustrate the disrupting cells observed in NBRIP medium (**Figure 2(c)**).

NBRIP medium is a medium known for its phosphate deficiency. It is generally supplemented with complex phosphate to evaluate the capacity of phosphate solubilizing microorganisms. We found no difference of growth in NBRIP medium or NBRIP supplemented with tricalcium phosphate (TCP), indicating that *Tetrahymena* has no solubilizing activity of complex phosphate. Thus, protozoa are confronted with a medium low in phosphate and limited in nutrients.

Although the growth in NBRIP medium is slowed down and illustrates the low density of the cells observed under an optical microscope, a resistance phenomenon is observed in this medium because viable cells are found even after 144H of culture. This resistance can be explained by the fact that protozoa have polyphosphate reserves [44] which can be used in the event of stress. In addition protozoa have pyrophosphate granules reserves whose appearance depend on calcium and magnesium rate and accumulate during the stationary phase or in case of large amount of phosphate in culture medium [45].

3.2. Enzymatic Activity Assays

Inorganic pyrophosphatases play an important role in the hydrolysis mechanism of inorganic pyrophosphate and their activity depends on cellular living conditions. Enzymatic activity of soluble pyrophosphatases (sPPases) and membrane pyrophosphatases (mPPases) is measured in both species *Tetrahymena thermophila* (TT) and *Tetrahymena pyriformis* (TP), according to standard (PPYE) or complex (NBRIP) media.

In standard medium, mPPases dependent on KCl activity is about 25 mU/mg in both species. In NBRIP medium, enzymatic activity also dependent on KCl decreased to 10 mU/mg (**Figure 3**).

The activity of sPPases under standard medium is in the order of 30 mU/mg in TT and 10 mU/mg in TP whereas in the NBRIP medium the activity of sPPases decreased considerably in TT (4 mU/mg) and TP where it is absent (**Figure 3**). These activities are sensitive to fluoride and seem to correspond to the family I sPPases activity.

Soluble pyrophosphatases play an important role in the regulation of pyrophosphate by preventing its accumulation which disturbs the metabolic reactions [15] [16]. The mPPases constitute ion pumps and use pyrophosphate as a substrate to create a gradient of Na^+ or H^+ ions [28] essential for cellular metabolism. Under stress, cellular metabolism is reduced and a decreased pyrophosphatases activity is normal because of the restriction of metabolic reactions. The absence of soluble pyrophosphatases activity in the NBRIP stress medium results in a small amount of pyrophosphate released through metabolic pathways. Membrane pyrophosphatases activity, however low, ensures the maintenance of ion gradient essential for cell survival. This decline accompanied a reduction of the growth rate of protozoa, the appearance of form of resistance and broken cells by osmotic pressure.




Figure 3. sPPases and mPPases activities of both species *Tetrahymena thermophila* (TT) and *Tetrahymena pyriformis* (TP) in PPYE (TT, TP) and NBRIP (TT + N, TP + N) media.

Phosphate, by its omnipresence in cells of living organisms, is therefore of capital interest particularly in cellular organelles such as mitochondria which are the site of several biosynthetic reactions involving the hydrolysis of pyrophosphate (activation of fatty acids, [46]). The pyrophosphate hydrolysis in these organelles is essentially assured by soluble inorganic pyrophosphatases (sPPases) and a decrease of metabolic reactions as NBRIP medium also shows a decreased activity of these enzymes.

Protozoa survival in the NBRIP medium can be explained by the fact that the cells have reserves of phosphate (polyphosphates, pyrophosphate granules) which are used to ensure their growth and compensate the lack of nutrients in stress medium. It is shown that the hydrolysis of these reserves is not assured by inorganic pyrophosphatases [45] and does not contradict the low pyrophosphatases activity found in NBRIP medium. Nevertheless, a subsequent study of these enzymes involved in the hydrolysis of these reserves would be of great interest for understanding the phosphate cycle from the *Tetrahymena* model.

4. Conclusion

This study shows the importance of phosphate as a cellular compound. Cellular damage that phosphate can generate is elucidated in the protozoa *Tetrahymena thermophila* and *Tetrahymena pyriformis* are considered as unicellular eukaryotic models. Our results indicate that the lack of phosphate affects the morphology, mobility, growth of protozoa and induces a decrease in inorganic pyrophosphatases activity. Proteomics and transcriptomics studies of these enzymes of interest in *Tetrahymena* will be important for further investigations for a better understanding of phosphate cycle in higher organisms.

References

- [1] Soetan, K.O., Olaiya, C.O. and Oyewole, O.E. (2010) The Importance of Mineral Elements for Humans, Domestic Animals and Plants: A Review. *African Journal of Food Science*, **4**, 200-222.
- [2] Kuro-o, M. (2010) A Potential Link between Phosphate and Aging—Lessons from Klotho-Deficient Mice. *Mechanisms of Ageing and Development*, **131**, 270-275. <https://doi.org/10.1016/j.mad.2010.02.008>
- [3] Malhotra, V.K. (1998) *Biochemistry for Students*. 10th Edition, Jaypee Brothers Medical Publishers, (P) Ltd, New Delhi.
- [4] Murray, R.K., Granner, D.K., Mayes, P.A. and Rodwell, V.W. (2000) *Harper's Biochemistry*. 25th Edition, McGraw-Hill, Health Profession Division.
- [5] Wetter, T.J., Gazdag, A.C., Dean, D.J. and Cartee, G.D. (1999) Effect of Calorie Restriction on *In Vivo* Glucose Metabolism by Individual Tissues in Rats. *American Journal of Physiology—Endocrinology and Metabolism*, **276**, E728-E738.
- [6] Lee, C.K., Klopp, R.G., Weindruch, R. and Prolla, T.A. (1999) Gene Expression Profile of Aging and Its Retardation by Caloric Restriction. *Science*, **285**, 1390-1393. <https://doi.org/10.1126/science.285.5432.1390>
- [7] Xie, W., Li, Y., Mechin, M.C. and Van De Werve, G. (1999). Up-Regulation of Liver Glucose-6-Phosphatase in Rats Fed with a P(i)-Deficient Diet. *Biochemical Journal*,

- 343**, 393-396. <https://doi.org/10.1042/bj3430393>
- [8] Xie, W., Tran, T.L., Finegood, D.T. and Van de Werve, G. (2000) Dietary P(i) Deprivation in Rats Affects Liver cAMP, Glycogen, Key Steps of Gluconeogenesis and Glucose Production. *Biochemical Journal*, **352**, 227-232. <https://doi.org/10.1042/bj3520227>
- [9] Nagai, T., Yamada, K., Kim, H.C., Kim, Y.S., Noda, Y., Imura, A., Nabeshima, Y. and Nabeshima, T. (2003) Cognition Impairment in the Genetic Model of Aging Klotho Gene Mutant Mice: A Role of Oxidative Stress. *The FASEB Journal*, **17**, 50-52.
- [10] Boer, V.M., Amini, S. and Botstein, D. (2008) Influence of Genotype and Nutrition on Survival and Metabolism of Starving Yeast. *Proceedings of the National Academy of Sciences U.S.A.*, **105**, 6930-6935. <https://doi.org/10.1073/pnas.0802601105>
- [11] Lahti, R., Pitkäranta, T., Valve, E., Ilta, I., Kukko-Kalske, E. and Heinonen, J. (1988) Cloning and Characterization of the Gene Encoding Inorganic Pyrophosphatase of *Escherichia coli* K-12. *Journal of Bacteriology*, **170**, 5901-5907. <https://doi.org/10.1128/jb.170.12.5901-5907.1988>
- [12] Perez-Castineira, J.R., Lopez-Marques, R.L., Villalba, J.M., Losada, M. and Serrano, A. (2002) Functional Complementation of Yeast Cytosolic Pyrophosphatase by Bacterial and Plant H⁺-Translocating Pyrophosphatases. *Proceedings of the National Academy of Sciences U.S.A.*, **99**, 15914-15919.
- [13] Terkeltaub, R.A. (2001) Inorganic Pyrophosphate Generation and Disposition in Pathophysiology. *American Journal of Physiology. Cell Physiology*, **281**, C1-C11.
- [14] Rea, P.A. and Poole, R.J. (1993) Vacuolar H⁺-Translocating Pyrophosphatase. *Annual Review of Plant Physiology and Plant Molecular Biology*, **44**, 157-180. <https://doi.org/10.1146/annurev.pp.44.060193.001105>
- [15] Kukko-Kalske, E. and Heinonen, J. (1985) Inorganic Pyrophosphate and Inorganic Pyrophosphatase in *Escherichia coli*. *The International Journal of Biochemistry*, **17**, 575-580. [https://doi.org/10.1016/0020-711X\(85\)90288-5](https://doi.org/10.1016/0020-711X(85)90288-5)
- [16] Chen, J., Brevet, A., Fromant, M., Le've'que, F., Schmitter, J.M., Blanquet, S. and Plateau, P. (1990) Pyrophosphatase Is Essential for Growth of *Escherichia coli*. *Journal of Bacteriology*, **172**, 5686-5689. <https://doi.org/10.1128/jb.172.10.5686-5689.1990>
- [17] Lee, H.S., Cho, Y., Kim, Y.J., Lho, T.O., Cha, S.S., Lee, J.H. and Kang, S.G. (2009) A Novel Inorganic Pyrophosphatase in *Thermococcus onnurineus* NA1. *FEMS Microbiology Letters*. **300**, 68-74. <https://doi.org/10.1111/j.1574-6968.2009.01766.x>
- [18] Pratt, A.C., Dewage, S.W., Pang, A.H., Biswas, T., Barnard-Britson, S., Cisneros, G.A. and Tsodikov, O.V. (2015) Structural and Computational Dissection of the Catalytic Mechanism of the Inorganic Pyrophosphatase from *Mycobacterium tuberculosis*. *Journal of Structural Biology*, **192**, 76-87. <https://doi.org/10.1016/j.jsb.2015.08.010>
- [19] Parfenyev, A.N., Salminen, A., Halonen, P., Hachimori, A., Baykov, A.A. and Lahti, R. (2001) Quaternary Structure and Metal Ion Requirement of Family II Pyrophosphatases from *Bacillus subtilis*, *Streptococcus gordonii*, and *Streptococcus mutans*. *The Journal of Biological Chemistry*, **276**, 24511-24518. <https://doi.org/10.1074/jbc.M101829200>
- [20] Zyryanov, A.B., Vener, A.V., Salminen, A., Goldman, A., Lahti, R. and Baykov, A.A. (2004) Rates of Elementary Catalytic Steps for Different Metal Forms of the Family II Pyrophosphatase from *Streptococcus gordonii*. *Biochemistry*, **43**, 1065-1074. <https://doi.org/10.1021/bi0357513>

- [21] Ahn, S., Milner, A.J., Futterer, K., Konopka, M., Ilias, M., *et al.* (2001) The “Open” and “Closed” Structures of the Type-C Inorganic Pyrophosphatases from *Bacillus subtilis* and *Streptococcus gordonii*. *Journal of Molecular Biology*, **313**, 797-811. <https://doi.org/10.1006/jmbi.2001.5070>
- [22] Baykov, A.A., Fabrichniy, I.P., Pohjanjoki, P., Zyryanov, A.B. and Lahti, R. (2000) Fluoride Effects along the Reactions Pathways of Pyrophosphatase: Evidence for a Second Enzyme Center Pyrophosphate Intermediate. *Biochemistry*, **39**, 11939-11947. <https://doi.org/10.1021/bi000627u>
- [23] Kurilova, S.A., Bogdanova, A.V., Nazarova, T.I. and Avaeva, S. (1984) Changes in the *Escherichia coli* Inorganic Pyrophosphatase Activity on Interaction with Magnesium, Zinc, Calcium and Fluoride Ions. *Bioorganicheskaya Khimiya*, **10**, 1153-1160.
- [24] Lundin, M., Baltscheffsky, H. and Ronne, H. (1991) Yeast PPA2 Gene Encodes a Mitochondrial Inorganic Pyrophosphatase That Is Essential for Mitochondrial Function. *The Journal of Biological Chemistry*, **266**, 12166-12168.
- [25] Serrano-Bueno, G., Hernandez, A., Lopez-Lluch, G., Perez-Castineira, J.R., Navas, P. and Serrano, A. (2013) Inorganic Pyrophosphatase Defects Lead to Cell Cycle Arrest and Autophagic Cell Death through NAD⁺ Depletion in Fermenting Yeast. *The Journal of Biological Chemistry*, **288**, 13082-13092. <https://doi.org/10.1074/jbc.M112.439349>
- [26] Hamler, R.L., Zhu, K., Buchanan, N.S., Kreunin, P., Kachman, M.T. and Miller, F.R. (2004) A Two-Dimensional Liquidphase Separation Method Coupled with Mass Spectrometry for Proteomic Studies of Breast Cancer and Biomarker Identification. *Proteomics*, **4**, 562-577. <https://doi.org/10.1002/pmic.200300606>
- [27] Gajadeera, C.S., Zhang, X., Wei, Y. and Tsodikov, O.V. (2015) Structure of Inorganic Pyrophosphatase from *Staphylococcus aureus* Reveals Conformational Flexibility of the Active Site. *Journal of Structural Biology*, **189**, 81-86. <https://doi.org/10.1016/j.jsb.2014.12.003>
- [28] Luoto, H.H., Baykov, A.A., Lahti, R. and Malinen, A.M. (2013) Membrane Integral Pyrophosphatase Subfamily Capable of Translocating Both Na⁺ and H⁺. *Proceedings of the National Academy of Sciences*, **110**, 1255-1260. <https://doi.org/10.1073/pnas.1217816110>
- [29] Perez-Castiñeira, J.R., Lopez-Marques, R.L., Losada, M. and Serrano, A. (2001) A Thermostable K⁺-Stimulated Vacuolar-Type Pyrophosphatase from the Hyperthermophilic Bacterium *Thermotoga maritima*. *FEBS Letters*, **496**, 6-11. [https://doi.org/10.1016/S0014-5793\(01\)02390-0](https://doi.org/10.1016/S0014-5793(01)02390-0)
- [30] Maeshima, M. (2000) Vacuolar H⁺-Pyrophosphatase. *Biochimica et Biophysica Acta*, **1465**, 37-51. [https://doi.org/10.1016/S0005-2736\(00\)00130-9](https://doi.org/10.1016/S0005-2736(00)00130-9)
- [31] Gaxiola, R.A., Palmgren, M.G. and Schumacher, K. (2007) Plant Proton Pumps. *FEBS Letters*, **581**, 2204-2214. <https://doi.org/10.1016/j.febslet.2007.03.050>
- [32] McIntosh, M.T. and Vaidya, A.B. (2002) Vacuolar Type H⁺ Pumping Pyrophosphatases of Parasitic Protozoa. *International Journal for Parasitology*, **32**, 1-14. [https://doi.org/10.1016/S0020-7519\(01\)00325-3](https://doi.org/10.1016/S0020-7519(01)00325-3)
- [33] Serrano, A., Pérez-Castiñeira, J.R., Baltscheffsky, M. and Baltscheffsky, H. (2007) H⁺-P Pases: Yesterday, Today and Tomorrow. *IUBMB Life*, **59**, 76-83. <https://doi.org/10.1080/15216540701258132>
- [34] Garcia-Contreras, R., Celis, H. and Romero, I. (2004) Importance of *Rhodospirillum rubrum* H⁺-Pyrophosphatase under Low-Energy Conditions. *Journal of Bacteriology*, **186**, 6651-6655. <https://doi.org/10.1128/JB.186.19.6651-6655.2004>
- [35] Ruehle, M.D., Orias, E. and Pearson, C.G. (2016) Tetrahymena as a Unicellular

- Model Eukaryote: Genetic and Genomic Tools. *Genetics*, **203**, 649-665.
<https://doi.org/10.1534/genetics.114.169748>
- [36] Leick, V., Bog-Hansen, T.C., Christensen S.T. and Kaufman, S.J. (1996) Concanavalin A Receptors and the Chemosensory Behaviour of *Tetrahymena thermophila*. *Experimental Biology Online*, **1**, 5-10.
<https://doi.org/10.1007/s00898-996-0005-0>
- [37] Tiedtke, A., Rehm, H.J. and Reed, G. (2001) Biotechnology with Protozoa. *Biotechnology*, **10**, 137-149.
- [38] Darcy, P., Kelly, J.P., Leonard, B.E. and Henry, J.A. (2002) The Effect of Lofepamine and Other Related Agents on the Motility of *Tetrahymena pyriformis*. *Toxicology Letters*, **128**, 207-214. [https://doi.org/10.1016/S0378-4274\(01\)00542-2](https://doi.org/10.1016/S0378-4274(01)00542-2)
- [39] Pousada, R.C., Cyrne, M.L. and Hayes, D. (1979) Characterization of Preribosomal Ribonucleoprotein Particles from *Tetrahymena pyriformis*. *European Journal of Biochemistry*, **102**, 389-397. <https://doi.org/10.1111/j.1432-1033.1979.tb04254.x>
- [40] Nautiyal, C.S. (1999) An Efficient Microbiological Growth Medium for Screening Phosphate Solubilizing Microorganisms. *FEMS Microbiology Letters*, **170**, 265-270.
<https://doi.org/10.1111/j.1574-6968.1999.tb13383.x>
- [41] Bradford, M. (1976) A Rapid and Sensitive Method for the Quantitation of Microgram Quantities of Protein Utilizing the Principle of Protein Dye Binding. *Analytical Biochemistry*, **72**, 248-254. [https://doi.org/10.1016/0003-2697\(76\)90527-3](https://doi.org/10.1016/0003-2697(76)90527-3)
- [42] Rathbun, W.B. and Betlach, M.V. (1969) Estimation of Enzymatically Produced Orthophosphate in the Presence of Cysteine and Adenosine Triphosphate. *Analytical Biochemistry*, **28**, 436-445. [https://doi.org/10.1016/0003-2697\(69\)90198-5](https://doi.org/10.1016/0003-2697(69)90198-5)
- [43] Rifkin, J.L. (1973) The Role of the Contractile Vacuole in the Osmoregulation of *Tetrahymena pyriformis*. *The Journal of Protozoology*, **20**, 108-114.
<https://doi.org/10.1111/j.1550-7408.1973.tb06012.x>
- [44] Harold, F.M. (1966) Inorganic Polyphosphates in Biology: Structure, Metabolism, and Function. *Bacteriological Reviews*, **30**, 772-794.
- [45] Munk, N. and Rosenberg, H. (1968) On the Deposition and Utilization of Inorganic Pyrophosphate in *tetrahymena pyriformis*. *Biochimica et Biophysica Acta*, **177**, 629-640. [https://doi.org/10.1016/0304-4165\(69\)90329-8](https://doi.org/10.1016/0304-4165(69)90329-8)
- [46] Mansurova, S.E. (1989) Inorganic Pyrophosphate in Mitochondrial Metabolism. *Biochimica et Biophysica Acta*, **977**, 237-247.
[https://doi.org/10.1016/S0005-2728\(89\)80078-7](https://doi.org/10.1016/S0005-2728(89)80078-7)