

Dividend Payments and Related Problems in a Markov-Dependent Insurance Risk Model under Absolute Ruin

Wenguang Yu¹, Yujuan Huang²

¹School of Statistics and Mathematics, Shandong University of Finance and Economics, Jinan, China; ²Department of Mathematics and Physics, Shandong Jiaotong University, Jinan, China.
Email: yuwg@mail.sdu.edu.cn, yujuanh518@163.com

Received September 5th, 2011; revised September 19th, 2011; accepted September 29th, 2011.

ABSTRACT

In this paper, we study the dividend payments prior to absolute ruin in a Markov-dependent risk process in which the claim occurrence and the claim amount are regulated by an external discrete time Markov chain. A system of integro-differential equations with boundary conditions satisfied by the moment-generating function, the n th moment of the discounted dividend payments prior to absolute ruin and the discounted penalty function, given the initial environment state, are derived. In the two-state risk model, explicit solutions to the integro-differential equations satisfied by the n th moment of the discounted dividend payments prior to absolute ruin are obtained when the claim size distribution is exponentially distributed. Finally, the matrix form of systems of integro-differential equations satisfied by the discounted penalty function are presented.

Keywords: Absolute Ruin, Markov-Dependent Insurance Risk Model, Debit Interest, Moment-Generating Function

1. Introduction

Recently ruin theory under regime-switching model is becoming a popular topic. This model is proposed in Reinhard [1] and Asmussen [2]. Asmussen [2] calls it a Markov-modulated risk model in which both the frequency of the claim arrivals and the distribution of the claim amounts are influenced by an external environment process. This model is a generalization of the classical compound Poisson risk model and the primary motivation for this generalization is the enhanced flexibility that it permits for the modeling of the claim arrival process and the claim severity distribution assumed in the classical risk process. The model builds a Markov chain whose states represent different states of an economy into the insurance risk model. The regime-switching of the states of the economy can be attributed to the structural changes in the (macro-) economic conditions, the changes in political regimes, the impact of (macro-) economic news and business cycles, etc. There are many papers published on ruin probabilities and the related problems under the Markov-modulated (or Markov regime-switching) risk model. Ng and Yang [3] give closed form solutions for the joint distribution of the surplus before and after

ruin when the initial surplus is zero or when the claim amount distributions are phase-type distributed. Li and Lu [4] study the moments of the present value of the dividend payments and the distribution of the total dividends prior to ruin for the Markov-modulated risk model modified by the introduction of a barrier dividend. Lu and Li [5] and Liu *et al.* [6] consider a regime-switching risk model with a threshold dividend strategy. Zhu and Yang [7] study a more general Markovian regime-switching risk model in which the premium, the claim intensity, the claim amount, the dividend payment rate and the dividend threshold level are influenced by an external Markovian environment process. Wei *et al.* [8] consider the Markov-modulated insurance risk model with tax.

Moreover, in recent years, semi-Markovian risk model has attracted attention in the literature. Albrecher and Boxma [9] study the expected discounted penalty function in a semi-Markovian dependent risk model in which at each instant of a claim, the underlying Markov chain jumps to a new state and the distribution of claim depends on this state. Liu *et al.* [10,11] consider the expected discounted penalty function under the constant

dividend barrier and dividends payments under the threshold strategy in a Markov-dependent risk model, respectively. They consider the structure of a semi-Markovian dependence type as follows. Let W_i denote the time between the arrival of the $(i-1)$ th and the i th claims and $W_0 = X_0$ a.s., then

$$\begin{aligned} & \Pr[W_{n+1} \leq x, X_{n+1} \leq y, Z_{n+1} = j | Z_n = i, \\ & \quad (W_s, X_s, Z_s), 0 \leq s \leq n] \\ &= \Pr(W_1 \leq x, X_1 \leq y, Z_1 = j | Z_0 = i) \\ &= (1 - e^{-\lambda_1 x}) p_{ij} F_j(y) \end{aligned}$$

where $\{Z_n; n \geq 0\}$ is an irreducible discrete time Markov chain with state space $S = \{1, 2, 3, \dots, m\}$ and transition matrix $\Lambda = (p_{i,j})_{i,j=1}^m$, X_n is the amount of the n th claim.

Thus at each instant of a claim, the Markov chain jumps to a state j and the distribution F_j of the claim depends on the new state j , and has a positive mean μ_j . Then, the next interarrival time is exponentially distributed with parameter λ_j . Note that given the states Z_{n-1} and Z_n , the quantities W_n and X_n are independent, but there is an autocorrelation among consecutive claim sizes and among consecutive interclaim times as well as cross-correlation between W_n and X_n .

Inspired by Albrecher and Boxma [9] and Liu *et al.* [10,11], in this paper we propose to generalize the semi-Markovian risk model to the absolute ruin risk model. In the new risk model, we assume that the insurer could borrow an amount of money equal to the deficit at a debit interest force β when the surplus is negative. Meanwhile, the insurer will repay the debts continuously from his/her premium income. When the negative surplus attains the level $-c/\beta$ or is below $-c/\beta$, the surplus is no longer able to be positive. Absolute ruin occurs at this moment. Moreover, when the surplus exceeds the constant barrier $b(\geq u)$, dividends are paid continuously so the surplus stays at the level b until a new claim occurs. Some recent references about absolute ruin risk model include Zhou and Zhang [12], Cai [13], Gerber and Yang [14], Yuen *et al.* [15], Yuan and Hu [16], Wang and Yin [17], Ming *et al.* [18], Wang *et al.* [19], Zhang *et al.* [20], Yu and Huang [21] and references therein.

The surplus process $\{U_b(t); t \geq 0\}$ under the Markov-dependent risk model is given by

$$dU_b(t) = [c + \beta U(t) I(U(t) < 0)] dt - d\left(\sum_{n=1}^{N(t)} X_n\right)$$

where $U(0) = u$ is the initial surplus, c the premium rate, β the debit interest, $\{N(t); t \geq 0\}$ the number of claims up to time t , and $I(B)$ means the indicator function of an event B . Furthermore, we assume the net profit condi-

tion holds, that is

$$\sum_{i=1}^m \pi_i \mu_i < c \sum_{i=1}^m \pi_i \frac{1}{\lambda_i}$$

where $\pi = (\pi_1, \pi_2, \dots, \pi_m)$ is the stationary distribution of process $\{Z_n, n \geq 0\}$.

Let $D(t)$ be the cumulative amount of dividends paid out up to time t and $\delta > 0$ the force of interest, then

$$D_{u,b} = \int_0^{T_b} e^{-\delta t} dD(t)$$

is the present value of all dividends until time of ruin T_b , where T_b denoted by $T_b = \inf\{t \geq 0 : U_b(t) \leq -c/\beta\}$ is the time of absolute ruin.

In the sequel we will be interested in the moment-generating function

$$M_i(u, y; b) = E[e^{yD_{u,b}} | Z_0 = i], \quad i \in S$$

and the n th moment function

$$V_{n,i}(u; b) = E[D_{u,b}^n | Z_0 = i], \quad n \in N, \quad i \in S$$

with $V_{0,i}(u; b) = 1$, and the expected discounted penalty function, for $i \in S$

$$\begin{aligned} \Phi_i(u, b) &= E[e^{-\delta T_b} \omega(U_b(T_b^-), |U_b(T_b)|)] \\ &\quad \times I(T_b < \infty) | U_b(0) = u, Z_0 = i \end{aligned}$$

where, $U_b(T_b^-)$ is the surplus prior to absolute ruin and $|U_b(T_b)|$ is the deficit at absolute ruin. The penalty function $\omega(x_1, x_2)$ is an arbitrary nonnegative measurable function defined on $(-c/\beta, +\infty) \times (c/\beta, +\infty)$. Throughout this paper we assume that $M_i(u, y; b)$, $V_{n,i}(u; b)$ and $\Phi_i(u, b)$ are sufficiently smooth functions in u and y , respectively.

Then, fix $n \in N$, the expected present value of the total dividend payments until ruin in the stationary case is given by

$$V_n(u, b) = \sum_{i=1}^m \pi_i V_{n,i}(u, b)$$

The rest of the paper is organized as follows. In Sections 2, we get integro-differential equations for the moment-generating function and boundary conditions in a Markov-dependent risk model. In section 3, the integro-differential equations satisfied by higher moment of the dividend payments and boundary conditions are derived. Examples for a two-state risk model are illustrated in section 4 when the claim size distribution is exponentially distributed. In the last section, we obtain the systems of integro-differential equations for the discounted penalty function and its matrix form.

2. Moment-Generating Function of $D_{u,b}$

In this section, we discuss the integro-differential equations satisfied by the moment-generating function at absolute ruin. We point out that $M_i(u, y; b)$ has different paths for $0 \leq u < b$ and $-c/\beta < u \leq 0$. For $i \in S$, we

$$c \frac{\partial M_{1,i}(u, y; b)}{\partial u} = \delta y \frac{\partial M_{1,i}(u, y; b)}{\partial y} + \lambda_i M_{1,i}(u, y; b) + \lambda_i \sum_{j=1}^m p_{i,j} \left[\int_0^u M_{1,j}(u-x, y; b) dF_j(x) + \int_u^{u+\frac{c}{\beta}} M_{2,j}(u-x, y; b) dF_j(x) + \bar{F}_j(u+c/\beta) \right] \tag{2.1}$$

and, for $-c/\beta < u \leq 0$,

$$(\beta u + c) \frac{\partial M_{2,i}(u, y; b)}{\partial u} = \delta y \frac{\partial M_{2,i}(u, y; b)}{\partial y} + \lambda_i M_{2,i}(u, y; b) - \lambda_i \sum_{j=1}^m p_{i,j} \left[\int_0^{u+\frac{c}{\beta}} M_{2,j}(u-x, y; b) dF_j(x) + \bar{F}_j(u+c/\beta) \right] \tag{2.2}$$

with boundary conditions, for $i \in S$,

$$\left. \frac{\partial M_{1,i}(u, y; b)}{\partial u} \right|_{u=b} = y M_{1,i}(b, y; b) \tag{2.3}$$

$$M_{2,i}(-c/\beta, y; b) = 1 \tag{2.4}$$

where, $\bar{F}_j(x) = 1 - F_j(x)$.

Proof. Fix $i \in S$, and $0 \leq u \leq b$. Considering a small

define

$$M_i(u, y; b) = \begin{cases} M_{1,i}(u, y; b), & 0 \leq u < b \\ M_{2,i}(u, y; b), & -c/\beta < u \leq 0 \end{cases}$$

Theorem 2.1 For $i \in S$, $0 \leq u \leq b$, we have

time interval $[0, t]$, such that $u + ct < b$. In view of the strong Markov property of the surplus process $\{U_b(t), t \geq 0\}$, we have

$$M_i(u, y; b) = E \left[M_i(U_b(t), ye^{-\alpha t}; b) \right] \tag{2.5}$$

Conditioning on the event occurring in the interval $[0, t]$, we obtain

$$M_{1,i}(u, y; b) = (1 - \lambda_i t) M_{1,i}(u + ct, ye^{-\delta t}; b) + \lambda_i t \sum_{j=1}^m p_{i,j} \left[\int_0^{u+ct} M_{1,j}(u+ct-x, ye^{-\delta t}; b) dF_j(x) + \int_{u+ct}^{u+ct+\frac{c}{\beta}} M_{2,j}(u+ct-x, ye^{-\alpha t}; b) dF_j(x) + \bar{F}_j(u+ct+c/\beta) \right] + o(t) \tag{2.6}$$

Taylor's expansion gives

$$M_{1,i}(u + ct, ye^{-\delta t}; b) = M_{1,i}(u, y; b) + ct \frac{\partial M_{1,i}(u, y; b)}{\partial u} - \delta y t \frac{\partial M_{1,i}(u, y; b)}{\partial y} + o(t) \tag{2.7}$$

Substituting (2.7) into (2.6), dividing both sides by t , and letting $t \rightarrow 0$, we obtain (2.1).

Similarly, when $-c/\beta < u \leq 0$, we still consider a small time interval $[0, t]$, with $t(t > 0)$ being sufficiently small so that the surplus will not reach 0 in the time interval. Let t_0 be the solution to

$$h_\beta(u, t) = ue^{\beta t} + c(e^{\beta t} - 1)/\beta = 0$$

Then $h_\beta(u, t)$ is the surplus at time $t \leq t_0$ if no claim occurs prior to time t_0 . We assume $t \leq t_0$. So conditioning on the time and the amount of the first claim, we have

$$M_{2,i}(u, y; b) = (1 - \lambda_i t) M_{2,i}(h_\beta(u, t), ye^{-\delta t}; b) + \lambda_i t \sum_{j=1}^m p_{i,j} \left[\int_0^{h_\beta(u,t)+\frac{c}{\beta}} M_{2,j}(h_\beta(u,t)-x, ye^{-\delta t}; b) dF_j(x) + \bar{F}_j(h_\beta(u,t)+c/\beta) \right] + o(t) \tag{2.8}$$

By Taylor's expansion

$$M_{2,i}(h_\beta(u, t), ye^{-\delta t}; b) = M_{2,i}(u, y; b) + (\beta u + c)t \frac{\partial M_{2,i}(u, y; b)}{\partial u} - \delta y t \frac{\partial M_{2,i}(u, y; b)}{\partial y} + o(t) \tag{2.9}$$

Substituting (2.9) into (2.8), dividing both sides by t , and letting $t \rightarrow 0$, we obtain (2.2).

When the initial surplus is b , we obtain

$$M_{1,i}(b, y; b) = (1 - \lambda_i t) e^{yct} M_{1,i}(b, ye^{-\delta t}; b) + \lambda_i t e^{yct} \sum_{j=1}^m p_{i,j} \left[\int_0^b M_{1,j}(b-x, ye^{-\delta t}; b) dF_j(x) + \int_b^{b+\frac{c}{\beta}} M_{2,j}(b-x, ye^{-\delta t}; b) dF_j(x) + \bar{F}_j(b+c/\beta) \right] + o(t) \tag{2.10}$$

Using Taylor's expansion, we have,

$$\delta y \frac{\partial M_{1,i}(b, y; b)}{\partial y} + (\lambda_i - cy) M_{1,i}(b, y; b) = \lambda_i \sum_{j=1}^m p_{i,j} \left[\int_0^b M_{1,j}(b-x, ye^{-\delta t}; b) dF_j(x) \right] = \lambda_i \sum_{j=1}^m p_{i,j} \left[\int_0^b M_{1,j}(b-x, ye^{-\delta t}; b) dF_j(x) + \bar{F}_j(b+c/\beta) \right] + o(t) \tag{2.11}$$

Letting $u \uparrow b$ in (2.1) and comparing it with (2.11), we obtain (2.3). Where “ \uparrow ” denoting increasing approach.

When $u = -c/\beta$, absolute ruin is immediate. Thus, no dividend is paid. So we obtain (2.4). Theorem 2.1 is proved.

Theorem 2.2 For $i \in S$,

$$M_{1,i}(0+, y; b) = M_{2,i}(0-, y; b) \tag{2.12}$$

Proof. For $-c/\beta < u \leq 0$, letting τ_0 be the time that the surplus reach 0 for the first time from $u < 0$ and using the Markov property of the surplus process, we obtain

$$\begin{aligned} M_{2,i}(u, y; b) &= E_i^u \left[I(\tau_0 < T_b) e^{yD_{u,b}} \right] + E_i^u \left[I(\tau_0 \geq T_b) e^{yD_{u,b}} \right] \\ &= E_i^u \left[I(\tau_0 < T_b) \exp \left\{ y \int_0^{T_b - \tau_0} e^{-\delta t} dD(t + \tau_0) \right\} \right] + P(\tau_0 \geq T_b) \\ &= E_i^u \left[I(\tau_0 < T_b) \exp \left\{ ye^{-\delta \tau_0} \int_{\tau_0}^{T_b} e^{-\delta t} dD(t) \right\} \right] + P(\tau_0 \geq T_b) \\ &\leq M_{1,i}(0, y; b) + P(\tau_0 \geq T_b) \end{aligned} \tag{2.13}$$

Similarly, we obtain

$$\begin{aligned} M_{2,i}(u, y; b) &\geq E_i^u \left[I(\tau_0 < T_b, \tau_0 = t_0) e^{yD_{u,b}} \right] + E_i^u \left[I(\tau_0 \geq T_b) e^{yD_{u,b}} \right] \\ &= M_{1,i}(0, y; b) E_i^u \left[e^{-\delta \tau_0} I(\tau_0 < T_b, \tau_0 = t_0) \right] + P(\tau_0 \geq T_b) \\ &= M_{1,i}(0, y; b) e^{-\delta \tau_0} P(T_1 > t_0) + P(\tau_0 \geq T_b) \\ &= e^{-(\lambda_i + \delta)t_0} M_{1,i}(0, y; b) + P(\tau_0 \geq T_b) \end{aligned} \tag{2.14}$$

where T_1 is the time of the first claim.

When $u \uparrow 0$, we notice that τ_0 and t_0 both go into zero. Letting $u \uparrow 0$ in (2.13) and (2.14) and in view of

$$\lim_{u \uparrow 0} P(\tau_0 \geq T_b) = 0$$

we obtain (2.12). Theorem 2.2 is proved.

3. Higher Moment of the Dividend Payments

We now derive a system of integro-differential equations satisfied by $V(u, b)$. By the definitions of $M(u, y; b)$ and $V(u, b)$, we obtain, for $i \in S$,

$$M_{1,i}(u, y; b) = 1 + \sum_{n=1}^{\infty} \frac{y^n}{n!} V_{n,1,i}(u, b) \tag{3.1}$$

$$M_{2,i}(u, y; b) = 1 + \sum_{n=1}^{\infty} \frac{y^n}{n!} V_{n,2,i}(u, b) \tag{3.2}$$

where, $V_{n,i}(u; b)$ is defined by

$$V_{n,i}(u; b) = \begin{cases} V_{n,1,i}(u; b), & 0 \leq u < b \\ V_{n,2,i}(u; b), & -c/\beta < u \leq 0 \end{cases}$$

Substituting (3.1) and (3.2) into (2.1) and (2.2), respectively, and comparing the coefficients of y^n yield the following integro-differential equations:

$$cV'_{n,1,i}(u, b) = (\lambda_i + n\delta)V_{n,1,i}(u, b) - \lambda_i \sum_{j=1}^m p_{i,j} \left[\int_0^u V_{n,1,j}(u-x, b) dF_j(x) + \int_u^{u+\frac{c}{\beta}} V_{n,2,j}(u-x, b) dF_j(x) \right] \tag{3.3}$$

for $0 \leq u < b$, and for $-c/\beta < u \leq 0$,

$$(\beta u + c)V'_{n,2,i}(u, b) = (\lambda_i + n\delta)V_{n,2,i}(u, b) - \lambda_i \sum_{j=1}^m p_{i,j} \int_0^{u+\frac{c}{\beta}} V_{n,2,j}(u-x, b) dF_j(x) \tag{3.4}$$

Substituting (3.1) into (2.3), similarly, we obtain

$$V'_{n,1,i}(u, b)|_{u=b} = nV_{n-1,1,i}(b, b) \tag{3.5}$$

Thus, $V_{1,1,i}(b, b) = 1$ is an obvious result since $V_{0,1,i}(b, b) = 1$.

Substituting (3.1) and (3.2) into (2.4) and (2.12), we obtain, for $n \in N^+$

$$V_{n,2,i}(-c/\beta, b) = 0 \tag{3.6}$$

$$V_{n,1,i}(0+, b) = V_{n,2,i}(0-, b) \tag{3.7}$$

Letting $u \downarrow 0$ in (3.3) and $u \uparrow 0$ in (3.4) and using (3.7), we obtain, for $n \in N^+$

$$V'_{n,1,i}(0+, b) = V'_{n,2,i}(0-, b) \tag{3.8}$$

where, “ \downarrow ” denoting decreasing approach.

4. Explicit Expressions for Exponential Claims to $V_n(u, b)$ for a Two-State Model

In this section, we consider a two-state Markov-dependent risk model. Then $\{Z_n; n \geq 0\}$ is a two-state Markov chain, which reflects the random environmental effects due to “normal” vs. “abnormal”, or “high season” vs. “low season” conditions. We derive the explicit formulae for $V_n(u, b)$ when the claim size is exponentially distributed $F_i(x) = 1 - e^{-x/\mu_i}$, $i = 1, 2$. Set $p_{11} = p_{22} = 0$, $p_{12} = p_{21} = 1$. In view of Equation (3.3) and the exponential density function, Equation (3.3) are reduced to, for $0 \leq u \leq b$

$$cV'_{n,1,1}(u, b) = (\lambda_1 + n\delta)V_{n,1,1}(u, b) - \frac{\lambda_1}{\mu_2} e^{-u/\mu_2} \int_0^u V_{n,1,2}(x, b) e^{x/\mu_2} dx \tag{4.1}$$

$$cV'_{n,1,2}(u, b) = (\lambda_2 + n\delta)V_{n,1,2}(u, b) - \frac{\lambda_2}{\mu_1} e^{-u/\mu_1} \int_0^u V_{n,1,1}(x, b) e^{x/\mu_1} dx - \frac{\lambda_2}{\mu_1} e^{-u/\mu_1} \int_{-\frac{c}{\beta}}^0 V_{n,2,1}(x, b) e^{x/\mu_1} dx \tag{4.2}$$

Applying the operator $\left(\frac{d}{du} + \frac{1}{\mu_2}\right)$ and $\left(\frac{d}{du} + \frac{1}{\mu_1}\right)$ on (4.1) and (4.2), respectively, and rearranging them, we obtain, for $0 \leq u \leq b$

$$V''_{n,1,1}(u, b) + \left(\frac{1}{\mu_2} - \frac{\lambda_1 + n\delta}{c}\right)V'_{n,1,1}(u, b) - \frac{\lambda_1 + n\delta}{\mu_2 c} V_{n,1,1}(u, b) = -\frac{\lambda_1}{\mu_2 c} V_{n,1,1}(u, b) \tag{4.3}$$

$$V''_{n,1,2}(u, b) + \left(\frac{1}{\mu_1} - \frac{\lambda_2 + n\delta}{c}\right)V'_{n,1,2}(u, b) - \frac{\lambda_2 + n\delta}{\mu_1 c} V_{n,1,2}(u, b) = -\frac{\lambda_2}{\mu_1 c} V_{n,1,1}(u, b) \tag{4.4}$$

They are second-order linear non-homogeneous differential equations with constant coefficients. For convenient writing, let

$$q_1 = \frac{1}{\mu_2} - \frac{\lambda_1 + n\delta}{c}, \quad q_2 = -\frac{\lambda_1 + n\delta}{\mu_2 c}$$

$$\eta_1 = \frac{1}{\mu_1} - \frac{\lambda_2 + n\delta}{c}, \quad \eta_2 = -\frac{\lambda_2 + n\delta}{\mu_1 c}$$

$$g_1(u) = -\frac{\lambda_1 V_{n,1,2}(u, b)}{\mu_2 c}, \quad g_2(u) = -\frac{\lambda_2 V_{n,1,1}(u, b)}{\mu_1 c}$$

Then Equation (4.3) and Equation (4.4) can be rewritten as

$$V''_{n,1,1}(u, b) + q_1 V'_{n,1,1}(u, b) + q_2 V_{n,1,1}(u, b) = g_1(u) \tag{4.5}$$

$$V''_{n,1,2}(u, b) + \eta_1 V'_{n,1,2}(u, b) + \eta_2 V_{n,1,2}(u, b) = g_2(u) \tag{4.6}$$

The corresponding homogeneous differential Equations of (4.5) and (4.6) are

$$V''_{n,1,1}(u, b) + q_1 V'_{n,1,1}(u, b) + q_2 V_{n,1,1}(u, b) = 0 \tag{4.7}$$

$$V''_{n,1,2}(u, b) + \eta_1 V'_{n,1,2}(u, b) + \eta_2 V_{n,1,2}(u, b) = 0 \tag{4.8}$$

The general solutions of Equations (4.7) and (4.8) are

$$V_{n,1,1}(u, b) = c_1 e^{\eta_1 u} + c_2 e^{\eta_2 u} \tag{4.9}$$

$$V_{n,1,2}(u, b) = c_3 e^{\eta_1 u} + c_4 e^{\eta_2 u} \tag{4.10}$$

where c_1, c_2, c_3, c_4 are arbitrary constants,

$$r_1 = \frac{-q_1 + \sqrt{q_1^2 - 4q_2}}{2}$$

$$r_2 = \frac{-q_1 - \sqrt{q_1^2 - 4q_2}}{2}$$

$$\eta_1 = \frac{-q_3 + \sqrt{q_3^2 - 4q_4}}{2}$$

$$\eta_2 = \frac{-q_3 - \sqrt{q_3^2 - 4q_4}}{2}$$

According to the variation of constants method, we assume $c_1(u)e^{\eta_1 u} + c_2(u)e^{\eta_2 u}$ and $c_3(u)e^{\eta_1 u} + c_4(u)e^{\eta_2 u}$ are special solutions of Equations (4.5) and (4.6), respectively. Then we have

$$\begin{cases} c_1'(u)e^{\eta_1 u} + c_2'(u)e^{\eta_2 u} = 0 \\ c_1'(u)r_1e^{\eta_1 u} + c_2'(u)r_2e^{\eta_2 u} = g_1(u) \end{cases} \quad (4.11)$$

$$\begin{cases} c_3'(u)e^{\eta_1 u} + c_4'(u)e^{\eta_2 u} = 0 \\ c_3'(u)\eta_1e^{\eta_1 u} + c_4'(u)\eta_2e^{\eta_2 u} = g_2(u) \end{cases} \quad (4.12)$$

Solving the above two equations, we obtain

$$c_1'(u) = g_1(u)(r_1 - r_2)e^{-\eta_1 u}$$

$$c_2'(u) = g_1(u)(r_2 - r_1)e^{-\eta_2 u}$$

$$c_3'(u) = g_2(u)(\eta_1 - \eta_2)e^{-\eta_1 u}$$

$$c_4'(u) = g_2(u)(\eta_2 - \eta_1)e^{-\eta_2 u}$$

then we have, for $0 \leq u \leq b$

$$c_1(u) = (r_1 - r_2) \int_0^u g_1(x) e^{-\eta_1 x} dx$$

$$c\Phi'_{1,i}(u, b) = (\lambda_i + \delta)\Phi_{1,i}(u, b) - \lambda_i \sum_{j=1}^m p_{i,j} \left[\int_0^u \Phi_{1,j}(u-x, b) dF_j(x) + \int_u^{u+\frac{c}{\beta}} \Phi_{2,j}(u-x, b) dF_j(x) + A_j(u) \right] \quad (5.1)$$

and, for $-c/\beta < u \leq 0$,

$$(\beta u + c)\Phi'_{2,i}(u, b) = (\lambda_i + \delta)\Phi_{2,i}(u, b) - \lambda_i \sum_{j=1}^m p_{i,j} \int_0^{u+\frac{c}{\beta}} \Phi_{2,j}(u-x, b) dF_j(x) + A_j(u) \quad (5.2)$$

with boundary conditions

$$\Phi'_{1,i}(b, b) = 0 \quad (5.3)$$

$$\Phi_{1,i}(0+, b) = \Phi_{2,i}(0-, b) \quad (5.4)$$

$$\Phi'_{1,i}(0+, b) = \Phi'_{2,i}(0-, b) \quad (5.5)$$

$$\begin{aligned} \Phi_{1,i}(u, b) = & (1 - \lambda_i t) e^{-\delta t} \Phi_{1,i}(u + ct, b) + \lambda_i t e^{-\delta t} \sum_{j=1}^m p_{i,j} \left[\int_0^{u+ct} \Phi_{1,j}(u+ct-x, b) dF_j(x) \right. \\ & \left. + \int_{u+ct}^{u+ct+\frac{c}{\beta}} \Phi_{2,j}(u+ct-x, b) dF_j(x) + \int_{u+\frac{c}{\beta}}^{\infty} \omega(u, x-u) dF_j(x) \right] + o(t) \end{aligned} \quad (5.6)$$

$$c_2(u) = (r_2 - r_1) \int_0^u g_1(x) e^{-r_2 x} dx$$

$$c_3(u) = (\eta_1 - \eta_2) \int_0^u g_2(x) e^{-\eta_1 x} dx$$

$$c_4(u) = (\eta_2 - \eta_1) \int_0^u g_2(x) e^{-\eta_2 x} dx$$

So the general solutions of Equations (4.5) and (4.6) are, for $0 \leq u \leq b$,

$$\begin{aligned} V_{n,1,1}(u, b) = & (r_1 - r_2) e^{\eta_1 u} \int_0^u g_1(x) e^{-\eta_1 x} dx \\ & + (r_2 - r_1) e^{\eta_2 u} \int_0^u g_1(x) e^{-\eta_2 x} dx + c_1 e^{\eta_1 u} + c_2 e^{\eta_2 u} \end{aligned} \quad (4.13)$$

$$\begin{aligned} V_{n,1,2}(u, b) = & (\eta_1 - \eta_2) e^{\eta_1 u} \int_0^u g_2(x) e^{-\eta_1 x} dx \\ & + (\eta_2 - \eta_1) e^{\eta_2 u} \int_0^u g_2(x) e^{-\eta_2 x} dx + c_3 e^{\eta_1 u} + c_4 e^{\eta_2 u} \end{aligned} \quad (4.14)$$

5. The Discounted Penalty Function

In this section, we derive integro-differential equations for the discounted penalty functions. For $i \in S$, define

$$\Phi_i(u, b) = \begin{cases} \Phi_{1,i}(u, b), & 0 \leq u < b \\ \Phi_{2,i}(u, b), & -c/\beta < u \leq 0 \end{cases}$$

Note that in the stationary case, we have

$$\Phi(u, b) = \sum_{i=1}^m \pi_i \Phi_i(u, b)$$

Theorem 5.1 For $i \in S$, $0 \leq u \leq b$,

where

$$A_i(u) = \int_{u+\frac{c}{\beta}}^{\infty} \omega(u, x-u) dF_i(x)$$

Proof. For $i \in S$ and $0 \leq u \leq b$. Similar to argument as in Section 2, we condition on the events that can occur in the small time interval $[0, t]$.

Since

$$e^{-\delta t} = 1 - \delta h + o(h)$$

we then get

$$\begin{aligned} \Phi_{1,i}(u, b) = & \left[1 - (\lambda_i + \delta)t \right] \Phi_{1,i}(u + ct, b) + \lambda_i t \sum_{j=1}^m p_{i,j} \int_0^{u+ct} \Phi_{1,j}(u + ct - x, b) dF_j(x) \\ & + \int_{u+ct}^{u+ct+\frac{c}{\beta}} \Phi_{2,j}(u + ct - x, b) dF_j(x) + \int_{u+\frac{c}{\beta}}^{\infty} \omega(u, x - u) dF_j(x) \Big] + o(t) \end{aligned} \tag{5.7}$$

Equation (5.7) can be rewritten as

$$\begin{aligned} \frac{\Phi_{1,i}(u + ct, b) - \Phi_{1,i}(u, b)}{t} = & (\lambda_i + \delta) \Phi_{1,i}(u + ct, b) - \lambda_i \sum_{j=1}^m p_{i,j} \left[\int_0^{u+ct} \Phi_{1,j}(u + ct - x, b) dF_j(x) \right. \\ & \left. + \int_{u+ct}^{u+ct+\frac{c}{\beta}} \Phi_{2,j}(u + ct - x, b) dF_j(x) + \int_{u+\frac{c}{\beta}}^{\infty} \omega(u, x - u) dF_j(x) \right] + \frac{o(t)}{t} \end{aligned} \tag{5.8}$$

Letting $t \rightarrow 0$ in (5.8), we obtain (5.1).

For $i \in S$ and $-c/\beta < u \leq 0$, we have

$$\begin{aligned} \Phi_{2,i}(u, b) = & (1 - \lambda_i t) e^{-\delta t} \Phi_{2,i}(h_\beta(u, t), b) + \lambda_i t e^{-\delta t} \\ & \times \sum_{j=1}^m p_{i,j} \left[\int_0^{h_\beta(u, t) + \frac{c}{\beta}} \Phi_{2,j}(h_\beta(u, t) - x, b) dF_j(x) + \int_{h_\beta(u, t) + \frac{c}{\beta}}^{\infty} \omega(h_\beta(u, t), x - h_\beta(u, t)) dF_j(x) \right] + o(t) \end{aligned} \tag{5.9}$$

By Taylor's expansion

$$\begin{aligned} \Phi_{2,i}(h_\beta(u, t), b) = & \Phi_{2,i}(u, b) \\ & + (\beta u + c)t \Phi'_{2,i}(u, b) + o(t) \end{aligned} \tag{5.10}$$

Substituting (5.10) into (5.9), dividing both sides by t , and letting $t \rightarrow 0$, we obtain (5.2). Theorem 5.1 is proved.

Integro-differential Equations (5.1) and (5.2) can easily be rewritten in matrix form.

Let

$$\Phi_j(u, b) = (\Phi_{j,1}(u, b), \dots, \Phi_{j,m}(u, b))^T, \quad j = 1, 2.$$

“ T ” denoting transpose. Rewritten (5.1) and (5.2) in matrix, then we have

Theorem 5.2 The integro-differential equation in matrix form for $\Phi_1(u, b)$ and $\Phi_2(u, b)$, for $0 \leq u \leq b$,

$$\begin{aligned} \Phi'_1(u, b) = & \mathbf{H}_1 \Phi_1(u, b) + \int_0^u \mathbf{G}_1(x) \Phi_1(u - x, b) dx \\ & + \int_u^{u+\frac{c}{\beta}} \mathbf{G}_1(x) \Phi_2(u - x, b) dx + \mathbf{A}_1(u) \end{aligned} \tag{5.11}$$

and, for $-c/\beta < u \leq 0$,

$$\begin{aligned} \Phi'_2(u, b) = & \mathbf{H}_2(u) \Phi_2(u, b) \\ & + \int_0^{u+\frac{c}{\beta}} \mathbf{G}_2(x) \Phi_2(u - x, b) dx + \mathbf{A}_2(u) \end{aligned} \tag{5.12}$$

where

$$\mathbf{H}_1 = \begin{pmatrix} \frac{\lambda_1 + \delta}{c} & & & \\ & \frac{\lambda_2 + \delta}{c} & & \\ & & \ddots & \\ & & & \frac{\lambda_m + \delta}{c} \end{pmatrix}$$

$$\mathbf{H}_2(u) = \begin{pmatrix} \frac{\lambda_1 + \delta}{\beta u + c} & & & \\ & \frac{\lambda_2 + \delta}{\beta u + c} & & \\ & & \ddots & \\ & & & \frac{\lambda_m + \delta}{\beta u + c} \end{pmatrix}$$

$$\mathbf{G}_1(x) = - \begin{pmatrix} \frac{\lambda_1}{c} & & & \\ & \ddots & & \\ & & \frac{\lambda_m}{c} & \\ & & & \end{pmatrix} \Lambda \begin{pmatrix} f_1(x) & & & \\ & \ddots & & \\ & & & f_m(x) \end{pmatrix}$$

$$\mathbf{G}_2(x) = - \begin{pmatrix} \frac{\lambda_1}{\beta u + c} & & & \\ & \ddots & & \\ & & \frac{\lambda_m}{\beta u + c} & \\ & & & \end{pmatrix} \times \Lambda \begin{pmatrix} f_1(x) & & & \\ & \ddots & & \\ & & & f_m(x) \end{pmatrix}$$

are all $m \times m$ matrices, and $A_1(u)$ and $A_2(u)$ defined by

$$A_1(u) = \int_{u+\frac{c}{\beta}}^{\infty} \omega(u, x-u) G_1(x) I dx$$

$$A_2(u) = \int_{u+\frac{c}{\beta}}^{\infty} \omega(u, x-u) G_2(x) I dx$$

are all m -dimensional vector, in which $I = (1, 1, \dots, 1)^T$ is an $m \times 1$ column vector. The continuity condition and derivative condition for $\Phi_1(u, b)$ and $\Phi_2(u, b)$ is

$$\Phi_1(0+, b) = \Phi_2(0-, b),$$

$$\Phi_1'(0+, b) = \Phi_2'(0-, b).$$

6. Acknowledgements

The authors are very grateful to the editor and two anonymous referees for their valuable comments and suggestions which led to the present improved version of the manuscript. This research was supported by Humanities and Social Sciences Project of the Ministry Education of China (No. 09YJC910004; No. 10YJC630092) and Shandong Provincial Natural Science Foundation of China (No. ZR2010GL013) and Research Program of Higher Education of Shandong Province (No. J10WF84) and Natural Science Foundation of Shandong Jiaotong University (No. Z201031).

REFERENCES

- [1] J. M. Reinhard, "On a Class of Semi-Markov Risk Models Obtained as Classical Risk Models in a Markovian Environment," *Astin Bulletin*, Vol. 14, 1984, pp. 23-43.
- [2] S. Asmussen, "Risk Theory in a Markovian Environment," *Scandinavian Actuarial Journal*, No. 2, 1989, pp. 69-100.
- [3] A. Ng and H. Yang, "On the Joint Distribution of Surplus Prior and Immediately after Ruin under a Markovian Regime Switching Model," *Stochastic Processes and Their Applications*, Vol. 116, No. 2, 2006, pp. 244-266. [doi:10.1016/j.spa.2005.09.008](https://doi.org/10.1016/j.spa.2005.09.008)
- [4] S. M. Li and Y. Lu, "Moments of the Dividend Payments and Related Problems in a Markov-Modulated Risk Model," *North American Actuarial Journal*, Vol. 11, No. 2, 2007, pp. 65-76.
- [5] Y. Lu and S. Li, "The Markovian Regime-Switching Risk Model with a Threshold Dividend Strategy," *Insurance: Mathematics and Economics*, Vol. 44, No. 2, 2009, pp. 296-303. [doi:10.1016/j.insmatheco.2008.04.004](https://doi.org/10.1016/j.insmatheco.2008.04.004)
- [6] J. Liu, J. C. Xu and H. C. Hu, "The Markov-Dependent Risk Model with a Threshold Dividend Strategy," *Wuhan University Journal of Natural Sciences*, Vol. 16, No. 3, 2011, pp. 193-198. [doi:10.1007/s11859-011-0736-9](https://doi.org/10.1007/s11859-011-0736-9)
- [7] J. Zhu and H. Yang, "Ruin Theory for a Markov Regime-Switching Model under a Threshold Dividend Strategy," *Insurance: Mathematics and Economics*, Vol. 42, No. 1, 2008, pp. 311-318. [doi:10.1016/j.insmatheco.2007.03.004](https://doi.org/10.1016/j.insmatheco.2007.03.004)
- [8] J. Q. Wei, H. L. Yang and R. M. Wang, "On the Markov-modulated Insurance Risk Model with Tax," *Blaetter der DGVFM*, Vol. 31, No. 1, 2010, pp. 65-78. [doi:10.1007/s11857-010-0104-4](https://doi.org/10.1007/s11857-010-0104-4)
- [9] H. Albrecher and O. Boxma, "On the Discounted Penalty Function in a Markov-Dependent Risk Model," *Insurance Mathematics and Economics*, Vol. 37, No. 2, 2005, pp. 650-672. [doi:10.1016/j.insmatheco.2005.06.007](https://doi.org/10.1016/j.insmatheco.2005.06.007)
- [10] J. Liu, J. C. Xu and Y. J. Hu, "On the Expected Discounted Penalty Function in a Markov-Dependent Risk Model with Constant Dividend Barrier," *Acta Mathematica Scientia*, Vol. 30B, No. 5, 2010, pp. 1481-1491.
- [11] J. Liu, J. C. Xu and H. C. Hu, "Dividend Payments with a Threshold Strategy in a Markov-Dependent Risk Model," *Wuhan University Journal of Natural Sciences*, Vol. 16, No. 1, 2011, pp. 11-15. [doi:10.1007/s11859-011-0703-5](https://doi.org/10.1007/s11859-011-0703-5)
- [12] M. Zhou and C. Zhang, "Absolute Ruin under Classical Risk Model," *Acta Mathematicae Applicatae Sinica*, Vol. 28, No. 4, 2005, pp. 57-80.
- [13] J. Cai, "On the Time Value of Absolute Ruin with Debit Interest," *Advances in Applied Probability*, Vol. 39, No. 2, 2007, pp. 343-359. [doi:10.1239/aap/1183667614](https://doi.org/10.1239/aap/1183667614)
- [14] H. U. Gerber and H. L. Yang, "Absolute Ruin Probabilities in a Jump Diffusion Risk Model with Investment," *North American Actuarial Journal*, Vol. 11, No. 3, 2007, pp. 159-169.
- [15] K. C. Yuen, M. Zhou and J. Y. Guo, "On a Risk Model with Debit Interest and Dividend Payments," *Statistics and Probability Letters*, Vol. 78, No. 15, 2008, pp. 2426-2432. [doi:10.1016/j.spl.2008.02.021](https://doi.org/10.1016/j.spl.2008.02.021)
- [16] H. L. Yuan and Y. J. Hu, "Absolute Ruin in the Compound Poisson Risk Model with Constant Dividend Barrier," *Statistics and Probability Letter*, Vol. 78, No. 14, 2008, pp. 2086-2094. [doi:10.1016/j.spl.2008.01.076](https://doi.org/10.1016/j.spl.2008.01.076)
- [17] C. W. Wang and C. C. Yin, "Dividend Payments in the Classical Risk Model under Absolute Ruin with Debit Interest," *Applied Stochastic Models in Business and Industry*, Vol. 25, No. 3, 2009, pp. 247-262. [doi:10.1002/asmb.722](https://doi.org/10.1002/asmb.722)
- [18] R. X. Ming, W. Y. Wang and L. Q. Xiao, "On the Time Value of Absolute Ruin with Tax," *Insurance: Mathematics and Economics*, Vol. 46, No. 1, 2010, pp. 67-84. [doi:10.1016/j.insmatheco.2009.09.004](https://doi.org/10.1016/j.insmatheco.2009.09.004)
- [19] C. W. Wang, C. C. Yin and E. Q. Li, "On the Classical Risk Model with Credit and Debit Interests under Absolute Ruin," *Statistics and Probability Letters*, Vol. 80, No. 15, 2010, pp. 427-436. [doi:10.1016/j.spl.2009.11.020](https://doi.org/10.1016/j.spl.2009.11.020)
- [20] Z. M. Zhang, H. L. Yang and H. Yang, "On the Absolute Ruin in a Map Risk Model with Debit Interest," *Advances in Applied Probability*, Vol. 43, No. 1, 2011, pp. 77-96.

[doi:10.1239/aap/1300198513](https://doi.org/10.1239/aap/1300198513)

- [21] W. G. Yu and Y. J. Huang, "Absolute Ruin for a Risk Model with Credit and Debit Interest under a Threshold

Dividend Strategy," *Far East Journal of Applied Mathematics*, Vol. 57, No. 2, 2011, pp. 125-137.