

# Photosynthetic Light Utilization Efficiency, Water Relations and Leaf Growth of C<sub>3</sub> and CAM Tropical Orchids under Natural Conditions

Shawn Tay<sup>1</sup>, Jie He<sup>1\*</sup>, Tim Wing Yam<sup>2</sup>

<sup>1</sup>Natural Sciences and Science Education Academic Group, National Institute of Education, Nanyang Technological University, Singapore

<sup>2</sup>Singapore Botanic Gardens, National Parks Board, Singapore

Email: \*jie.he@nie.edu.sg

Received 21 September 2015; accepted 20 November 2015; published 25 November 2015

Copyright © 2015 by authors and Scientific Research Publishing Inc.

This work is licensed under the Creative Commons Attribution International License (CC BY).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

---

## Abstract

Native orchid species of Singapore in their natural conditions experience stress from high irradiance, high temperatures and periods of extended low rainfall, which impact orchid plant physiology and lead to reduced growth and productivity. In this study, it was found that there was a reduction in photochemical efficiency of photosystem II (PSII) in 6 native orchid species under high light (HL) and *Bulbophyllum membranaceum* under low light (LL). There was chronic photoinhibition in these 6 orchid species over a period of 3 months after transplanting onto the tree trunks without watering and fertilization, especially in *Coelogynes mayeriana* and *Bulbophyllum membranaceum* under both HL and LL. This chronic photoinhibition caused by sustained period of water deficit in their natural conditions was later reversed by natural re-watering conditions from higher rainfall. These results indicate that water deficit has a greater impact on photosynthetic light utilization efficiency than excess light. The present study also showed that after natural re-watering, relative water content (RWC) of leaves and pseudobulbs generally increased. During the natural re-watering, total leaf area also gradually increased and reached maximum expansion after 7 weeks under both HL and LL, with some exceptions due to leaf abscission or decline in total leaf area, possibly a strategy for water conservation.

## Keywords

Orchid, CAM, C<sub>3</sub>, Photosynthetic Light Utilization Efficiency, Water Relations and Leaf Growth

---

\*Corresponding author.

## 1. Introduction

Singapore has a tropical rainforest climate with no distinctive seasons, temperatures ranging from 22°C to 35°C, high relative humidity averages ranging from 60% to 100%, and annual rainfall about 2300 mm, with November to January being the wettest months. Considerable efforts have been made in the past decade to re-introduce native orchid species back into Singapore as part of conservation [1]-[3]. Many of these re-introduced species are epiphytes adapted to relatively dry habitats, native to low-elevation forests, and are dependent upon their microclimate. Crucial to these epiphytic orchids is the requirement for water economy where high evapotranspiration combined with scarce water supply challenges plant water balance [4]. Epiphytic orchids grow well under optimal environmental conditions and propagate naturally. However, when they experience stress from high irradiance, high temperatures and periods of extended low rainfall, these factors significantly impact their physiology and lead to reduced growth and productivity.

High irradiance and high temperature cause a reduction in photosynthetic quantum yield, loss in leaf chlorophyll, and a greater susceptibility to photoinhibition in orchids [5]-[7]. The reduction in photosynthesis can be attributed to dynamic photoinhibition, photooxidation, photobleaching of chlorophyll and chloroplast damage leading to deleterious effects of the leaf photosynthetic apparatus [7]-[9]. Furthermore, orchid leaves exposed to drastic increase in photon flux densities experienced a drop in CO<sub>2</sub> gas exchange and leaf chlorosis [10].

Epiphytic orchids experiencing stress from water deficit show significant decrease in relative water content (RWC) in leaves and pseudobulbs, reduction in photosynthetic light utilization, and greater susceptibility to photoinhibition caused by high irradiance [11]. In other plants, periodic drought stress also leads to a closure of stomata [12], through the reduction of shoot water potential directly or reduced turgor [13]. As a result, CO<sub>2</sub> depletes in the intercellular spaces and at the chloroplast level, limiting photosynthesis. Plants grown under low irradiance may also experience a reduction in the photosynthetic rate, resulting in a decline in growth rate [5] [14]. Therefore it is necessary to strike a good balance in maximising photon energy utilization of leaves to increase photosynthetic activity whilst minimising photoinhibition [6].

Orchids are also able to tolerate and acclimatise to stressful conditions, dynamically achieved by the physiological and morphological adaptations. In some orchid species, carbon fixation through Crassulacean acid metabolism [15] provides a means of water conservation during times of water deficit, while preventing loss of carbon [16]-[19]. Thick-leaved orchids, such as *Bulbophyllum vaginatum*, *Dendrobium leonis* and *Phalaenopsis cornu-cervi*, have features that are characteristic of plants that fix carbon primarily through Crassulacean acid metabolism [15], while thin-leaved orchids such as *Coelogynes rochussenii*, *Coelogynes mayeriana* and *Bulbophyllum membranaceum* are plants that fix carbon primarily through C<sub>3</sub> pathway [20]-[22].

There is little information about the photosynthetic light utilization efficiency and water relations of tropical orchid species in their natural environments, especially with regards to their responses to varying environmental conditions and the stress that comes with it. The information would be useful for improving the approach towards conservation of orchid species in their natural environments. Therefore, to complement the conservation efforts in Singapore, this study aims to better understand the photosynthetic light utilization efficiency, water relations and leaf growth of six selected C<sub>3</sub> and CAM native tropical orchids under natural conditions. The results benefit sustainable conservation of these orchid species under natural conditions.

## 2. Materials and Methods

### 2.1. Plant Cultivation under Natural Conditions

Mature plants of *Coelogynes rochussenii*, *Coelogynes mayeriana*, *Bulbophyllum vaginatum*, *Bulbophyllum membranaceum*, *Dendrobium leonis* and *Phalaenopsis cornu-cervi* planted by the National Parks Board (NParks) Singapore in National Institute of Education (NIE), Singapore, were used in this study. These epiphytic orchid plants were planted on the trunks of rain trees, *Albizia saman* (syn. *Samanea saman*), under natural high light (HL) and low light (LL) conditions on 12 January 2015. LL was achieved by natural shading provided by the trees and positioning the plants out of direct sunlight. These orchids were neither watered nor fertilized after planting, and depended upon rainfall as the only source of water. The maximal PPFD ranged from 90 to 200  $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$  for LL and from 400 to 1300  $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$  for HL, with an ambient day temperature of about 30°C to 35°C during the photoperiod. 6 plants of each species for each light condition were planted on the trees.

## 2.2. Measurement of Photosynthetic Photon Flux Density (PPFD)

PPFD was measured using a photosynthetically available radiation quantum sensor and reading unit (Skye Instruments Ltd, Llandrindod, UK). The stabilized PPFD, within a range of 0 to 1999  $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ , was measured from 6 different positions above the leaves for HL and LL respectively, and an average was calculated from the 6 values measured.

## 2.3. Measurement of Chl Fluorescence $F_v/F_m$ Ratio

Diurnal changes of  $F_v/F_m$  ratios were taken every two hours from 0800 to 1600 h with the Plant Efficiency Analyser, PEA (Hansatech Instruments Ltd, England) on sunny and cloudy days, while weekly  $F_v/F_m$  ratios were taken at the time corresponding to the maximal midday PPFD. 6 samples were taken from each species under each light condition. The leaves were pre-darkened with clips for 15 min prior to measurements. Dark adapted leaves were placed under the light pipe to measure  $F_o$ , initial chlorophyll (Chl) fluorescence during the very early onset of illumination with excitation light.  $F_m$ , maximum Chl fluorescence was determined by 0.8 seconds of saturated pulse ( $>6000 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ ). The variable fluorescence yield  $F_v$ , was determined by  $F_m - F_o$ . The efficiency of excitation energy captured by open PSII reaction centres in dark adapted leaves was estimated by the fluorescence  $F_v/F_m$  ratio.

## 2.4. Measurement of Chl Reading by Hand Held Chlorophyll Meter (SPAD-502)

Leaf Chl content was read by using a hand-held chlorophyll meter (SPAD-502, Minolta, Japan), which measures the absorbances of the leaf at wavelengths 645 nm and 663 nm. Using these two absorbances, the meter calculates a numerical SPAD value which is proportional to the amount of Chl present in the leaf, and expresses in terms of SPAD units. 6 samples were taken from each species under each light condition.

## 2.5. Measurement of Midday RWC

Square cuts (1 cm by 1 cm) were made out of the leaves and 5 mm thick slices were made out of the pseudobulbs. These sample were first weighed with an analytical balance to determine fresh weight (FW), followed by immersion in water in the dark for 3 hours (for pseudobulb of *C. mayeriana* and leaf of *B. membranaceum*) or 24 hours (for the rest of the samples) prior to measurement of their saturated weight (SW). The samples were then dried in the oven at 80°C for at least 72 hours to obtain their dry weight. RWC was expressed as  $\text{RWC} = (\text{FW} - \text{DW})/(\text{SW} - \text{DW}) \times 100\%$ . 10 samples each for leaves and for pseudobulbs were obtained from each species.

## 2.6. Measurement of Total Leaf Area

Young, budding leaves from 6 plants of each orchid species, cultivated under the respective light conditions (HL or LL), were selected and tracked over 16 weeks. The individual leaf area was measured every week by capturing the image of selected leaves against a measured scale and then using Image J software to calculate the leaf area from the image according the method outlined by Schneider *et al.* [23]. The total leaf area was calculated as a sum of the individual leaf areas. Total leaf area is one of the parameters for leaf growth.


## 2.7. Statistical Analysis

One-way ANOVA was used to test for significant differences among different growth irradiances, using Tukey's multiple comparison tests to discriminate the means (IBM SPSS Statistics for Macintosh, Version 22.0, 2013).

## 3. Results

### 3.1. Photosynthetic Light Utilization Efficiency

Diurnal PPFD was much higher for the plants grown under HL than LL on a sunny day at 1000 h, 1200 h and 1400 h (Figure 1(a) and Figure 1(b)) and the diurnal  $F_v/F_m$  ratio at these timings were also correspondingly lower in HL than LL (Figure 1(c) and Figure 1(d)), with the biggest difference in *B. vaginatum* and *B. membranaceum*. This suggests a reduction in the efficiency of open PSII reaction centres of the orchids under exposure




**Figure 1.** Diurnal changes of PPFD and  $F_v/F_m$  ratio on sunny days for *C. rochussenii* ( $\blacktriangle$ ), *C. mayeriana* ( $\blacksquare$ ), *B. vaginatum* ( $\diamond$ ), *B. membranaceum* ( $\square$ ), *D. leonis* ( $\bullet$ ) and *P. cornu-cervi* ( $\blacklozenge$ ) grown under HL and LL.


to HL (PPFDs of 400 to 1300  $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ ) towards midday, particularly more severe in *B. vaginatum* and *B. membranaceum*, among the 6 orchid species examined. A reduction in the efficiency of open PSII reaction centres was also shown in *B. membranaceum* under exposure to LL (with a maximal PPFD of 200  $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ ) during midday.  $F_v/F_m$  ratio at 1600 h of all plants grown under HL except *C. mayeriana*, and *B. membranaceum* grown under LL did not recover to the same levels as at 0800 h, suggesting that these plants were suffering from chronic photoinhibition.

However, there was evidence of recovery from this chronic photoinhibition as shown in the difference in  $F_v/F_m$  ratios at midday, between 1 and 3 months of growing under HL and LL. Significant increase in the midday  $F_v/F_m$  ratios from 1 to 3 months of cultivation were observed in all species under both HL (Figure 2(a)) and LL (Figure 2(b)) except for *C. mayeriana* under LL (One-way ANOVA,  $p < 0.05$ ,  $n = 6$ ). Between species, *B. vaginatum* and *B. membranaceum* after 1 month under HL show significantly lower  $F_v/F_m$  ratios as compared with the other species, but recovered to levels comparable to *D. leonis* and *P. cornu-cervi* after 3 months under HL (Tukey's multiple comparison test,  $p < 0.05$ ,  $n = 6$ ). No significant difference in  $F_v/F_m$  ratio was found between species after 1 month and 3 months under LL. Further evidence of recovery from this chronic photoinhibition was provided in the data collected over 16 weeks for the changes to midday maximal PPFD and  $F_v/F_m$  ratio. Over the 16 weeks of cultivation under HL, midday maximal PPFD exceeded 800  $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$  only in weeks 2 to 7 and week 13 (Figure 3(a)) while little fluctuation in midday PPFD ( $< 200 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ ) was seen in LL (Figure 3(b)). Based on rainfall data of the micro-climate of the area where the orchids were cultivated, weeks 2 to 7 coincided with a period of low rainfall (Figure 3(a)), while the drop in midday maximal PPFD ( $< 800 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ ) from week 8 onwards, with the exception of week 13, coincided with an increase in weekly total rainfall. Corresponding with the fluctuations in midday maximal PPFD under HL and the weekly total rainfall, the  $F_v/F_m$  ratio decreased from weeks 2 to 8 for all 6 orchid species, under both HL (Figure 3(c) and Figure 3(e)) and LL (Figure 3(d) and Figure 3(f)), and started increasing back to original levels or higher thereafter. The decrease in  $F_v/F_m$  ratio was greater in HL than LL, for *B. vaginatum*, *B. membranaceum* and *P. cornu-cervi*, *C. rochussenii*, *C. mayeriana* and *D. leonis*. Therefore, the period of higher weekly total rainfall from week 8 onwards provided re-watering conditions, and this corresponded with plants showing recovery from chronic photoinhibition, where the  $F_v/F_m$  ratio that was decreasing below 0.700 in the first 8 weeks started to increase back to levels above 0.700 after week 8.

In contrast to the changes in  $F_v/F_m$  ratio over the 16 weeks of cultivation, difference between Chl content under HL and LL were lesser, with the exception of *B. membranaceum* which showed a higher Chl content under



**Figure 2.**  $F_v/F_m$  ratio at midday maximal PPFD after 1 month (open bars) and 3 months (solid bars) of cultivation under HL and LL (*Cr*: *C. rochussenii*, *Cm*: *C. mayeriana*, *Bv*: *B. vaginatum*, *Bm*: *B. membranaceum*, *Dl*: *D. leonis*, *Pc*: *P. cornu-cervi*). Significant difference observed between the  $F_v/F_m$  ratio after 1 and 3 months, under HL and LL, in all species except for *Cm* under LL (One-way ANOVA,  $p < 0.05$ ,  $n = 6$ ). Different letters above bars indicate significant difference in  $F_v/F_m$  ratio between species after 1 month ((a), (b)) and 3 months ((c), (d), (e)) under HL (Tukey's multiple comparison,  $p < 0.05$ ,  $n = 6$ ). No significant difference found between species after 1 month and 3 months under LL.



**Figure 3.** Changes of midday PPFD and  $F_v/F_m$  ratios of different orchid species over 16 weeks under HL and LL. Rainfall data adapted with permission from Meteorological Services Singapore (Historical Daily Records, 2015).

LL (Figure 4(d)) than HL (Figure 4(c)). Chl content fluctuated little between 20 to 40 SPAD units for *C. rochussenii*, *C. mayeriana* and *P. cornu-cervi* (Figure 4(a) and Figure 4(b)), whereas Chl content showed greater fluctuations between 30 to 70 SPAD units for *B. vaginatum* and *D. leonis* (Figure 4(c) and Figure 4(d)).

### 3.2. Water Relations

To further understand the significance of the natural re-watering conditions provided by the higher weekly total rainfall from week 8 onwards, the water status of the plants was examined. RWC of leaves (Figure 5(a) and Figure 5(b)) and pseudobulbs (Figure 5(c) and Figure 5(d)) increased significantly in all 6 orchid species, between 1 to 4 months of cultivation under HL and LL, with the exception of the pseudobulbs of *B. vaginatum* and *B. membranaceum* under HL (One-way ANOVA,  $p < 0.05$ ,  $n = 10$ ). Between species, leaves of *B. vaginatum*, *B. membranaceum*, *D. leonis* and *P. cornu-cervi* after 1 month under HL and LL had significantly lower RWC compared to *C. rochussenii* and *C. mayeriana* (Tukey's multiple comparison test,  $p < 0.05$ ,  $n = 10$ ). However, after 4 months, RWC in the leaves of *P. cornu-cervi* under HL and LL and *B. membranaceum* under LL increased back to levels comparable with *C. rochussenii* and *C. mayeriana*. After 1 and 4 months under HL and LL, pseudobulbs of *C. rochussenii*, *B. vaginatum* and *B. membranaceum* remained significantly lower than that of *C. mayeriana*.

### 3.3. Leaf Growth

The difference in leaf growth under HL and LL was also examined through the changes in total leaf area for all 6 species. Total leaf area generally increased for the first 7 weeks for all 6 orchid species, with the exception of *D. leonis* cultivated under HL (Figure 6(e)), where there was a sharp drop in total leaf area during weeks 5 to 7. The sharp drop was due to a loss of the selected leaves, and this could be further attributed to the sustained low rainfall period coupled with HL from weeks 2 to 9. Total leaf area continued to be sustained at around the same level from week 7 onwards for the rest of the plants, with the exception of *C. rochussenii* (Figure 6(a)) and *B. membranaceum* (Figure 6(d)) cultivated under LL, and *C. mayeriana* (Figure 6(b)) and *P. cornu-cervi* (Figure 6(f)) cultivated under HL. For the plants with total leaf area remaining unchanged from week 7 onwards, it could be attributed to the leaves reaching their maximum expanded area.






Figure 4. Changes of Chl reading of different orchid species over 16 weeks under HL and LL.



**Figure 5.** RWC of leaves and pseudobulbs after 1 month (open bars) and 4 months (solid bars) of cultivation under HL and LL (*Cr*: *C. rochussenii*, *Cm*: *C. mayeriana*, *Bv*: *B. vaginatum*, *Bm*: *B. membranaceum*, *Dl*: *D. leonis*, *Pc*: *P. cornu-cervi*). Significant difference was observed between 1 and 4 months under HL and LL, in all species except for *Bv* pseudobulb under HL (One-way ANOVA,  $p < 0.05$ ,  $n = 10$ ). Different letters above bars indicate significant difference in RWC between species after 1 and 4 months under HL ((a), (b), (c)) and LL ((d), (e), (f)) (Tukey’s multiple comparison,  $p < 0.05$ ,  $n = 10$ ).



**Figure 6.** Total leaf area of different orchid species over 16 weeks under HL (◆) and LL (■).

## 4. Discussion

Some 70% of the native orchid species of Singapore are mostly vascular epiphytes [1]. Vascular epiphytes are dependent upon their micro-climate in photosynthetic light utilization and water economy [4]. In a review by Ort [24], it was reported that light intensities often exceed photosynthetic capacity in plants, while stomatal conductance limits CO<sub>2</sub> uptake into leaves, rendering even moderate irradiances in excess of photosynthetic capacity. Plants grown under high irradiance may lead to photoinhibition of the leaves, and this will result in a reduction in growth and productivity. Both C<sub>3</sub> and CAM orchids have been reported to show stress from exposure to high irradiances under natural conditions, such as decrease in photosynthetic light utilization, photoinhibition, photooxidative damage, and leaf chlorosis [5] [10] [11] [25]-[29]. Our present study reports a reduction in the efficiency of open PSII reaction centres of all the 6 orchid species under exposure to HL and C<sub>3</sub> orchid *B. membranaceum* under exposure to LL, towards midday (Figure 1), indicating photoinhibition. This photoinhibition is particularly more severe in *B. vaginatum* and *B. membranaceum* grown under HL. All plants under HL were also not able to fully recover from the photoinhibition over the first month (Figure 2(a), open bars), indicating chronic photoinhibition.

However, photoinhibition of PSII *in vivo* provides photoprotection and is an acclimation strategy for plants under stress from excessive irradiances [30]. Orchids have protective strategies to overcome the stress from high irradiances and to help balance maximising photon energy utilization while minimising photoinhibition [6]. Previous studies reported higher tolerance to high PPFD and faster recovery from photoinhibition in C<sub>3</sub> *Oncidium* orchid as compared to CAM *Phalaenopsis* orchid [29], greater photoprotection in leaves of C<sub>3</sub> orchid *C. rochussenii* under HL compared to LL [28], and recovery from photoinhibition in C<sub>3</sub> orchid *Gramatophyllum speciosum* in HL [27]. The results of our study seem to contradict these cases because the diurnal F<sub>v</sub>/F<sub>m</sub> ratios decline sharply towards 1200 h in all the orchids exposed to HL (Figure 1(c)) as compared to LL (Figure 1(d)). All 6 species also showed chronic photoinhibition after 1 month under HL, but recovered after 3 months (Figure 2). However, in all previous studies, water deficit was not severe and this could be the reason why the C<sub>3</sub> orchids could better utilize photosynthetic light energy as compared to the CAM orchids.

Ort [24] argued that at high irradiances, water status take physiological precedence over maximizing photosynthesis. It has been reported that in thin-leaved, C<sub>3</sub> epiphytic orchid *Catasetum viridiflavum*, and other epiphytes, re-watered plants show recovery from photoinhibition despite high PPFD, indicating that water deficit has a greater impact on photoinhibition than high light [31] [32]. Furthermore, strong photoinhibition and reduced photosynthetic capacity was observed in epiphytic CAM orchid hybrid, *Cattleya forbesii* Lindl. × *Laelia tenebrosa* Rolfe., following sustained water deficit for 42 days [11]. It has also been reported that under water deficit stress, *C. rochussenii* under HL had a reduced photosynthetic light utilization than plants under LL, and water deficit stress took precedence over light stress in reducing the Chl reading of *C. rochussenii*. [28]. Drought stress coupled with high light occurs when water supply is insufficient to sustain plant growth [27] [33]. Our present study showed that a sustained period of water deficit caused a decline in photosynthetic light utilization in all 6 orchid species, followed by a period of natural re-watering conditions that enabled recovery from this chronic photoinhibition (Figure 3). The changes in photosynthetic light utilization were less likely to be attributed to changes in Chl content since there was little difference in Chl content between HL and LL (Figure 4), with the exception of *B. membranaceum* which showed a higher Chl content under LL than HL. Therefore, the results suggest that water deficit has a greater impact on photosynthetic light utilization efficiency than HL, and that *B. vaginatum* and *B. membranaceum* thrive better in LL, well-watered conditions.

Water deficit that drives closure of stomata result in reduction in CO<sub>2</sub> uptake and subsequent down-regulation of photosynthetic efficiency [34], which is alleviated in epiphytic orchids by carbon fixation through CAM, while achieving water economy [4] [35] [36]. Epiphytic orchids have also been reported to lose water from the leaf slowly, yet maintain high values of RWC [37]. However, another study reported a reduction in RWC and osmotic potential during a period of water shortage in some tropical epiphytes [38] [39]. Moreover, studies had shown that pseudobulbs helped to reduce the loss of leaf water content during periods of water deficit stress by storing water in the pseudobulbs and then supplying this water to the leaves, thereby providing orchids with the ability to survive [11] [40]-[43]. A previous study had also reported that *C. rochussenii* grown under LL had a significantly higher leaf RWC than those under HL [28]. Our present study showed that after a period of natural re-watering, RWC of leaves and pseudobulbs increased significantly in all 6 orchid species, with the exception of the pseudobulbs of *B. vaginatum* and *B. membranaceum* under HL and *C. mayeriana* under LL (Figure 5).

Large pseudobulbs in  $C_3$  orchids *C. rochussenii* and *C. mayeriana* could have stored water and helped maintain leaf water content so as to alleviate photoinhibition. The small size and thin leaves of *B. membranaceum* could have disadvantaged it in its ability to store water in the pseudobulbs to overcome periods of water deficit and for the leaves to overcome the stress from high irradiance, which explains the severity of photoinhibition. On the other hand, in both *D. leonis* and *P. cornu-cervi*, having fleshy leaves and the ability to fix carbon through CAM could be significant strategies in response to photoinhibition. Therefore, both of these species are more adaptable to both HL and LL conditions, and could have a wider range of tolerable irradiance.

Water deficits lead to reduction in leaf area, leaf abscission, and dieback of individual plant parts. However, reduction in leaf area and leaf abscission strongly reduced water loss, and epiphytic orchids that encounter shortages of water may stop growing when water is in short supply, only to resume growth when water supply is restored [31] [44]. Our study showed a general increase in total leaf area for the 6 species for the first 7 weeks under HL and LL and maintained onwards as they had reached maximum leaf expansion. However, the drop in total leaf area in some species, such as *D. leonis*, was due to leaf abscission and decline in total leaf area. This could be a strategy for water conservation through reducing the total number of leaves, and therefore number of stomata, as well as reducing the surface area, leading to lesser water loss by evapotranspiration.

## 5. Conclusion

The results suggest that water deficit has a greater impact on photosynthetic light utilization efficiency than excess light. Over a period of water deficit, 6 orchid species experienced a decrease in  $F_v/F_m$  ratio and RWC of leaves and pseudobulbs, which could both be reversed by natural re-watering conditions from higher rainfall. Over 16 weeks, total leaf area increased and leaves reached maximum expansion after 7 weeks under both HL and LL, with some exceptions due to leaf abscission or decline in total leaf area, possibly a strategy for water conservation. Future work would be required to fully understand the adaptation strategies in response to stress from HL and water deficit that may occur in the natural environment. This will aid in improving the approach of re-introduction of native orchid species in Singapore.

## Acknowledgements

This project was carried out in National Institute of Education (NIE), Nanyang Technological University (NTU), Singapore. We would like to thank NIE, NTU for providing the space for orchid cultivation and also the facilities required for this study, as well as the Singapore Botanic Gardens and the National Parks Board for providing the plant materials. We would like to also thank Miss Choong Tsui Wei for her valuable inputs and assistance in optimizing the methods of this study and analyzing the results.

## References

- [1] Yam, T.W. (2013) Native Orchids of Singapore: Diversity, Identification and Conservation. National Parks Board, Singapore.
- [2] Yam, T.W., *et al.* (2011) Conservation and Reintroduction of Native Orchids of Singapore—The Next Phase. *European Journal of Environmental Sciences*, **1**, 38-47.
- [3] Yam, T.W. and Thame, A. (2005) Conservation and Reintroduction of the Native Orchids of Singapore. *Selbyana*, **26**, 75-80.
- [4] Benzing, D. (1998) Vulnerabilities of Tropical Forests to Climate Change: The Significance of Resident Epiphytes. *Climatic Change*, **39**, 519-540. <http://dx.doi.org/10.1023/A:1005312307709>
- [5] He, J., Khoo, G.H. and Hew, C.S. (1998) Susceptibility of CAM Dendrobium Leaves and Flowers to High Light and High Temperature under Natural Tropical Conditions. *Environmental and Experimental Botany*, **40**, 255-264. [http://dx.doi.org/10.1016/S0098-8472\(98\)00042-2](http://dx.doi.org/10.1016/S0098-8472(98)00042-2)
- [6] He, J. and Teo, L.C.D. (2007) Susceptibility of Green Leaves and Green Flower Petals of CAM Orchid Dendrobium cv. Burana Jade to High Irradiance under Natural Tropical Conditions. *Photosynthetica*, **45**, 214-221. <http://dx.doi.org/10.1007/s11099-007-0035-z>
- [7] Lichtenhaler, H.K., Buschmann, C., Döll, M., Fietz, H.-J., Bach, T., Kozel, U., *et al.* (1981) Photosynthetic Activity, Chloroplast Ultrastructure, and Leaf Characteristics of High-Light and Low-Light Plants and of Sun and Shade Leaves. *Photosynthesis Research*, **2**, 115-141. <http://dx.doi.org/10.1007/BF00028752>
- [8] Chow, W.S. (1994) Photoprotection and Photoinhibitory Damage. In: Bittar, E.E. and Barber, J. Eds., *Advances in*

*Molecular and Cell Biology*, Elsevier, London, 151-196. [http://dx.doi.org/10.1016/s1569-2558\(08\)60397-5](http://dx.doi.org/10.1016/s1569-2558(08)60397-5)

- [9] Osmond, C.B. (1994) What Is Photoinhibition? Some Insights from Comparisons of Shade and Sun Plants. Bios Scientific Publication, Oxford.
- [10] Johnson, S.R. (1993) Photosynthesis and Aspects of Phenology of the Rapidly Dispersing Orchid *Oeceoclades maculata*. *Lindleyana*, **8**, 69-72.
- [11] Stancato, G.C., Mazzafera, P. and Buckeridge, M.S. (2001) Effect of a Drought Period on the Mobilisation of Non-Structural Carbohydrates, Photosynthetic Efficiency and Water Status in an Epiphytic Orchid. *Plant Physiology and Biochemistry*, **39**, 1009-1016. [http://dx.doi.org/10.1016/s0981-9428\(01\)01321-3](http://dx.doi.org/10.1016/s0981-9428(01)01321-3)
- [12] Pinheiro, C. and Chaves, M.M. (2011) Photosynthesis and Drought: Can We Make Metabolic Connections from Available Data? *Journal of Experimental Botany*, **62**, 869-882.
- [13] Turner, N.C. and Jones, M.M. (1980) Turgor Maintenance by Osmotic Adjustment: A Review and Evaluation. In: Turner, N.C. and Kramer, P.J., Eds., *Adaptation of Plants to Water and High Temperature Stress*, Willey & Sons, New York, 87-103.
- [14] Lüttge, U., Ball, E., Fetene, M. and Medina, E. (1991) Flexibility of Crassulacean Acid Metabolism in *Kalanchoe Pinnata* (Lam.) Pers. I. Response to Irradiance and Supply of Nitrogen and Water. *Journal of Plant Physiology*, **137**, 259-267. [http://dx.doi.org/10.1016/S0176-1617\(11\)80129-X](http://dx.doi.org/10.1016/S0176-1617(11)80129-X)
- [15] Cameron, K.M., Chase, M.W., Whitten, W.M., Kores, P.J., Jarrell, D.C., Albert, V.A., *et al.* (1999) A Phylogenetic Analysis of the Orchidaceae: Evidence from rbcL Nucleotide Sequences. *American Journal of Botany*, **86**, 208-224. <http://dx.doi.org/10.2307/2656938>
- [16] Adams III, W.W. and Osmond, C.B. (1988) Internal CO<sub>2</sub> Supply during Photosynthesis of Sun and Shade Grown CAM Plants in Relation to Photo Inhibition. *Plant Physiology*, **86**, 117-123. <http://dx.doi.org/10.1104/pp.86.1.117>
- [17] Benzing, D.H. (1989) The Evolution of Epiphytism. In: Lüttge, U., Ed., *Vascular Plants as Epiphytes: Evolution and Ecophysiology*, Springer-Verlag, Berlin, 15-41. [http://dx.doi.org/10.1007/978-3-642-74465-5\\_2](http://dx.doi.org/10.1007/978-3-642-74465-5_2)
- [18] Silvera, K., Santiago, L.S., Cushman, J.C. and Winter, K. (2009) Crassulacean Acid Metabolism and Epiphytism Linked to Adaptive Radiations in the Orchidaceae. *Plant Physiology*, **149**, 1838-1847. <http://dx.doi.org/10.1104/pp.108.132555>
- [19] Silvera, K., Santiago, L.S., Cushman, J.C. and Winter, K. (2010) The Incidence of Crassulacean Acid Metabolism in Orchidaceae Derived from Carbon Isotope Ratios: A Checklist of the Flora of Panama and Costa Rica. *Botanical Journal of the Linnean Society*, **163**, 194-222. <http://dx.doi.org/10.1111/j.1095-8339.2010.01058.x>
- [20] Arditti, J. and Woolhuse, W.H. (1980) Aspects of the Physiology of Orchids. In: Woolhouse, H., Ed., *Advances in Botanical Research*, Academic Press, London, 421-655. [http://dx.doi.org/10.1016/s0065-2296\(08\)60091-9](http://dx.doi.org/10.1016/s0065-2296(08)60091-9)
- [21] Hew, C.S. and Yong, J.W.H. (2004) *The Physiology of Tropical Orchids in Relation to the Industry*. World Scientific Publishing Co. Pte. Ltd., Singapore.
- [22] Neales, T.F. and Hew, C.S. (1975) Two Types of Carbon Fixation in Tropical Orchids. *Planta*, **123**, 303-306. <http://dx.doi.org/10.1007/BF00390710>
- [23] Schneider, C.A., Rasband, W.S. and Eliceiri, K.W. (2012) NIH Image to Image J: 25 Years of Image Analysis. *Nature Methods*, **9**, 671-675. <http://dx.doi.org/10.1038/nmeth.2089>
- [24] Ort, D.R. (2001) When There Is Too Much Light. *Plant Physiology*, **125**, 29-32. <http://dx.doi.org/10.1104/pp.125.1.29>
- [25] Ding, T.H., Ong, H.T. and Yong, H.C. (1980) Factors Affecting Flower Development and Production of Golden Shower (*Oncidium goldiana*). *Proceedings of the 3rd ASEAN Orchid Congress*, Kuala Lumpur, 22-26 August, 65-78.
- [26] He, J., Norhafis, H. and Qin, L. (2013) Responses of Green Leaves and Green Pseudobulbs of CAM Orchid *Cattleya laeliocattleya* Aloha Case to Drought Stress. *Journal of Botany*, **2013**, 1-9.
- [27] He, J., Yong, T.Z. and Yam, T.W. (2014) Orchid Conservation in Singapore under Natural Conditions: Responses of *Grammatophyllum speciosum* to Growth Irradiances. *Plant Science International*, **1**, 11-23. <http://dx.doi.org/10.12735/psi.v1n1p11>
- [28] Tan, S.H.G. (2012) Photosynthetic Characteristics of Green Leaves and Green Pseudobulbs of *Coelogyne rochussenii* Orchid and Their Responses to Different Growth Irradiance and Drought Stress. National Institute of Education, Nanyang Technological University, Singapore.
- [29] Wang, S.Z. (2007) Effects of Growth Irradiance and Virus Infection on the Physiology of C<sub>3</sub> *Oncidium* and CAM *Phalaenopsis* Orchids under Natural Tropical Conditions. National Institute of Education, Nanyang Technological University, Singapore.
- [30] Anderson, J.M., Park, Y.I. and Chow, W.S. (1997) Photoinactivation and Photoprotection of Photosystem II in Nature. *Physiologia Plantarum*, **100**, 214-223. <http://dx.doi.org/10.1111/j.1399-3054.1997.tb04777.x>

- [31] Zotz, G. and Tyree, M. (1996) Water Stress in the Epiphytic Orchid, *Dimerandra emarginata* (G. Meyer) Hoehne. *Oecologia*, **107**, 151-159. <http://dx.doi.org/10.1007/BF00327898>
- [32] Zotz, G. and Winter, K. (1994) Annual Carbon Balance and Nitrogen-Use Efficiency in Tropical C3 and CAM Epiphytes. *New Phytologist*, **126**, 481-492. <http://dx.doi.org/10.1111/j.1469-8137.1994.tb04245.x>
- [33] Ashraf, M. (2010) Inducing Drought Tolerance in Plants: Recent Advances. *Biotechnology Advances*, **28**, 169-183. <http://dx.doi.org/10.1016/j.biotechadv.2009.11.005>
- [34] Cornic, G. and Fresneau, C. (2002) Photosynthetic Carbon Reduction and Carbon Oxidation Cycles Are the Main Electron Sinks for Photosystem II Activity during a Mild Drought. *Annals of Botany*, **89**, 887-894. <http://dx.doi.org/10.1093/aob/mcf064>
- [35] Haag-Kerwer, A., Franco, A.C. and Lüttge, U. (1992) The Effect of Temperature and Light on Gas Exchange and Acid Accumulation in the C3-CAM Plant *Clusia minor* L. *Journal of Experimental Botany*, **43**, 345-352. <http://dx.doi.org/10.1093/aob/mcf064>
- [36] Rodrigues, M.A., et al. (2013) Spatial Patterns of Photosynthesis in Thin- and Thick-Leaved Epiphytic Orchids: Unravelling C3-CAM Plasticity in an Organ-Compartmented Way. *Annals of Botany*, **112**, 17-29. <http://dx.doi.org/10.1093/aob/mct090>
- [37] Goh, C.J. and Kluge, M. (1989) Gas Exchange and Water Relations in Epiphytic Orchids. In: Lüttge, U., Ed., *Vascular Plants as Epiphytes, Evolution and Ecophysiology*, Springer-Verlag, Heidelberg, 139-166. <http://dx.doi.org/10.1007/bf00327898>
- [38] Sinclair, R. (1983) Water Relations of Tropical Epiphytes: I. Relationships between Stomatal Resistance, Relative Water Content and the Components of Water Potential. *Journal of Experimental Botany*, **34**, 1652-1663. <http://dx.doi.org/10.1093/jxb/34.12.1652>
- [39] Sinclair, R. (1983) Water Relations of Tropical Epiphytes: II. Performance during Droughting. *Journal of Experimental Botany*, **34**, 1664-1675. <http://dx.doi.org/10.1093/jxb/34.12.1664>
- [40] Chiang, Y.-L. and Chen, Y.-R. (1968) Observations on *Pleione formosana* Hayata. *Taiwania*, **14**, 271-301.
- [41] Ertelt, J.B. (1992) Horticultural Aspects of Growing and Displaying a Wide Variety of Epiphytes. *Selbyana*, **13**, 95-98.
- [42] Stern, W.L. and Morris, M.W. (1992) Vegetative Anatomy of Stanhopea (Orchidaceae) with Special Reference to Pseudobulb Water-Storage Cells. *Lindleyana*, **7**, 34-53.
- [43] Zheng, X.N., et al. (1992) Response of Cymbidium Sinense to Drought Stress. *The Journal of Horticultural Science & Biotechnology*, **67**, 295-300.
- [44] Yam, T.W. (2011) Conservation and Reintroduction of the Native Epiphytic Orchids of Singapore—A Physiological and Developmental Biology Perspective. In: *Proceedings of the International Conference on Biological Science*, Faculty of Biology Universitas Gadjah Mada, Yogyakarta, 2-9.