

A Recursive Approach to the Kauffman Bracket

Abdul Rauf Nizami, Mobeen Munir, Umer Saleem, Ansa Ramzan

Division of Science and Technology, University of Education, Lahore, Pakistan

Email: arnizami@ue.edu.pk, mobeenmunir@gmail.com, umerlinks@hotmail.com, ansaramzan@yahoo.com

Received 26 July 2014; revised 28 August 2014; accepted 10 September 2014

Copyright © 2014 by authors and Scientific Research Publishing Inc.

This work is licensed under the Creative Commons Attribution International License (CC BY).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

Abstract

We introduce a simple recursive relation and give an explicit formula of the Kauffman bracket of two-strand braid link $\widehat{x_1^n}$. Then, we give general formulas of the bracket of the sequence of links of three-strand braids $\alpha(n) = x_1 x_2 x_1 x_2 \cdots$. Finally, we give an interesting result that the Kauffman bracket of the three-strand braid link $\widehat{x_1^m x_2^n}$ is actually the product of the brackets of the two-strand braid links $\widehat{x_1^m}$ and $\widehat{x_1^n}$. Moreover, a recursive relation for $\langle \widehat{x_1^a x_2^b x_1^c x_2^d} \rangle$ is also given.

Keywords

Recursive Relation, Kauffman Bracket, Braid Link

1. Introduction

The Kauffman bracket polynomial was introduced by L. H. Kauffman in 1987 [1] in concern with link invariants. The bracket polynomial soon became popular due to its connections with the Jones polynomial, dichromatic polynomial, and the Potts model. While the HOMFLY polynomial and the bracket polynomial are distinct with different topological properties, there is a very beautiful relationship between them due to F. Jaeger [2], and it is also observed in a special case by Reshetikhin [3].


The Kauffman bracket (polynomial) is actually not a link invariant because it is not invariant under the first Reidemeister move. However, it has many applications and it can be extended to a popular link invariant, the Jones polynomial. In the present work we shall confine ourselves to the Kauffman bracket to avoid this work from unnecessary length and to leave it for applications.

This paper is organized as follows: In Section 2 we shall give the basic ideas about knots, braids, and the Kauffman bracket. In Section 3 we shall present the main results.


2. Basic Notions

2.1. Links

A *link* is a disjoint union of circles embedded in \mathbb{R}^3 . A one-component link is called a *knot*. Links are usually studied via projecting them on a plan; a projection with extra information of *overcrossing* and *undercrossing* is called the *link diagram*.



Two links are *isotopic* if and only if one of them can be transformed to the other by a diffeomorphism of the ambient space onto itself. A fundamental result by Reidemeister [4] about the isotopic link diagrams is: *Two unoriented links L_1 and L_2 are equivalent if and only if a diagram of L_1 can be transformed into a diagram of L_2 by a finite sequence of ambient isotopies of the plane and the local (Reidemeister) moves of the following three types:*




The set of all links that are equivalent to a link L is called a *class* of L . By a link L we shall always mean the class of L .

The main question of knot theory is *Which two links are equivalent and which are not?* To address this question one needs a *knot invariant*, a function that gives one value on all links that belong to a single class and gives different values (but not always) on knots that belong to different classes. The present work is basically concerned with this question.

2.2. Braids


Braids were first studied by Emil Artin in 1925 [5] [6], which now play an important role in knot theory, see [7]-[9] for detail.

An n -strand *braid* is a set of n non intersecting smooth paths connecting n points on a horizontal plane to n points exactly below them on another horizontal plane in an arbitrary order. The smooth paths are called *strands* of the braid.




The *product* ab of two n -strand braids is defined by putting the braid b above the braid a and then gluing their common end points.

A braid with only one crossing is called *elementary* braid. The i th elementary braid x_i on n strands is:



A useful property of elementary braids is that every braid can be written as a product of elementary braids. For instance, the above 2-strand braid is $x_i^{-3} = (x_i^{-1})(x_i^{-1})(x_i^{-1})$.

The closure of a braid b is the link \hat{b} obtained by connecting the lower ends of b with the corresponding upper ends.




An important result by Alexander [10] connecting knots and braids is: Each link can be represented as the closure of a braid. This result motivated knot theorists to study braids to solve problems of knot theory.

Remark 2.1 In the last section, all the concerned links will be closures of products of elementary braids.

2.3. The Kauffman Bracket

Before the definition it is better to understand the two types of splitting of a crossing, the A-type and the B-type splittings:



In the following, the symbols \bigcirc and \sqcup represent respectively the unknot and the disconnected sum.


Definition 2.2 The Kauffman bracket is the function $\langle \cdot \rangle: \text{Links} \rightarrow \mathbb{Z}[a, a^{-1}]$ defined by the axioms:

$$\langle L \rangle = a \langle L_A \rangle + a^{-1} \langle L_B \rangle$$

$$\langle L \sqcup \bigcirc \rangle = (-a^2 - a^{-2}) \langle L \rangle$$

$$\langle \bigcirc \rangle = 1.$$

Here L , L_A , and L_B are three links which are isotopic everywhere except at one crossing where they look as in the figure:




Proposition 2.3 The Kauffman polynomial is invariant under second and third Reidemeister moves but not under the first Reidemeister move [11].

3. Main Results

In this section we shall introduce a recursive relation for the Kauffman bracket, shall give an explicit formula of $\langle \widehat{x_1^n} \rangle$, and shall express $\langle \widehat{x_1^m x_2^n} \rangle$ as the product of $\langle \widehat{x_1^m} \rangle$ and $\langle \widehat{x_1^n} \rangle$.

First of all we give the Kauffman bracket of the k -twist unknot U_k :



Lemma 3.1 *The Kauffman bracket of the k -twist unknot is*

$$\langle U_k \rangle = (-1)^k a^{3k}.$$

Proof. We prove it by induction on k :

The case $k=0$ holds by definition as U_0 is the unknot without any crossings. Now, with the assumption that the result holds for an arbitrary k , we have

$$\begin{aligned} \langle U_{k+1} \rangle &= \langle \text{diagram} \rangle = a \langle \text{diagram} \rangle + a^{-1} \langle \text{diagram} \rangle \\ &= a(-a^2 - a^{-2}) \langle U_k \rangle + a^{-1} \langle U_k \rangle \\ &= -a^3 \langle U_k \rangle - a^{-3} [(-1)^k a^{3k}] \\ &= (-1)^{k+1} a^{3(k+1)}. \quad \square \end{aligned}$$

Theorem 3.2 (A recursive relation) *The following relation holds for any $n \geq 2$:*

$$\langle \widehat{x_1^n} \rangle = (-1)^{n-1} a^{3n-2} + a^{-1} \langle \widehat{x_1^{n-1}} \rangle. \tag{3.1}$$

Proof. We prove it using directly the definition and Lemma 3.1:

$$\begin{aligned} \langle \widehat{x_1^n} \rangle &= \langle \text{diagram} \rangle = a \langle \text{diagram} \rangle + a^{-1} \langle \text{diagram} \rangle \\ &= a \langle U_{n-1} \rangle + a^{-1} \langle \widehat{x_1^{n-1}} \rangle \\ &= a [(-1)^{n-1} a^{3(n-1)}] + a^{-1} \langle \widehat{x_1^{n-1}} \rangle \\ &= (-1)^{n-1} a^{3n-2} + a^{-1} \langle \widehat{x_1^{n-1}} \rangle. \quad \square \end{aligned}$$

From this recursive relation, we get the explicit formula for the 2-strand braid link $\widehat{x_1^n}$:

Proposition 3.3 *The Kauffman bracket of the link $\widehat{x_1^n}, n \geq 2$, is*

$$\langle \widehat{x_1^n} \rangle = -a^{-n-2} + \sum_{k=1}^{n-1} (-1)^{n+k-2} a^{(3n+2)-4k}.$$

Proof. We prove it by induction on n .

For $n = 2$, we have

$$\begin{aligned}
 \langle \widehat{x_1^2} \rangle &= \langle \text{Diagram 1} \rangle = a \langle \text{Diagram 2} \rangle + a^{-1} \langle \text{Diagram 3} \rangle \\
 &= a \left[a \langle \text{Diagram 4} \rangle + a^{-1} \langle \text{Diagram 5} \rangle \right] + a^{-1} \left[a \langle \text{Diagram 6} \rangle + a^{-1} \langle \text{Diagram 7} \rangle \right] \\
 &= a[a(-a^2 - a^{-2}) + a^{-1}(1)] + a^{-1}[a(1) + a^{-1}(-a^2 - a^{-2})] \\
 &= -a^4 - a^{-4},
 \end{aligned}$$

which satisfies the recursive relation.

With the assumption that the relation holds for an arbitrary n , we, using Theorem 3.2, get

$$\begin{aligned}
 \langle \widehat{x_1^{n+1}} \rangle &= (-1)^{(n+1)-1} a^{3(n+1)-2} + a^{-1} \langle \widehat{x_1^n} \rangle \\
 &= (-1)^n a^{3(n+1)-2} + a^{-1} \left[(-1)^{n-1} a^{3n-2} + (-1)^{n-2} a^{3n-6} \right. \\
 &\quad \left. + (-1)^{n-3} a^{3n-10} + \dots + (-1)^{2n-3} a^{-n+6} - a^{-n-2} \right] \\
 &= (-1)^n a^{3(n+1)-2} + \left[(-1)^{n-1} a^{3n-3} + (-1)^{n-2} a^{3n-7} \right. \\
 &\quad \left. + (-1)^{n-3} a^{3n-11} + \dots + (-1)^{2n-3} a^{-n+5} - a^{-n-3} \right] \\
 &= (-1)^n a^{3(n+1)-2} + (-1)^{n-1} a^{3(n+1)-6} + (-1)^{n-2} a^{3(n+1)-10} \\
 &\quad + (-1)^{n-3} a^{3(n+1)-14} + \dots + (-1)^{2(n+1)-3} a^{-(n+1)+6} - a^{-(n+1)-2}.
 \end{aligned}$$

This completes the proof. \square

In the following we give the Kauffman bracket polynomial of the closure of the braid $\alpha(n) = x_1 x_2 x_1 x_2 \dots (n \text{ factors})$; this sequence contains the powers of the Garside element $\Delta = x_1 x_2 x_1 = x_2 x_1 x_2 : \alpha(3k) = \Delta^k$.

Proposition 3.4 *The Kauffman bracket of $\alpha(n) = x_1 x_2 x_1 \dots (n\text{-times})$ satisfies the recurrence relations:*

$$\begin{aligned}
 \langle \Delta^{2k} \rangle &= a^6 \langle \Delta^{2(k-1)} \rangle - a^{16-6k} - a^{8-6k} + a^{4-6k} + a^{-4-6k} \\
 \langle \Delta^{2k} x_1 \rangle &= a^6 \langle \Delta^{2(k-1)} x_1 \rangle - a^{15-6k} - a^{7-6k} + a^{3-6k} + a^{-5-6k} \\
 \langle \Delta^{2k} x_1 x_2 \rangle &= a^6 \langle \Delta^{2(k-1)} x_1 x_2 \rangle - a^{14-6k} - a^{6-6k} + a^{2-6k} + a^{-6-6k} \\
 \langle \Delta^{2k+1} \rangle &= a^6 \langle \Delta^{2k-1} \rangle - a^{13-6k} - a^{5-6k} + a^{1-6k} + a^{-7-6k} \\
 \langle \Delta^{2k+1} x_2 \rangle &= a^6 \langle \Delta^{2k-1} x_2 \rangle - a^{12-6k} - a^{4-6k} + a^{-6k} + a^{-8-6k} \\
 \langle \Delta^{2k+1} x_2 x_1 \rangle &= a^6 \langle \Delta^{2k-1} x_2 x_1 \rangle - a^{11-6k} - a^{3-6k} + a^{-1-6k} + a^{-9-6k}
 \end{aligned}$$

Proof. Simply, apply the definition for different values of k , and write recursively each next bracket in terms of the previous one. \square

Lemma 3.5 *The Kauffman brackets for $k = 0$ are:*

$$\begin{aligned}
 \langle \Delta^0 \rangle &= a^4 + 2 + a^{-4} \\
 \langle \Delta^0 x_1 \rangle &= a^{-1} + a^{-5} \\
 \langle \Delta^0 x_1 x_2 \rangle &= a^{-6}
 \end{aligned}$$

$$\begin{aligned} \langle \Delta \rangle &= a + a^{-7} \\ \langle \Delta x_2 \rangle &= -a^4 + 1 + a^{-8} \\ \langle \Delta x_2 x_1 \rangle &= a^7 - a^3 + a^{-1} + a^{-9} \end{aligned}$$

Proof. The proofs of first three cases are given (proofs of remaining cases are similar):

$$\begin{aligned} \langle \Delta^0 \rangle &= \langle \bigcirc \rangle = (-a^2 - a^{-2})^2 \langle \bigcirc \rangle = a^4 + 2 + a^{-4} \\ \langle \Delta^0 x_1 \rangle &= \langle \bigcirc \rangle = (-a^2 - a^{-2}) \langle \bigcirc \rangle = (-a^2 - a^{-2}) [a \langle \bigcirc \rangle + a^{-1} \langle \bigcirc \rangle] \\ &= (-a^2 - a^{-2}) [a + a^{-1}(-a^2 - a^{-2})] = a^{-1} + a^{-5} \\ \langle \Delta^0 x_1 x_2 \rangle &= \langle \bigcirc \rangle = a \langle \bigcirc \rangle + a^{-1} \langle \bigcirc \rangle \\ &= a \langle x_1 \rangle + a^{-1}(-a^2 - a^{-2}) \langle x_1 \rangle = a^{-6} \quad \square \end{aligned}$$

Theorem 3.6 For any $k \geq 0$ the Kauffman bracket of $\alpha(n) = x_1 x_2 x_1 \dots (n\text{-times})$ is given by:


$$\begin{aligned} \langle \Delta^{2k} \rangle &= 2a^{6k} + a^{-6k+4} + a^{-6k-4} \\ \langle \Delta^{2k} x_1 \rangle &= -a^{6k+3} + a^{6k-1} + a^{-6k+3} + a^{-6k-5} \\ \langle \Delta^{2k} x_1 x_2 \rangle &= -a^{6k+2} + a^{-6k+2} + a^{-6k-6} \\ \langle \Delta^{2k+1} \rangle &= a^{-6k+1} + a^{-6k-7} \\ \langle \Delta^{2k+1} x_2 \rangle &= -a^{6k+4} + a^{-6k} + a^{-6k-8} \\ \langle \Delta^{2k+1} x_2 x_1 \rangle &= a^{6k+7} - a^{6k+3} + a^{-6k-1} + a^{-6k-9}. \end{aligned}$$

Proof. We prove it by induction on k . The case $k = 0$ is covered by Lemma 3.5, and the inductive step can be checked with Proposition 3.4.

For instance,

$$\begin{aligned} \langle \Delta^{2k+2} \rangle &= a^6 \langle \Delta^{2k} \rangle - a^{10-6k} - a^{2-6k} + a^{-2-6k} + a^{-10-6k} \\ &= a^6 [2a^{6k} + a^{4-6k} + a^{-4-6k}] - a^{10-6k} - a^{2-6k} + a^{-2-6k} + a^{-10-6k} \\ &= 2a^{6+6k} + a^{10-6k} + a^{2-6k} - a^{10-6k} - a^{2-6k} + a^{-10-6k} + a^{-2-6k} \\ &= 2a^{6(k+1)} + a^{4-6(k+1)} + a^{-4-6(k+1)}. \quad \square \end{aligned}$$

In connected sum $\widehat{x_1^n} \# U_k$ of the braid link $\widehat{x_1^n}$ with the trivial knot U_k has the diagram:



Lemma 3.7

$$\langle \widehat{x_1^n} \# U_k \rangle = (-1)^k a^{3k} \langle \widehat{x_1^n} \rangle.$$

Proof. We prove it by induction on k :

For $k = 1$, we have

$$\begin{aligned} \langle \widehat{x_1^n} \# U_1 \rangle &= \langle \text{diagram} \rangle = a \langle \text{diagram} \rangle + a^{-1} \langle \text{diagram} \rangle \\ &= a(-a^2 - a^{-2}) \langle \widehat{x_1^n} \rangle + a^{-1} \langle \widehat{x_1^n} \rangle \\ &= -a^3 \langle \widehat{x_1^n} \rangle. \end{aligned}$$

Now, with the assumption that the result holds for an arbitrary k , we have

$$\begin{aligned} \langle \widehat{x_1^n} \# U_{k+1} \rangle &= \langle \text{diagram} \rangle \\ &= a \langle \text{diagram} \rangle + a^{-1} \langle \text{diagram} \rangle \\ &= a(-a^2 - a^{-2}) \langle \widehat{x_1^n} \# U_k \rangle + a^{-1} \langle \widehat{x_1^n} \# U_k \rangle \\ &= -a^3 \langle \widehat{x_1^n} \# U_k \rangle \\ &= -a^3 [(-1)^k a^{3k} \langle \widehat{x_1^n} \rangle] \\ &= (-1)^{k+1} a^{3(k+1)} \langle \widehat{x_1^n} \rangle, \end{aligned}$$

as required. \square

The following result confirms that the Kauffman bracket of $\langle \widehat{x_1^m x_2^n} \rangle$ is actually the product $\langle \widehat{x_1^m} \rangle \langle \widehat{x_1^n} \rangle$.

Theorem 3.8 For any $m, n \geq 2$,

$$\langle \widehat{x_1^m x_2^n} \rangle = \langle \widehat{x_1^m} \rangle \langle \widehat{x_1^n} \rangle.$$

Proof. We prove it by induction on n :

When $n = 2$,

$$\begin{aligned} \langle \widehat{x_1^m x_2^2} \rangle &= \langle \text{diagram} \rangle = a \langle \text{diagram} \rangle + a^{-1} \langle \text{diagram} \rangle \\ &= a^2 \langle \text{diagram} \rangle + \langle \text{diagram} \rangle + \langle \text{diagram} \rangle + a^{-2} \langle \text{diagram} \rangle \\ &= a^2(-a^2 - a^{-2}) \langle \widehat{x_1^m} \rangle + \langle \widehat{x_1^m} \rangle + \langle \widehat{x_1^m} \rangle + a^{-2}(-a^2 - a^{-2}) \langle \widehat{x_1^m} \rangle \\ &= \langle \widehat{x_1^m} \rangle [-a^4 - 1 + 1 + 1 - 1 - a^{-4}] \\ &= \langle \widehat{x_1^m} \rangle [-a^4 - a^{-4}] = \langle \widehat{x_1^m} \rangle \langle \widehat{x_1^2} \rangle. \end{aligned} \tag{3.2}$$

Suppose the result holds for $n = k$, that is $\langle \widehat{x_1^m x_2^k} \rangle = \langle \widehat{x_1^m} \rangle \langle \widehat{x_1^k} \rangle$.

Now, using Lemma 3.7, we have

$$\begin{aligned}
 \langle \widehat{x_1^m x_2^{k+1}} \rangle &= a \langle \widehat{x_1^m \# U_k} \rangle + a^{-1} \langle \widehat{x_1^m x_2^k} \rangle \\
 &= a \left[(-1)^k a^{3k} \langle \widehat{x_1^m} \rangle \right] + a^{-1} \langle \widehat{x_1^m} \rangle \langle \widehat{x_1^k} \rangle \\
 &= \langle \widehat{x_1^m} \rangle \left[(-1)^k a^{3k+1} + a^{-1} \langle \widehat{x_1^k} \rangle \right] \\
 &= \langle \widehat{x_1^m} \rangle \left[(-1)^k a^{3k+1} + (-1)^{k-1} a^{3k-3} + (-1)^k a^{3k-7} \right. \\
 &\quad \left. + (-1)^{k+1} a^{3k-11} + \dots + (-1)^{2k-2} a^{-k+5} - a^{-k-3} \right] \\
 &= \langle \widehat{x_1^m} \rangle \left[(-1)^{k-2} a^{3(k+1)-2} + (-1)^{k-1} a^{3(k+1)-6} \right. \\
 &\quad \left. + (-1)^k a^{3(k+1)-10} + \dots + (-1)^{2k-2} a^{-(k+1)+6} - a^{-(k+1)-2} \right] \\
 &= \langle \widehat{x_1^m} \rangle \langle \widehat{x_1^{k+1}} \rangle.
 \end{aligned}$$

This completes the proof. \square

Corollary 3.9

$$\langle \widehat{x_1^m x_2^n} \rangle = \langle \widehat{x_1^n x_2^m} \rangle.$$

Proof. It is obvious: $\langle \widehat{x_1^m x_2^n} \rangle = \langle \widehat{x_1^m} \rangle \langle \widehat{x_1^n} \rangle = \langle \widehat{x_1^n} \rangle \langle \widehat{x_1^m} \rangle = \langle \widehat{x_1^n x_2^m} \rangle$.

Corollary 3.10


$$\deg \langle \widehat{x_1^m x_2^n} \rangle = 3(m+n) - 4$$

and

$$\text{span} \langle \widehat{x_1^m x_2^n} \rangle = 4(m+n).$$

Proof. The result follows immediately from Theorem 3.8 as $\deg \langle \widehat{x_1^n} \rangle = 3n - 2$ and $\text{span} \langle \widehat{x_1^n} \rangle = -n - 2$. \square

For the following, let us fix the notation L_{abc} for the link with the understanding that the link contains a , b , and c crossings of type x_1, x_2 , and x_1 , respectively, and that $L_{abc} \neq \widehat{x_1^a x_2^b x_1^c}$.



Proposition 3.11 The Kauffman bracket of the link L_{abc} is

$$\langle L_{abc} \rangle = \langle \widehat{x_1^a} \rangle \langle \widehat{x_1^b} \rangle \langle \widehat{x_1^c} \rangle + (a^{-b+2} + a^{-b-2}) \langle \widehat{x_1^a} \rangle \langle \widehat{x_1^c} \rangle + a^{-b} \langle \widehat{x_1^{a+c}} \rangle$$

Proof. We prove it by induction on b :
 For $b = 1$, we have

$$\begin{aligned} \langle L_{a1c} \rangle &= \langle \widehat{x_1^a} \rangle \langle \widehat{x_1^1} \rangle \langle \widehat{x_1^c} \rangle + (a + a^{-3}) \langle \widehat{x_1^a} \rangle \langle \widehat{x_1^c} \rangle + a^{-1} \langle \widehat{x_1^{a+c}} \rangle \\ &= a^{-3} \langle \widehat{x_1^a} \rangle \langle \widehat{x_1^c} \rangle + (a + a^{-3}) \langle \widehat{x_1^a} \rangle \langle \widehat{x_1^c} \rangle + a^{-1} \langle \widehat{x_1^{a+c}} \rangle \\ &= a \langle \widehat{x_1^a} \rangle \langle \widehat{x_1^c} \rangle + a^{-1} \langle \widehat{x_1^{a+c}} \rangle. \end{aligned}$$

Now, with the assumption that the result holds for an arbitrary k , we have $b = k + 1$

$$\begin{aligned} \langle L_{a(k+1)c} \rangle &= a \langle \text{Diagram 1} \rangle + a^{-1} \langle \text{Diagram 2} \rangle \\ &= a \langle \widehat{x_1^a} \# U_k \# \widehat{x_1^c} \rangle + a^{-1} \langle \widehat{x_1^a x_2^k x_1^c} \rangle \\ &= a \langle \widehat{x_1^a} \# U_k \rangle \langle \widehat{x_1^c} \rangle + a^{-1} \langle \widehat{x_1^a x_2^k x_1^c} \rangle \\ &= a(-1)^k a^{3k} \langle \widehat{x_1^a} \rangle \langle \widehat{x_1^c} \rangle + a^{-1} \left(\langle \widehat{x_1^a} \rangle \langle \widehat{x_1^k} \rangle \langle \widehat{x_1^c} \rangle + (a^{-k+2} + a^{-k-2}) \langle \widehat{x_1^a} \rangle \langle \widehat{x_1^c} \rangle + a^{-k} \langle \widehat{x_1^{a+c}} \rangle \right) \\ &= (-1)^k a^{3k+1} \langle \widehat{x_1^a} \rangle \langle \widehat{x_1^c} \rangle + a^{-1} \langle \widehat{x_1^a} \rangle \langle \widehat{x_1^k} \rangle \langle \widehat{x_1^c} \rangle + (a^{-k+1} + a^{-k-3}) \langle \widehat{x_1^a} \rangle \langle \widehat{x_1^c} \rangle + a^{-k-1} \langle \widehat{x_1^{a+c}} \rangle \\ &= \left((-1)^k a^{3k+1} + (-1)^{k+1} a^{3k-3} + \dots - a^{-k+5} - a^{-k-3} \right) \langle \widehat{x_1^a} \rangle \langle \widehat{x_1^c} \rangle \\ &\quad + (a^{-k-1+2} + a^{-k-1-2}) \langle \widehat{x_1^a} \rangle \langle \widehat{x_1^c} \rangle + a^{-k-1} \langle \widehat{x_1^{a+c}} \rangle \\ &= \langle \widehat{x_1^a} \rangle \langle \widehat{x_1^{k+1}} \rangle \langle \widehat{x_1^c} \rangle + (a^{-(k+1)+2} + a^{-(k+1)-2}) \langle \widehat{x_1^a} \rangle \langle \widehat{x_1^c} \rangle + a^{-(k+1)} \langle \widehat{x_1^{a+c}} \rangle \end{aligned}$$

as required. \square

Proposition 3.12 The Kauffman bracket of the link $\widehat{x_1^a x_2^b x_1^c x_2^d}$ is


$$\begin{aligned} \langle \widehat{x_1^a x_2^b x_1^c x_2^d} \rangle &= \left(\sum_{i=1}^d (-1)^{d+i} a^{3d-4i+2} \right) \langle \widehat{x_1^a} \rangle \langle \widehat{x_1^b} \rangle \langle \widehat{x_1^c} \rangle \\ &\quad + \left((-1)^{d+1} a^{-b+3d} + a^{-b-d} \right) \langle \widehat{x_1^a} \rangle \langle \widehat{x_1^c} \rangle \\ &\quad + \left(\sum_{i=1}^d (-1)^{d+i} a^{3d-4i+2-b} + a^{-d} \langle \widehat{x_1^b} \rangle \right) \langle \widehat{x_1^{a+c}} \rangle. \end{aligned}$$

Proof. We prove it by induction on d :

For $d = 1$, we have

$$\begin{aligned} \langle \widehat{x_1^a x_2^b x_1^c x_2^1} \rangle &= \left(\sum_{i=1}^1 (-1)^{d+i} a^{3d-4i+2} \right) \langle \widehat{x_1^a} \rangle \langle \widehat{x_1^b} \rangle \langle \widehat{x_1^c} \rangle + \left((-1)^{d+1} a^{-b+3} + a^{-b-1} \right) \langle \widehat{x_1^a} \rangle \langle \widehat{x_1^c} \rangle \\ &\quad + \left(\sum_{i=1}^1 (-1)^{1+i} a^{3-4i+2-b} + a^{-1} \langle \widehat{x_1^b} \rangle \right) \langle \widehat{x_1^{a+c}} \rangle \\ &= a \langle \widehat{x_1^a} \rangle \langle \widehat{x_1^b} \rangle \langle \widehat{x_1^c} \rangle + (a^{-b+3} + a^{-b-1}) \langle \widehat{x_1^a} \rangle \langle \widehat{x_1^c} \rangle \\ &\quad + (a^{-b+1} + a^{-1} \langle \widehat{x_1^b} \rangle) \langle \widehat{x_1^{a+c}} \rangle. \end{aligned}$$

Now, with the assumption that the result holds for $d = k$, we have



$$\begin{aligned}
\langle \widehat{x_1^a x_2^b x_1^c x_2^{k+1}} \rangle &= a \langle \text{Diagram 1} \rangle + a^{-1} \langle \text{Diagram 2} \rangle = a(L_{abc} \# U_k) + a^{-1} \langle \widehat{x_1^a x_2^b x_1^c x_2^k} \rangle \\
&= a \left[(-1)^k a^{3k} \left(\langle \widehat{x_1^a} \rangle \langle \widehat{x_1^b} \rangle \langle \widehat{x_1^c} \rangle + (a^{-b+2} + a^{-b-2}) \langle \widehat{x_1^a} \rangle \langle \widehat{x_1^c} \rangle + a^{-b} \langle \widehat{x_1^{a+c}} \rangle \right) \right] \\
&\quad + a^{-1} \left[\left(\sum_{i=1}^k (-1)^{k+i} a^{3k-4i+2} \right) \langle \widehat{x_1^a} \rangle \langle \widehat{x_1^b} \rangle \langle \widehat{x_1^c} \rangle + \left((-1)^{k+1} a^{-b+3k} + a^{-b-k} \right) \langle \widehat{x_1^a} \rangle \langle \widehat{x_1^c} \rangle \right] \\
&\quad + \left(\sum_{i=1}^k (-1)^{k+i} a^{3k-4i+2-b} + a^{-k} \langle \widehat{x_1^b} \rangle \right) \langle \widehat{x_1^{a+c}} \rangle \\
&= (-1)^k a^{3k+1} \langle \widehat{x_1^a} \rangle \langle \widehat{x_1^b} \rangle \langle \widehat{x_1^c} \rangle + \left((-1)^k a^{3k+3-b} + (-1)^k a^{3k-1-b} \right) \langle \widehat{x_1^a} \rangle \langle \widehat{x_1^c} \rangle \\
&\quad + (-1)^k a^{3k+1-b} + \sum_{i=1}^k (-1)^{k+i} a^{3k-4i+1} \langle \widehat{x_1^a} \rangle \langle \widehat{x_1^b} \rangle \langle \widehat{x_1^c} \rangle \\
&\quad + \left((-1)^{k+1} a^{3k-1-b} + a^{-k-1-b} \right) \langle \widehat{x_1^a} \rangle \langle \widehat{x_1^c} \rangle + \left(\sum_{i=1}^k (-1)^{k+i} a^{3k-4i+1-b} + a^{-k-1} \langle \widehat{x_1^b} \rangle \right) \langle \widehat{x_1^{a+c}} \rangle \\
&= \left(\sum_{i=1}^{k+1} (-1)^{(k+1)+i} a^{3(k+1)-4i+2} \right) \langle \widehat{x_1^a} \rangle \langle \widehat{x_1^b} \rangle \langle \widehat{x_1^c} \rangle + \left((-1)^{(k+1)+1} a^{-b+3(k+1)} + a^{-b-(k+1)} \right) \langle \widehat{x_1^a} \rangle \langle \widehat{x_1^c} \rangle \\
&\quad + \left(\sum_{i=1}^{k+1} (-1)^{(k+1)+i} a^{3(k+1)-4i+2-b} + a^{-(k+1)} \langle \widehat{x_1^b} \rangle \right) \langle \widehat{x_1^{a+c}} \rangle,
\end{aligned}$$

as was required. \square

References

- [1] Kauffman, L.H. (1987) State Models and the Jones Polynomial. *Topology*, **26**, 395-407. [http://dx.doi.org/10.1016/0040-9383\(87\)90009-7](http://dx.doi.org/10.1016/0040-9383(87)90009-7)
- [2] Jaeger, F. (1990) A Combinatorial Model for the Homy Polynomial. *European Journal of Combinatorics*, **11**, 549-555.
- [3] Reshetikhin, N.Y. (1988) Quantized Universal Enveloping Algebras, the Yang-Baxter Equation and Invariants of Links, I and II. LOMI Reprints E-4-87 and E-17-87, Steklov Institute, Leningrad, USSR.
- [4] Reidemeister, K. (1948) Knot Theory. Chelsea Publ and Co., New York.
- [5] Artin, E. (1925) Theorie der Zöpfe. *Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg*, **4**, 27-72. <http://dx.doi.org/10.1007/BF02950718>
- [6] Artin, E. (1947) Theory of Braids. *Annals of Mathematics*, **48**, 101-126. <http://dx.doi.org/10.2307/1969218>
- [7] Birman, J.S. (1974) Braids, Links, and Mapping Class Groups. Princeton University Press, Princeton.
- [8] Manturov, V.O. (2004) Knot Theory. Chapman and Hall/CRC, Boca Raton. <http://dx.doi.org/10.1201/9780203402849>
- [9] Murasugi, K. (1996) Knot Theory and Its Applications. Birkhäuser, Boston.
- [10] Alexander, J. (1923) Topological Invariants of Knots and Links. *Transactions of the American Mathematical Society*, **20**, 275-306.
- [11] Adams, C.C. (1994) The Knot Book. W H Freeman and Company, New York.