

# Relative effects of anaerobically-digested and conventional liquid swine manure, and N fertilizer on crop yield and greenhouse gas emissions

Reynald L. Lemke<sup>1\*</sup>, Sukhdev S. Malhi<sup>2</sup>, Fernando Selles<sup>3</sup>, Mark Stumborg<sup>1</sup>

<sup>1</sup>Agriculture and Agri-Food Canada, Saskatoon, Canada; \*Corresponding Author: [reynald.lemke@agr.gc.ca](mailto:reynald.lemke@agr.gc.ca)

<sup>2</sup>Agriculture and Agri-Food Canada, Melfort, Canada

<sup>3</sup>Agriculture and Agri-Food Canada, Brandon, Canada

Received 23 June 2012; revised 29 July 2012; accepted 8 August 2012

## ABSTRACT

Anaerobic digestion is a promising technology that could provide an option for managing animal waste with reduced greenhouse gas emissions. A three-year (2006-2008) field experiment was conducted at Star City, Saskatchewan, Canada, to compare the effects of land-applied anaerobically digested swine manure (ADSM), conventionally treated swine manure (CTSM) and N fertilizer on grain yield of barley, applied N use efficiency (ANUE,  $\text{kg}\cdot\text{grain}\cdot\text{kg}^{-1}$  of applied  $\text{N}\cdot\text{ha}^{-1}$ ), ammonia ( $\text{NH}_3$ ) volatilization and nitrous oxide ( $\text{N}_2\text{O}$ ) emissions. Treatments included spring and autumn applications of CTSM and ADSM at a 1× rate (10,000 and 7150  $\text{L}\cdot\text{ha}^{-1}$ , respectively) applied every year, a 3× rate (30,000 and 21,450  $\text{L}\cdot\text{ha}^{-1}$ , respectively) applied once at the beginning of the experiment, plus a treatment receiving commercial fertilizer (UAN at 60  $\text{kg}\cdot\text{N}\cdot\text{ha}^{-1}\cdot\text{yr}^{-1}$ ) and a zero-N control. There was a significant grain yield response of barley to applied N in all three years. The ANUE of ADSM or CTSM applied once at the 3× rate were lower than annual applications at the 1× rate (grain yield by 595  $\text{kg}\cdot\text{ha}^{-1}$  and NFUE by 6  $\text{kg}\cdot\text{grain}\cdot\text{kg}^{-1}$  of applied  $\text{N}\cdot\text{ha}^{-1}$ ). On average, agronomic performance of ADSM was similar to CTSM. The APNU of N fertilizer was greater than the 3× rate but lower than the 1× rate of ADSM or CTSM. Ammonia loss from ADSM was similar to CTSM, except for much higher loss of  $\text{NH}_3\text{-N}$  from CTSM at the 3× rate applied in the autumn (8100  $\text{g}\cdot\text{N}\cdot\text{ha}^{-1}$ ) compared to the other treatments (1100 - 2600  $\text{g}\cdot\text{N}\cdot\text{ha}^{-1}$ ). The percentage of applied N lost as  $\text{N}_2\text{O}$  gas was generally higher for treatments receiving CTSM (4.0%) compared to ADSM (1.4%). In conclusion, the findings suggest that ADSM is equal or slightly better than CTSM in

terms of agronomic performance, but has lower environmental impact.

**Keywords:** Ammonia Volatilization; Anaerobic Digestion; Barley Yield; Nitrogen Fertilizer Use Efficiency; Nitrous Oxide; Swine Manure

## 1. INTRODUCTION

In 2009, over 28 million hogs were marketed by Canadian farmers, with nearly one-half of that industry located in the prairie region. Approximately 90% of intensive livestock operations in the prairie region store manure in liquid form in a holding tank or lagoon [1] until it can be land-applied. Considerable amounts of methane ( $\text{CH}_4$ ) are emitted to the atmosphere during storage [2] and, while land application of liquid swine manure provides an effective source of nutrients for crop production [3], high ammonia ( $\text{NH}_3$ ) volatilization rates can occur following application [4-6]. In addition, soil-emitted nitrous oxide can also be stimulated [7]. Economically feasible, environmentally friendly, and socially acceptable management of animal wastes from intensive livestock operations is a key element for the future viability of this industry.

Anaerobic digestion of liquid swine manure is a promising technology that could provide a cost effective option for reducing greenhouse gas (GHG) emissions from liquid swine manure management by avoiding lagoon storage and the associated  $\text{CH}_4$  emissions, and utilizing the biogas produced during digestion to displace fossil-fuels. Biogas digestion has the potential to directly or indirectly influence  $\text{NH}_3$  volatilization and  $\text{N}_2\text{O}$  emissions. Anaerobic digestion decreases slurry viscosity and volatile fatty acid content, while increasing slurry pH and inorganic C content [8,9]. Reduced viscosity could decrease  $\text{NH}_3$  volatilization from pig slurry [4,10], whereas increased pH and carbonate content could stimulate  $\text{NH}_3$

volatilization [11,12]. Rubaek *et al.* [13] measured similar  $\text{NH}_3$  volatilization following application of undigested and anaerobically digested pig slurry on grasslands in the United Kingdom. Similarly, Chantigny *et al.* [14] found no differences between undigested and anaerobically digested pig slurry from a site in Quebec, Canada. Conversely in another study also in Quebec, Chantigny *et al.* [7] measured lower  $\text{NH}_3$  volatilization losses from digested compared to raw swine manure.

Land-applied liquid animal manures generally promotes soil-emitted nitrous oxide [7]. Anaerobic digestion of manures results in more recalcitrant products which may reduce the rate of microbial degradation and oxygen consumption in the soil [15-17], leading to less anoxic microsites which favor denitrifying activity. However, the inorganic nitrogen (N) content of digested manures tends to be higher, which could favour higher nitrification rates, and coincident  $\text{N}_2\text{O}$  production, and higher nitrate production—which would increase denitrification potential. The reported effects of manure treatments on nitrous oxide emissions are variable. Some authors reported similar  $\text{N}_2\text{O}$  emissions from raw compared to anaerobically digested manure [18], others have reported decreased emissions after anaerobic digestion [13,19,20]. Conversely, Chantigny *et al.* [7] reported increased emissions from digested compared to raw liquid swine manure, when the manures were injected into the soil.

Many factors, including environmental, soil, and application technique could potentially interact with manure type to influence  $\text{NH}_3$  volatilization and  $\text{N}_2\text{O}$  emission following land-application of the material. To the authors' knowledge, there is very limited research information internationally [13] comparing the agronomic and environmental performance of land-applied raw versus anaerobically digested swine manure, and no published results for the Canadian prairie region. The objective of this study was to compare agronomic performance and gaseous N loss of land-applied anaerobically digested swine manure (ADSM) to conventionally treated (raw) swine manure (CTSM).

## 2. MATERIALS AND METHODS

A 3-year (2006-2008) field experiment was conducted at Star City (Typic Haplocryalf) Saskatchewan, Canada. Precipitation during the growing season (May, June, July and August) from 2006 to 2008, and long-term (30-year) average for the same period taken from the nearest Environment Canada Meteorological Station (AAFC Melfort), are presented in **Table 1**. Precipitation in the growing season was slightly below average in 2006, slightly above average in 2007 and much below average (especially in May during seeding) in 2008. Eleven treatments (**Table 2**) were arranged in a randomized complete block

**Table 1.** Monthly cumulative precipitation during 2006, 2007 and 2008 at Star City, Saskatchewan.

Year	Precipitation (mm)				
	May	June	July	August	Total
2006	63	73	39	46	221
2007	71	119	47	40	277
2008	6	32	117	22	177
30-year mean	46	66	76	57	245

**Table 2.** List of treatments and the corresponding total amount of N applied during a three-year field study at Star City, Saskatchewan.

Time of application	Product applied	Application rate	Total N applied (3-year cumulative)
Autumn	<sup>a</sup> ADSM-3x	21,450 L·ha <sup>-1</sup>	214
	ADSM-1x	7150 L·ha <sup>-1</sup>	205
	CTSM-3x	30,000 L·ha <sup>-1</sup>	403
	CTSM-1x	10,000 L·ha <sup>-1</sup>	360
	ADSM-3x	21,450 L·ha <sup>-1</sup>	257
Spring	ADSM-1x	7150 L·ha <sup>-1</sup>	255
	CTSM-3x	30,000 L·ha <sup>-1</sup>	343
	CTSM-1x	10,000 L·ha <sup>-1</sup>	326
	UAN	60 kg·N·ha <sup>-1</sup>	180
	Control	0	0

<sup>a</sup>ADSM = Anaerobically digested swine manure, CTSM = Conventionally treated swine manure, UAN = Urea ammonium nitrate (liquid).

design with four replicates. Liquid manures were applied by the Prairie Agricultural Machinery Institute (PAMI) using a customized applicator which injects the material to a 10 cm depth. All plots were seeded to barley (*Hordeum vulgare* L.) in each of the three years (AC Rosser in 2006 and 2007; Newdale in 2008). Seeding dates and rates, weed control and harvesting operations followed standard agronomic practice.

Conventionally treated swine manure was obtained from a commercial 1200 sow farrow-to-finish barn. Initial batches of the ADSM were obtained from a full-scale pilot mesophilic digester situated by the commercial barn. Later batches (autumn 2007 and spring 2008) were obtained from a small-scale pilot mesophilic digester operated by PAMI. Operating conditions for the small-scale digester were purposefully maintained to be comparable with the full scale version.

Previous research has indicated that application rates of CTSM, providing between 75 and 150 kg·N·ha<sup>-1</sup>, are most effective for agronomic performance in this region

[21]. Based on analysis of the CTSM to be applied in the first autumn (2005) of the study, an application rate of 10,000 L·ha<sup>-1</sup>, a typical rate used by producers in Saskatchewan, would provide about 100 kg·N·ha<sup>-1</sup>. Similarly, based on analysis of the ADSM supplied for application in the fall of 2005, an application rate of 7150 L·ha<sup>-1</sup> provided a comparable amount of N. Treatments receiving CTSM and ADSM at 3× this rate were also applied. The “1×” rate was applied in each of the three years while the “3×” rate was applied only once at the beginning of the study. While not recommended, the latter treatment is a common practice employed by producers in this region. Rates were held constant on a volume basis throughout the study. However, the N concentration contained in both the CTSM and ADSM varied considerably from application period to application period. The cumulative N applied over the life of the study is presented in **Table 2**. To account for the differences in the actual N applied, grain yields and NH<sub>3</sub> and N<sub>2</sub>O losses were normalized by expressing them as a ratio of N applied prior to statistical analysis. Applied N use efficiency (ANUE) of barley grain yield for the various treatments was calculated as: [(3-year total grain yield ha<sup>-1</sup> for treatment) – (3-year total grain yield ha<sup>-1</sup> for check)] ÷ (3-year total N applied to treatment).

Ammonia volatilization was measured using the “double-sponge open-chamber” technique [22], with measurements made on a set schedule for 2 - 3 weeks following application of the treatments. Briefly, a white polyvinyl chloride tube 20 cm long and 15 cm in diameter was inserted in the soil to a depth of 5 cm. A foam disk impregnated with an acid solution is inserted inside the chamber to absorb NH<sub>3</sub> evolved from the soil. A second disc closes the top of the chamber to allow for exchange of air between the chamber and the surroundings while scrubbing out atmospheric NH<sub>3</sub>. The discs were prepared by washing twice with distilled water, twice with 0.001 M H<sub>2</sub>SO<sub>4</sub> and twice with a glycerol-phosphoric acid solution. The lower disc was placed 5 cm above the soil surface, and the upper disc was placed 5 cm below the top of the cylinder. White plastic shields, supported at the corners by reinforcing bars, were placed 30 cm above the tops of the cylinders to protect the discs from rainfall but still allow air movement. Discs are exchanged at 1, 2, 4, 8 and 16 d after manure or fertilizer application, and rinsed in 0.5 M KCl. The concentration of ammonium in the extractant was determined with a Technicon Autoanalyzer [23]. Cumulative losses for each sampling period were calculated by interpolating between data points and integrating over time assuming a constant flux. Cumulative losses were normalized by subtracting the NH<sub>3</sub> lost from the check (no N applied) treatment and dividing that difference by the total N applied.

Nitrous oxide gas samples were collected using a

non-flow through non-steady state chamber method [24]. Sample collection protocols were similar to those described by Rochette *et al.* [25]. Briefly, plexi-glass frames (22 cm × 45.5 cm and 10 cm high) were permanently installed in the soil between crop rows but covering manure or fertilizer injection bands, and lids were sealed to the frames for the collection period. Gas samples were drawn from the chamber headspace at three equally spaced time intervals, over a 60-minute period, by fully filling disposable 20-mL polypropylene syringes and transferring to pre-evacuated 13 mL exetainer™ glass tubes for transport to the laboratory. The concentration of N<sub>2</sub>O in the sample containers was determined using a gas chromatograph equipped with a <sup>63</sup>Ni electron capture detector (ECD). The calculated minimum detectable difference for the system was <10 ppbv. Nitrous oxide flux rate was calculated as the first derivative of the second-order polynomial equation that best described the concentration versus time relationship, with adjustments for non-standard conditions of humidity, temperature and barometric pressure as described by Rochette and Hutchinson [26]. Time zero values were estimated using a method similar to that described by Anthony *et al.* [27]. A series of ambient air samples was collected at each sampling time. The mean of these samples was used as the time zero concentration. Gas sampling was done at least weekly, with increased frequency when expected emission activity was high (after snow melt and application of manure or fertilizer) and reduced frequency during the latter part of the season when soil-water contents were low. Seasonal estimates of N<sub>2</sub>O emissions were calculated by interpolating between data points and integrating over time assuming a constant flux [28]. The percentage of applied N lost as N<sub>2</sub>O-N was calculated by subtracting the N<sub>2</sub>O-N lost from the check (no N applied) treatment and dividing that difference by the total N applied.

The data on various parameters were subjected to analysis of variance (ANOVA) using procedures as outlined in SAS [29]. Significant ( $P \leq 0.05$ ) differences between treatments were determined using least significant difference (LSD<sub>0.05</sub>).

### 3. RESULTS AND DISCUSSION

Rainfall was somewhat lower than the long-term mean during the July-August period in 2006 and 2007 (**Table 1**), but above average precipitation during the early part of the season (May-June) carried the crop through with good grain yields in 2006 (**Table 3**), and modest grain yields in 2007. Extremely low rainfall was received in May and June, above average rainfall in July, followed by very dry conditions through August of 2008. This somewhat erratic rainfall pattern resulted in modest grain

**Table 3.** Barley grain yields from various treatments for three years at Star City, Saskatchewan.

Time	N source/rate	2006	2007	2008	Mean
		kg·ha <sup>-1</sup>			
Autumn	<sup>a</sup> ADSM-3x	6268	2213	3325	3935
	ADSM-1x	5609	2792	3497	3966
	CTSM-3x	5837	3256	3924	4339
	CTSM-1x	6375	4257	4699	5110
Spring	ADSM-3x	6250	2502	3258	4003
	ADSM-1x	6202	3050	4504	4585
	CTSM-3x	5946	2913	3653	4171
	CTSM-1x	6437	3228	4725	4797
	UAN	5387	2119	3725	3744
	Control	3487	1241	2629	2452
LSD <sub>0.05</sub>		322 <sup>***b</sup>	443 <sup>***</sup>	305 <sup>***</sup>	194 <sup>***</sup>

<sup>a</sup>ADSM = Anaerobically digested swine manure, CTSM = Conventionally treated swine manure, UAN = Urea ammonium nitrate (liquid). <sup>\*\*\*b</sup>Refers to significant at P < 0.001.

yields. Compared to the zero-N control, there was a significant increase in grain yield of barley from application of ADSM, CTSM and N fertilizer in all three years. Similarly, other researchers have also found swine manure very effective in increasing crop yields in the years of application [3,21]. In general, the 1× application rates of ADSM or CTSM had the highest ANUE, the 3× application rates the lowest, and UAN was intermediate (**Table 4**). Further, ADSM tended to have similar or slightly better ANUE compared to CTSM.

Ammonia volatilization losses were generally quite low. Cumulative losses over all sampling periods ranged from less than a kilogram to about 3 kg of N·ha<sup>-1</sup> (**Table 5**). The exception was the autumn applied CTSM-3x treatment which lost over 8 kg·N·ha<sup>-1</sup>. When these losses were compared on a relative basis, (g NH<sub>3</sub>-N·kg<sup>-1</sup> applied NH<sub>4</sub>-N), the autumn applied CTSM-3x treatment was significantly higher than all other treatments. In contrast, Rubaek *et al.* [13] did not find any difference in NH<sub>3</sub> volatilization loss from undigested versus anaerobically digested pig slurry on grassland in UK. This discrepancy between the two studies could be due to the differences in soil-climatic conditions and crop type.

Nitrous oxide emissions responded to the treatments in a relatively consistent fashion. Emissions were highest from the CTSM treatments, with particularly high losses in the first year of the study on the treatment receiving CTSM at the 3× rate (**Table 6**). When emissions were expressed as a percentage of applied N lost as N<sub>2</sub>O,

**Table 4.** Applied N use efficiency (ANUE) of barley grain yield for varying rates and sources of applied N at Star City, Saskatchewan.

Time	N source/rate	<sup>b</sup> ANUE
		kg grain·kg <sup>-1</sup> applied N·ha <sup>-1</sup>
Autumn	<sup>a</sup> ADSM-3x	21bc
	ADSM-1x	22b
	CTSM-3x	14e
	CTSM-1x	22b
Spring	ADSM-3x	18d
	ADSM-1x	<sup>c</sup> 25a
	CTSM-3x	15e
Spring	CTSM-1x	22b
	UAN	19cd

<sup>a</sup>ADSM = Anaerobically digested swine manure, CTSM = Conventionally treated swine manure, UAN = Urea ammonium nitrate (liquid); <sup>b</sup>[(3-yr total grain yield ha<sup>-1</sup> for treatment) - (3-year total grain yield ha<sup>-1</sup> for control)] ÷ (3-year total N applied ha<sup>-1</sup> to treatment); <sup>c</sup>The values are significantly different, when not followed by the same letter, based on LSD<sub>0.05</sub>.

**Table 5.** Estimated ammonia-N (NH<sub>3</sub>-N) loss over three sampling periods from various treatments at Star City, Saskatchewan.

Time	N source/rate	Net NH <sub>3</sub> -N loss	<sup>b</sup> NH <sub>3</sub> -N loss response
		g·N·ha <sup>-1</sup>	g NH <sub>3</sub> -N·kg <sup>-1</sup> applied N·ha <sup>-1</sup>
Autumn	<sup>a</sup> ADSM-3x	2600b	<sup>c</sup> 13ab
	ADSM-1x	1200b	6b
	CTSM-3x	8100a	24a
	CTSM-1x	3000b	10b
Spring	ADSM-3x	1100b	5b
	ADSM-1x	1700b	8b
	CTSM-3x	1700b	6b
	CTSM-1x	2500b	10ab
	UAN	800b	6b

<sup>a</sup>ADSM = Anaerobically digested swine manure, CTSM = Conventionally treated swine manure, UAN = Urea ammonium nitrate (liquid); <sup>b</sup>[(Cumulative NH<sub>3</sub>-N lost from treatment) - (Cumulative NH<sub>3</sub>-N lost from control)] ÷ [Cumulative NH<sub>4</sub>-N applied]; <sup>c</sup>The values are significantly different, when not followed by the same letter, based on LSD<sub>0.05</sub>.

losses were significantly higher from the treatments receiving CTSM at the 1× and 3× rate compared to treatments receiving ADSM at the 1× rate and the UAN treatment (**Table 7**). The treatment receiving ADSM at the 3× rate was intermediate and significantly different from CTSM at the 3× rate applied in the autumn. Lower N<sub>2</sub>O emissions from digested compared to raw swine manure was also reported by Chantigny *et al.* [7] and

**Table 6.** Estimated annual and three-year cumulative N<sub>2</sub>O-N loss from various treatments at Star City, Saskatchewan.

Time	N source/rate	2006	2007	2008	3-year total
		kg·N·ha <sup>-1</sup>			
Autumn	<sup>a</sup> ADSM-3x	2.9b	1.1b	3.4ab	7.4b
	ADSM-1x	1.3b	2.2b	2.0b	5.5b
	CTSM-3x	16.3a	1.8b	3.4ab	21.5a
	CTSM-1x	3.6b	5.9a	7.3a	16.8a
Spring	ADSM-1x	1.7b	2.1	2.2b	6.0b
	CTSM-1x	3.2b	5.4a	6.7a	15.3a
	UAN	1.1b	1.1b	2.3b	4.5b
	Control	0.8b	0.8b	1.6b	3.2b

<sup>a</sup>ADSM = Anaerobically digested swine manure, CTSM = Conventionally treated swine manure, UAN = Urea ammonium nitrate (liquid); <sup>c</sup>The values in each column separately are significantly different, when not followed by the same letter, based on LSD<sub>0.05</sub>.

**Table 7.** Percentage of applied N lost as N<sub>2</sub>O-N over three years at Star City, Saskatchewan.

Time	N source/rate	N <sub>2</sub> O-N loss as a percentage of applied N
		%
Autumn	ADSM-3x	2.0bc
	ADSM-1x	1.1c
	<sup>a</sup> CTSM-3x	4.5a
	CTSM-1x	3.8ab
Spring	ADSM-1x	1.1c
	CTSM-1x	3.7ab
	UAN	0.7c

<sup>a</sup>ADSM = Anaerobically digested swine manure, CTSM = Conventionally treated swine manure, UAN = Urea ammonium nitrate (liquid); <sup>c</sup>The values are significantly different, when not followed by the same letter, based on LSD<sub>0.05</sub>.

Vallejo *et al.* [20], and similar results have been reported for digested cattle manure [13,19,30]. Nyberg *et al.* [31] reported that some compounds present in anaerobically digested manure may have a depressive effect on soil ammonia oxidizers, thereby reducing the supply of substrate for N<sub>2</sub>O production through nitrification and denitrification. Vallejo *et al.* [20] argued that because most easily degradable C present in manure is decomposed during anaerobic digestion, the C remaining in the digested manure is more stable and, therefore, less likely to stimulate denitrification and N<sub>2</sub>O production as compared with the undigested manure.

#### 4. CONCLUSION

There was a significant grain yield response of barley

to applied N in all three years. The ANUE of barley for single applications of ADSM or CTSM at the 3× rate was lower than three annual applications at the 1× rate, while UAN was intermediate. The ANUE of ADSM and CTSM applied in autumn was equal to spring when applied at 1× rate and, in general, agronomic performance of ADSM was similar or better than CTSM. The APNU of N fertilizer was greater than the 3× rate but lower than the 1× rate of ADSM or CTSM. Ammonia losses for all treatments were low (<1 kg·N·yr<sup>-1</sup>) except for CTSM at the 3× rate applied in the autumn (>81 kg·N·yr<sup>-1</sup>). In general, NH<sub>3</sub> loss from ADSM was similar to CTSM, except for CTSM at the 3× rate applied in the autumn. The percentage of applied N lost as N<sub>2</sub>O was generally higher for treatments receiving CTSM compared to ADSM or UAN, while N<sub>2</sub>O losses from ADSM and UAN were similar. In summary, the findings suggest that ADSM is equal or better than CTSM in terms of agronomic performance, and has a lower environmental impact with regard to gaseous N loss.

#### 5. ACKNOWLEDGEMENTS

We are grateful for the funding provided by the Environmental Technologies Assessment for Agriculture (ETAA) program, the excellent collaboration from the Prairie Agricultural Machinery Institute (PAMI), for access to raw and digested material provided by Cudworth Pork Investors Group Ltd. and Clear-Green Environmental Inc.

#### REFERENCES

- [1] Koroluk, R., Culver, D., Lefebvre A. and McRae, T. (2000) Management of farm nutrient and pesticides inputs. In: McRae, T., Smith, C.A.S. and Gregorich, L.J., Eds., *Environmental Sustainability of Canadian Agriculture: Report of the Agri-Environmental Indicator Project*. Agriculture and Agri-Food Canada, Ottawa.
- [2] Laguë, C., Gaudet, É., Agnew J. and Fonstad, T.A. (2005) Greenhouse gas emissions from liquid swine manure storage facilities in Saskatchewan. *Transactions of the American Society of Agricultural Engineering*, **48**, 2289-2296.
- [3] Mooleki, S.P., Schoenau, J.J., Hultgreen, G., Wen, G. and Charles, J.L. (2002) Effect of rate, frequency and method of liquid swine manure application on soil nitrogen availability, crop performance and N use efficiency in east-central Saskatchewan. *Canadian Journal of Soil Science*, **82**, 457-467. doi:10.4141/S01-057
- [4] Pain, B.F., Phillips, V.R., Clarkson, C.R. and Klarenbeek, J.V. (1989) Loss of nitrogen through volatilization during and following the application of pig or cattle slurry to grassland. *Journal of Science, Food and Agriculture*, **47**, 1-12. doi:10.1002/jsfa.2740470102
- [5] Gordon, R., Jamieson, R., Rodd, V., Patterson, G. and Harz, T. (2001) Effects of surface manure application timing on ammonia volatilization. *Canadian Journal of Soil Science*, **81**, 525-533. doi:10.4141/S00-092

- [6] Rochette, P., Chantigny, M.H., Angers, D.A., Bertrand, N. and Cote, D. (2001) Ammonia volatilization and soil nitrogen dynamics following fall application of pig slurry on canola crop residues. *Canadian Journal of Soil Science*, **81**, 515-523. [doi:10.4141/S00-044](https://doi.org/10.4141/S00-044)
- [7] Chantigny, M.H., Angers, D.A., Rochette, P., Belanger, G. and Masse, D. (2007) Gaseous nitrogen emissions and forage nitrogen uptake on soils fertilized with raw and treated swine manure. *Journal of Environmental Quality*, **36**, 1864-1872. [doi:10.2134/jeq2007.0083](https://doi.org/10.2134/jeq2007.0083)
- [8] Kirchmann, H. and Lundvall, A. (1993) Relationship between N immobilization and volatile fatty acids in soil after application of pig and cattle slurry. *Biology and Fertility of Soils*, **15**, 161-164. [doi:10.1007/BF00361605](https://doi.org/10.1007/BF00361605)
- [9] Sommer, S.G. and Husted, S. (1995) The chemical buffer system in raw and digested animal slurry. *Journal of Agricultural Science*, **124**, 45-53. [doi:10.1017/S0021859600071239](https://doi.org/10.1017/S0021859600071239)
- [10] Sommer, S.G. and Olesen, J.E. (1991) Effects of drymatter content and temperature on ammonia loss from surface-applied cattle slurry. *Journal of Environmental Quality*, **20**, 679-683. [doi:10.2134/jeq1991.00472425002000030029x](https://doi.org/10.2134/jeq1991.00472425002000030029x)
- [11] Sommer, S.G. and Sherlock, R.R. (1996) pH and buffer component dynamics in the surface layers of animal slurries. *Journal of Agricultural Science*, **127**, 109-116. [doi:10.1017/S002185960007742X](https://doi.org/10.1017/S002185960007742X)
- [12] Sommer, S.G. and Hutchings, N.J. (2001) Ammonia emission from field applied manure and its reduction. *European Journal of Agronomy*, **15**, 1-15. [doi:10.1016/S1161-0301\(01\)00112-5](https://doi.org/10.1016/S1161-0301(01)00112-5)
- [13] Rubaek, G.H., Henriksen, K., Petersen, J., Rasmussen B. and Sommer, S.G. (1996) Effects of application technique and anaerobic digestion on gaseous nitrogen loss from animal slurry applied to ryegrass (*Lolium perenne*). *Journal of Agricultural Science*, **126**, 481-492. [doi:10.1017/S0021859600075572](https://doi.org/10.1017/S0021859600075572)
- [14] Chantigny, M.H., Rochette, P., Angers, D.A., Masse, D. and Cote, D. (2004) Ammonia volatilization and selected soil characteristics following application of anaerobically digested pig slurry. *Soil Science Society of America Journal*, **68**, 306-312.
- [15] Kirchmann, H. and Bernal, M.P. (1997) Organic waste treatment and C stabilization efficiency. *Soil Biology and Biochemistry*, **29**, 1747-1753. [doi:10.1016/S0038-0717\(97\)00065-5](https://doi.org/10.1016/S0038-0717(97)00065-5)
- [16] Clemens, J. and Huschka, A. (2001) The effect of biological oxygen demand of cattle slurry and soil moisture on nitrous oxide emissions. *Nutrient Cycling in Agroecosystems*, **59**, 193-198. [doi:10.1023/A:1017562603343](https://doi.org/10.1023/A:1017562603343)
- [17] Oenema, O., Wrage, N., Velthof, G.L., Groenigen, J.W., Dolfing, J. and Kuikman, P.J. (2005) Trends in global nitrous oxide emissions from animal production systems. *Nutrient Cycling in Agroecosystems*, **72**, 51-65. [doi:10.1007/s10705-004-7354-2](https://doi.org/10.1007/s10705-004-7354-2)
- [18] Wulf, S., Maeting, M. and Clemens, J. (2002) Application technique and slurry co-fermentation effects on ammonia, nitrous oxide, and methane emissions after spreading: II. Greenhouse gas emissions. *Journal of Environmental Quality*, **31**, 1795-1801. [doi:10.2134/jeq2002.1795](https://doi.org/10.2134/jeq2002.1795)
- [19] Petersen, S.O. (1999) Nitrous oxide emissions from manure and inorganic fertilizers applied to spring barley. *Journal of Environmental Quality*, **28**, 1610-1618. [doi:10.2134/jeq1999.00472425002800050027x](https://doi.org/10.2134/jeq1999.00472425002800050027x)
- [20] Vallejo, A., Skiba, U.M., Garcia-Torres, L., Arce, A., Lopez-Fernandez, S. and Sanchez-Martin, L. (2006) Nitrogen oxides emission from soils bearing a potato crop as influenced by fertilization with treated pig slurries and composts. *Soil Biology and Biochemistry*, **38**, 2782-2793. [doi:10.1016/j.soilbio.2006.04.040](https://doi.org/10.1016/j.soilbio.2006.04.040)
- [21] Schoenau, J.J., Mooleki, P. and Qian, P. (2001) Maximizing the economic and environmental Benefit of Land Application of Animal Manure. Report # 96000131, Saskatchewan Agriculture, Food and Rural Revitalization. [www.agriculture.gov.sk.ca/19960131.pdf](http://www.agriculture.gov.sk.ca/19960131.pdf)
- [22] Grant, C.A., Jia, S., Brown, K.R. and Bailey, L.D. (1996) Volatile losses of NH<sub>3</sub> from surface-applied urea and urea-ammonium nitrate with and without the urease inhibitors NBPT or ammonium thiosulphate. *Canadian Journal of Soil Science*, **76**, 417-419. [doi:10.4141/cjss96-050](https://doi.org/10.4141/cjss96-050)
- [23] Technicon Industrial Systems. (1977) Industrial/simultaneous determination of nitrogen and/or phosphorus in BD acid digests. Industrial method No. 334-74W/Bt, Tarrytown.
- [24] Livingston, G.P. and Hutchinson, G.L. (1995) Enclosure-based measurement of trace gas exchange: Applications and sources of error. In: Matson, P.A. and Harris, R.C., Eds., *Biogenic Trace Gases: Measuring Emissions from Soil and Water*, Blackwell Science Ltd., Oxford, 14-51.
- [25] Rochette, P., Angers, D.A., Belanger, G., Chantigny, M.H., Prevost, D. and Levesque, G. (2004) Emissions of N<sub>2</sub>O from alfalfa and soybean crops in eastern Canada. *Soil Science Society of America Journal*, **68**, 93-506. [doi:10.2136/sssaj2004.0493](https://doi.org/10.2136/sssaj2004.0493)
- [26] Rochette, P. and Hutchinson, G.L. (2005) Measuring soil respiration using chamber techniques. In: Hatfield, J. and Baker, J.M., Eds., *Micrometeorological Studies of the Soil Plant-Atmosphere Continuum*, American Society of Agronomy, Madison, 227-266.
- [27] Anthony, W.H., Hutchinson, G.L. and Livingston, G.P. (1995) Chamber measurement of soil-atmosphere gas exchange: Linear vs diffusion-based flux models. *Soil Science Society of America Journal*, **59**, 1308-1310. [doi:10.2136/sssaj1995.03615995005900050015x](https://doi.org/10.2136/sssaj1995.03615995005900050015x)
- [28] Lemke, R.L., Izaurralde, R.C., Nyborg, M. and Solberg, E.D. (1999) Tillage and N-source influence soil-emitted nitrous oxide in the Alberta Parkland region. *Canadian Journal of Soil Science*, **79**, 15-24. [doi:10.4141/S98-013](https://doi.org/10.4141/S98-013)
- [29] SAS Institute, Inc. (2004) SAS product documentation. Version 8. SAS Institute, Inc., Cary. <http://support.sas.com/documentation/onlinedoc/index.html>
- [30] Amon, B., Kryvoruchko, V., Amon, T. and Zechmeister-Boltenstern, S. (2006) Methane, nitrous oxide and am-

monia emissions during storage and after application of dairy cattle slurry and influence of slurry treatment. *Agriculture, Ecosystem and Environment*, **112**, 153-162.  
[doi:10.1016/j.agee.2005.08.030](https://doi.org/10.1016/j.agee.2005.08.030)

(2004) Presence of potential ammonia oxidation (PAO) inhibiting substances in anaerobic digestion residues. *Applied Soil Ecology*, **26**, 107-112.  
[doi:10.1016/j.apsoil.2003.12.002](https://doi.org/10.1016/j.apsoil.2003.12.002)

[31] Nyberg, K., Sundh, I., Johansson, M. and Schnürer, A.