

Molecular Study of the Genetic Variability of Pumpkins Landraces from Brazilian Amazon

Lúcia Helena Pinheiro Martins¹, Maria Teresa Gomes Lopes², Hiroshi Noda³, Pedro de Queiroz Costa Neto², Ayrton Luiz Urizzi Martins², Fabíola Viana de Almeida⁴, Liane Cristine Rebouças Demosthenes⁵

¹Núcleo de Etnoecologia da Amazônia (NETNO) Universidade Federal do Amazonas, Manaus, Brazil

²Faculdade de Ciências Agrárias (FCA), Universidade Federal do Amazonas, Manaus, Brazil

³Instituto Nacional de Pesquisas da Amazônia (INPA), Manaus, Brazil

⁴Instituto Federal de Educação Tecnológica do Amazonas/Campus Parintins, IFAM/Parintins, Brazil

⁵Instituto de Ciências Exatas e Tecnologia (ICET), Universidade Federal do Amazonas, Itacoatiara, Brazil

Email: luciahp.martins@yahoo.com.br

How to cite this paper: Martins, L.H.P., Lopes, M.T.G., Noda, H., de Queiroz Costa Neto, P., Martins, A.L.U., de Almeida, F.V. and Demosthenes, L.C.R. (2018) Molecular Study of the Genetic Variability of Pumpkins Landraces from Brazilian Amazon. *Agricultural Sciences*, 9, 511-523.

<https://doi.org/10.4236/as.2018.94035>

Received: March 14, 2018

Accepted: April 28, 2018

Published: April 30, 2018

Copyright © 2018 by authors and Scientific Research Publishing Inc.

This work is licensed under the Creative Commons Attribution International License (CC BY 4.0).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

Abstract

The *Cucurbita maxima* Duchesne is a vegetable crop plant cultivated and maintained by traditional Amazon communities, Brazil. The situation is worsened by the possibility of disappearance of local populations and genetic variability of this specie, taking into account the today changes promoted in family farming. The aim of this study was to estimate the current levels of genetic variability of local cultivars through the use of molecular markers (Amplified Fragment Length Polymorphism—AFLP). We chose to collect in two distinct micro regions in order to identify possible influences of geographic isolation and different levels of market requirements in the conservation of the genetic variability of the *C. maxima*. For the molecular analysis, bulk samples of fresh leaves of 15 plants/half-sibling family were collected in paper bags. There were 34 samples from the half-sib families. The analysis of the results half-sib obtained by methods of estimation of genetic variation by molecular markers shows that the forms of cultivation and management adopted by family farmers maintain the identities of the local/landraces (native cultivars) and, at the same time, the levels of diversity for the assurance of adaptability macro-environmental.

Keywords

Crop Genetic Variability, Molecular Marker, Amplified Fragment Length Polymorphism, *Cucurbita maxima* Duchesne

1. Introduction

The *Cucurbita maxima* Duchesne is a vegetable crop plant cultivated and maintained by traditional Amazon communities, Brazil. The *C. maxima* species known as Jerimum Caboclo. It is present in the production system of the regional family agriculture and in the list of plants of the diet of these people [1]. It can be consumed in various ways, its ripe and immature fruits, flowers and leaves as vegetables, edible seeds and ripe fruits as forage [2] [3]. Although Brazil is considered a center of diversity of pumpkins and squash, there is little knowledge about the germplasm maintained by family farming in the Amazon. The situation is worsened by the possibility of disappearance of local populations and genetic variability of this specie, taking into account the today changes promoted in family farming.

Five *Cucurbita* species are cultivated plants (*C. argyrosperma* Huber., *C. ficifolia* Bouché, *C. maxima* Duchesne, *C. moschata* Duchesne and *C. pepo* L.) [4]. The native species of the Americas were part of the food base, already in the time of the Olmec civilization, later incorporated by the Aztec, Inca and Maia civilizations. Even today, species of Cucurbitaceae are in the system of ancestral indigenous production complex in Mexico called *milpa*. The pumpkins (*C. moschata* and *C. argyrosperma*) are cultivated associated with legumes such as beans (*Phaseolus* spp. or *Vigna unguiculata* (L.) Walp.) and maize (*Zea mays* L.) [5]. In Brazil, *C. moschata* and *C. maxima* were part of the diet of indigenous peoples before colonization [6].

All species of the genus *Cucurbita* are diploid, with 20 pairs of chromosomes ($2n = 40$). The plants are annual, presenting herbaceous stem, creeping, scandal or subshrub, equipped with tendrils and adventitious roots. The branches can reach 10 meters in length. The leaves are large, dark green in color. The fruits have varied shapes and sizes; In *C. maxima*, the peduncle is of circular section [7].

The pumpkins are cross-breeding plants, and pollination is carried out by insects (entomophilic pollination) [8]. The reproductive system is characterized by cross-pollination which is favored by monoic flowers. The flowers open in the morning and have the bees as the main group of pollinators.

The production ratio of male and female flowers varies greatly. Environments of high temperatures favor the formation of male flowers and milder temperatures, feminine flowers. The development of the plant is differentiated; however, synchronization occurs in the anthesis of male and female flowers [9].

The selection allowed the maintenance of great genetic variability among and within the cultivated species. Variability is associated with a diversity of uses attributed to interests and needs in relation to different formats and sizes [10].

The molecular marker AFLP (Amplified Fragment Length Polymorphism) technique has been used in the evaluation of genetic diversity of lineages, clones and germplasm bank accesses. The technique has advantages such as the reproducibility of amplified fragments and the speed of analysis due to the fluores-

cence primer labeling system [11].

The advantage of generating a large number of polymorphisms per reaction and the need for prior knowledge of DNA sequence data for the construction of the primers used [12]. The technique consists in: digestion of the total genomic DNA with restriction enzymes, ligation of adapters with known sequence at the two ends of each fragment and selective amplification of the fragment set using primers [13] [14]. The amplified fragment marker may be an ideal tool for genetic studies and, among them, for genetic diversity.

So many techniques are available for the detection of genetic polymorphism, that is, the determination of genetic variability at the DNA sequence level. Molecular markers are DNA characteristics that differentiate two or more individuals and are inherited genetically [15]. They can be defined as any and all molecular phenotypes derived from an expressed gene or reference points on the chromosomes [16].

The AFLP technique has great capacity for the detection of genetic variability and use in characterization of cultivars. It is based on selective PCR amplification of restriction fragments from a total digest of genomic DNA [13]. It consists of the cleavage of the subject's genomic DNA using two restriction enzymes, followed by the use of specific adapters, which are attached to the ends of the cleaved DNA fragments. PCR amplifications of the DNA fragments and high resolution gel electrophoresis are performed to visualize the generated fragments [17]. Among the advantages of using AFLP are the high degree of polymorphism detected and the number of markers obtained by gel analyzed [18].

The aim of this study was to estimate the current levels of genetic variability of local cultivars through the use of molecular markers (Amplified Fragment Length Polymorphism—AFLP).

2. Materials and Methods

The properties of family farmers who were cultivating *C. maxima* were identified in rural communities of Benjamin Constant and Iranduba cities, Amazonas (Table 1). The samples were fruits from plants grown by family farms and adapted to the Amazonian floodplain environment, with climatic characteristics equatorial, hot and humid. We chose to collect in two distinct micro regions in order to identify possible influences of geographic isolation and different levels of market requirements in the conservation of the genetic variability of the *C. maxima*.

Six plants were sampled, distributed in such a way that, on the whole, the constitution of the seeds of the six fruits contained the contribution of pollen from all the existing plants in the cultivated area. Crop lands were stratified and sampling were adopted the procedures for outcrossing species, such as the case of pumpkin landraces, to collect material in field conditions with gametic control [19].

The objective of this step was to obtain frequent and continuous fresh vegetable material (fresh leaves) for the DNA extraction step. The seeds of the

Table 1. General identification of landraces *Cucurbita maxima* collected in Benjamin Constant and Iranduba, AM. 2012/2013.

City	Locality	Geographic Coordinates	Landraces	Half-sib families
Benjamin Constant	São Luís	04°24'14.7" 069°55'09.8"	SL	SL1, SL2, SL3, SL4, SL5, SL7
Benjamin Constant	Novo Lugar	04°22'48.3" 069°57'54.1"	NL	NL1, NL2, NL3, NL4, NL5, NL7
Benjamin Constant	São José	04°20'27.5" 069°57'15.8"	SJ	SJ3, SJ4, SJ5, SJ6, SJ8, SJ9
Benjamin Constant	Novo Paraíso	4°19'30.48" 69°59'8.77"	NP	NP1, NP2, NP3, NP4, NP5, NP6
Iranduba	Ilha da Paciência	03°19'16.32" 60°11'09.83"	IA	IA1, IA2, IA3, IA5, IA7, IA8
Iranduba	Ilha da Paciência	03°19'16.32" 60°11'09.83"	IB	IB1, IB2, IB3, IB4, IB5, IB6

fruit samples (from) were seeded in plastic cups of 300 mL containing substrate for the production of seedlings of vegetables and organic compound (3: 1), in a greenhouse of the Sector of Horticulture of the Institute of Agricultural Sciences of the Federal University of Amazonas. All half-sib families collected previously were represented in the sowing plastic cups. Sowing occurred in a staggered way, that is, each week, three half-sib families previously identified with the original collection codes were chosen and placed for sowing, with 15 replicates.

For the molecular analysis, bulk samples of fresh leaves of 15 plants/half-sibling family were collected in paper bags. There were 34 samples from the half-sib families.

The pool leaves corresponding to each family were initially stripped and cut into small pieces, weighing about 60 mg and pooled to be later macerated in a pistil crucible. Each sample for extraction represented 15 sheets of different individuals from one family.

For the extraction, the protocol adapted from [16] was used [20]. The DNA pellet was resuspended in 30 or 50 µL of TE buffer solution plus RNase and incubated at 37°C for 1 hour for further storage at -20°C.

Quantification of the DNA by the direct reading equipment, the reading method in Nanodrop (Model Spectrophotometer 2000 - Thermo Scientific) was used.

2.1. Molecular Marker AFLP

The digestion reaction step consisted of the use of 350 ng of genomic DNA in a solution prepared with 5.0 µL of One Phor All buffer, 0.5 µL of BSA 'bovine serum albumin', 0.25 µL of the Mse-I enzyme and 0.5 µL of the Eco RI enzyme by adding Milli-Q water to complete a final volume of 50 µL. Reactions occurred in 3 hours at a temperature of 37°C and final 15 minutes at 70°C in the Esco Swift

Max Pro.

At adapter attachment, the digestion fragments were ligated to Mse-I and Eco RI specific adapters, added to the T4 DNA Ligase enzyme (Thermo Scientific) and the T4 DNA Ligase buffer. Added 40 μL of digested DNA and filled with Milli-Q water to a final volume of 50 μL . The samples were incubated in a thermocycler (Esco Swift Max Pro) for 3 hours at 23°C.

Pre-amplification was performed with specific primers Mse-I + A (50 ng/ μL) and Eco RI + C (ng/ μL). 0.5 μL of each primer (Eco RI + C rare cut and frequently Mse-I + A, 2.0 μL of 5 mM dNTPs (Promega), 2.0 μL 10x buffer (UniScience), 0.6 μL 50 mM MgCl₂ (UniScience), 2.4 μL of Taq DNA polymerase (5 U) (Ferments), 2.0 μL of ligated DNA and 10 μL of Milli-Q water for one The PCR program [21] was developed in 26 cycles of amplification after denaturation at 94° for 2 minutes. The cycles were constituted of 94° for 1 minute, annealing at 56° per 1 minute and extension at 72° for 1 minute. The final cycle was performed at 72° for 5 minutes (protocol adapted [13]). Selection was tested from 14 of the oligonucleotide combinations and four combinations were selected because of the better number of polymorphic *loci* and visualization quality of the bands (Table 2).

The selective amplification was done to a final solution volume of 20 μL , with 0.2 μL of Taq DNA Polymerase (5 U/ μL) (Fermentas), 2.0 μL of 10x buffer (UniScience), 0.6 μL of MgCl₂ 2.50 (UniScience), 0.8 μL of 5 mM dNTPs, 1.0 μL Eco-RI, 1.2 μL Mse-I, 3.0 μL of pre-amplified DNA and 11.2 μL Milli-Q water. The PCR program consisted of denaturation at 94°C for 2 minutes; 12 cycles of 30 seconds at temperatures of 94°C, 65°C for 30 seconds each and 72°C for one minute; then 23 cycles at 94°C, 56°C for 30 seconds and 72°C for one minute; finally, a step at 72°C for two minutes. The samples were stored at -20°C until used in the next step. The samples were heated for five minutes at a temperature of 95°C for denaturation. At each electrophoresis, a pre-run was performed for cleaning and heating the gel for one hour at 80 W of power, in 1X TEB buffer in electrophoresis system (GT Gene model, BioRad source—Power Pac HV, 3000 V) [22].

For the application of the denatured samples, 8 μL of loading buffer was applied to the Polyacrylamide gel and 50 μL Ladder (50 - 800 bp) DNA marker

Table 2. Sequence of primer combinations chosen for the selective amplification of AFLP markers. Plant Genetic Improvement Laboratory. UFAM. 2015.

<i>Primers AFLP combination</i>	
Mse-I + CTC	Eco RI + AGC
Mse-I + CTC	Eco RI + AGT
Mse-I + CTC	Eco RI + ACA
Mse-I + CAT	Eco RI + ACA

from Promega and 2 μ L from Loading Dye (Promega). The samples were applied on 6% polyacrylamide gel in tempered glass plates of dimensions 38 \times 50 cm, submitted to electrophoresis, in a source of 3000 V, at 300 mA of intensity, 80 W of power and 2500 V of tension by a period of 4 hours.

The gels were staining with silver nitrate [23]. The steps involved solutions: fixation (acetic acid and absolute ethanol); pre-treatment (nitric acid); impregnation (silver nitrate); of development (sodium carbonate); and blocking (glacial acetic acid). The steps were sandwiched with distilled water.

2.2. Analysis of Amplified Fragments

Bands were visualized in the range of 100 to 750 bp of the combinations used and counted for presence (1) or absence (0), generating binary data matrices, both by combination and by all combinations. The number of polymorphic loci was calculated from those in which at least one of the samples had a difference in presence or absence of band. The percentage of polymorphic sites by combining from the total bands in each of four combinations was estimated by genetic analysis of the binary matrix.

2.3. Grouping Analysis

The matrix of similarity with the data of the general matrix of the four joint combinations was generated by the *Jaccard* coefficient [24] and a dendrogram was elaborated by the UPGMA algorithm (Unweighted Pair-Group Average). The *Jaccard* coefficient (S_j) prioritizes the samples by the presence similarity and is based on the equation:

$$S_j = \frac{a}{a+b+c}$$

S_j : Jaccard coefficient

a : number of coincidences of type 1-1 for each pair of accesses;

b : number of dislocations of type 1-0 for each pair of accesses;

c : number of disagreements of type 0-1 for each pair of accesses.

The coefficient of correlation coefficient was estimated by the NTSYS-pc software version 2.1/2000 [25]. The reliability and robustness of the dendrogram were tested by the bootstrap analysis with 1000 replications by NTSYS-pc version 2.1/2000 [25].

2.4. Correspondence Analysis

A simple correspondence analysis of the 34 families of *C. maxima* half-sib based on three main axes was performed on 246 discrete distribution characteristics by the NTSYS-pc version 2.1/2000 statistical package [25]. We attempted to identify the set of variables that could form the composite variables that best explained the DNA samples. Correspondence analysis is a method of sorting for categorical data counting data [26].

3. Results and Discussion

3.1. Polymorphism of *Cucurbita maxima*

The most informative combinations of the 14 tested were selected because of the better response of polymorphic *loci*. The total *loci* of all four combinations was 246 and polymorphism was 81, equivalent to 34%. The highest percentage of polymorphism was detected by the combination Mse-I + CTC/Eco RI + ACA with 48% (**Table 3**).

Molecular studies with 40 microsatellite *loci* in *C. pepo* Rio Grande do Sul cultivated showed that there is great genetic variability landraces, distributed among different materials, although it also exists within populations. The main contributing causes are the alogamy of the species, the exchange of seeds between the farmers and the cultivation of more than one variety within the same area. The study also pointed out that there is no subdivision of populations according to the collection site [27].

Other molecular studies with genetic markers have also indicated the existence of variability between and within the accesses collected at the different sites investigated. For example, *C. moschata* collected in Colombia Department detected high genetic diversity in agreement with the wide morphoagronomic diversity registered for the species. Most of the variation was related among individuals within the same Department (88.76%) [28]. In the classification analysis, the 121 introductions were grouped in three groups, the first one of the Andean Region, the second formed by some introductions of the Andean Region and the Andean Coast and the third harboring most of the introductions of the Atlantic Coast [28].

3.2. Genetic Variability among Half-Sib Families Based on the AFLP Marker

In the analysis of the generated dendrogram we can observe the formation of six groups that coincide with the local cultivars separated by the geographic distances (**Figure 1**). The cut line was 0.92 of similarity (dashed line). The analysis of the melt-level behavior [29] was used to determine the groups number in the final partition.

Table 3. Total number and polymorphic fragments by AFLP molecular markers in *Cucurbita maxima* observed in four primer combinations. Laboratory of Plant Genetic Improvement. UFAM. 2015.

Combination of AFLP <i>primers</i>		<i>C. maxima</i> variability		
		Total number <i>Loci</i>	Number of polimorphic <i>loci</i>	Polymorphism %
Mse-I + CTC	Eco RI + AGC	53	9	17
Mse-I + CTC	Eco RI + AGT	78	28	36
Mse-I + CTC	Eco RI + ACA	81	39	48
Mse-I + CAT	Eco RI + ACA	34	5	15




Figure 1. Dendrogram of 34 samples of half-sib families from *Cucurbita maxima* formed by the coefficient of similarity of *Jaccard*, by the algorithm UPGMA. Cophenetic correlation (0.8445). Manaus, 2015.

Group I brings together all the families of half-sib of the local variety Iranduba A (IA), originating in the Lower Solimões River. Group II is formed by five families of half-sib of the local variety Novo Lugar (NL), located in Ilha do Aramaçá, Benjamin Constant, Upper Solimões River.

Group III gathers all the families of half-sib local variety São Luís (SL), locality located to the left margin of the channel of the Solimões-Amazonas River (Upper Solimões River). This group also houses a family of half-sib (NL7) of the local NL variety, separated by the geographical distance of 5.7 km in a straight line. The fourth group is home to three half-sib families (SJ3, SJ4 and SJ5) from the local variety of São José (SJ), all from Ilha do Aramaçá. In group V all half-sib families of the local variety Novo Paraíso (NP) are reunited, in addition to SJ8 and SJ9. Novo Paraíso is located on Bom Intento Island, Benjamin Constant, and São José on a nearby island, 3.9 km in a straight line. Group VI houses all the families of half-sib of the local variety Iranduba B (IB) located in the Ilha da Paciência, Iranduba (Lower Solimões River). We can infer that in the molecular study, the largest distance (5.7 km) between different local cultivars occurred in Group III, which brought together the families of half-sib of São Luís and only one of Novo Lugar, different from the morphological characteristics that grouped geographically distant local cultivars. This result also contributes to confirm the existence of morphoagronomic and genetic variability among and within local cultivars of *C. maxima* landraces cultivated by family farms.

The groups were formed according to the geographically separated local cultivars. Even those different varieties that participated in the groups of another

Table 4 Similarity matrix detected in 34 samples from half-sib families of *Cucurbita maxima* Duchesne by molecular marker AFLP using *Jaccard* coefficient. Manaus, 2015.

	IA1	IA2	IA3	IA5	IA7	IA8	IB1	IB2	IB3	IB4	IB5	IB6	NI1	NI2	NI3	NI4	NI5	NI7	NP2	NP3	NP4	NP5	NP6	SJ3	SJ4	SJ5	SJ8	SJ9	SL1	SL2	SL3	SL4	SL5	SL7			
IA1																																					
IA2	0.9310																																				
IA3	0.9375	0.9645																																			
IA5	0.9650	0.9375	0.9716																																		
IA7	0.9789	0.9510	0.9577	0.9580																																	
IA8	0.9384	0.9789	0.9580	0.9315	0.9448																																
IB1	0.8472	0.8592	0.8786	0.8531	0.8662	0.8414																															
IB2	0.8219	0.8723	0.8652	0.8403	0.8403	0.8542	0.9685																														
IB3	0.8163	0.8794	0.8723	0.8472	0.8345	0.8611	0.9609	0.9921																													
IB4	0.8493	0.8611	0.8936	0.8811	0.8681	0.8562	0.9538	0.9538	0.9615																												
IB5	0.8414	0.8662	0.8723	0.8472	0.8601	0.8611	0.9609	0.9764	0.9688	0.9615																											
IB6	0.8750	0.8611	0.8936	0.8811	0.8944	0.8562	0.9538	0.9389	0.9318	0.9695	0.9615																										
NI1	0.9301	0.9296	0.9091	0.8966	0.9366	0.9371	0.8310	0.8182	0.8125	0.8082	0.8252	0.8333																									
NI2	0.9301	0.9296	0.9091	0.8966	0.9366	0.9371	0.8310	0.8182	0.8125	0.8082	0.8252	0.8333	1.0000																								
NI3	0.9301	0.9296	0.9091	0.8966	0.9366	0.9371	0.8310	0.8182	0.8125	0.8082	0.8252	0.8333	1.0000	1.0000																							
NI4	0.9236	0.9231	0.9028	0.8904	0.9301	0.9306	0.8380	0.8252	0.8194	0.8151	0.8322	0.8403	0.9926	0.9926	0.9926																						
NI5	0.8600	0.8591	0.8400	0.8289	0.8658	0.8667	0.7891	0.7770	0.7718	0.7566	0.7718	0.7800	0.9225	0.9225	0.9225	0.9161																					
NI7	0.7945	0.8182	0.8239	0.8000	0.8125	0.8138	0.8222	0.8222	0.8162	0.8116	0.8296	0.8248	0.8551	0.8551	0.8551	0.8623	0.8500																				
NP2	0.8125	0.7986	0.8042	0.8056	0.8056	0.8069	0.8284	0.8421	0.8358	0.8444	0.8636	0.8582	0.7958	0.7958	0.7958	0.8028	0.7793	0.8682																			
NP3	0.8138	0.8125	0.8182	0.8069	0.8069	0.8207	0.8296	0.8296	0.8235	0.8188	0.8370	0.8321	0.8357	0.8357	0.8357	0.8429	0.8310	0.8984	0.9360																		
NP4	0.8276	0.8264	0.8322	0.8207	0.8345	0.8309	0.8175	0.8116	0.8071	0.8248	0.8333	0.8500	0.8500	0.8571	0.8451	0.8846	0.9213	0.9839																			
NP5	0.8662	0.8264	0.8451	0.8462	0.8592	0.8345	0.8175	0.7914	0.7857	0.8201	0.8116	0.8333	0.8369	0.8369	0.8310	0.8194	0.8421	0.9365	0.9370	0.9375																	
NP6	0.8462	0.8069	0.8380	0.8392	0.8392	0.8151	0.7971	0.7842	0.7786	0.8000	0.8043	0.8261	0.8429	0.8429	0.8369	0.8380	0.8485	0.9134	0.9291	0.9297	0.9449																
SJ3	0.8690	0.8681	0.9007	0.8750	0.8881	0.8759	0.8613	0.8478	0.8417	0.8633	0.8686	0.8905	0.8531	0.8531	0.8601	0.8356	0.9160	0.8797	0.8806	0.8815	0.8676	0.8741															
SJ4	0.8345	0.8333	0.8652	0.8403	0.8531	0.8414	0.8657	0.8519	0.8456	0.8676	0.8731	0.8815	0.8440	0.8440	0.8440	0.8264	0.9070	0.8992	0.9000	0.9008	0.8864	0.8931	0.9615														
SJ5	0.8562	0.8425	0.8611	0.8493	0.8750	0.8503	0.8613	0.8478	0.8417	0.8633	0.8686	0.8768	0.8403	0.8403	0.8403	0.8472	0.8231	0.8872	0.8797	0.8806	0.8676	0.8676	0.8741	0.9549	0.9466												
SJ8	0.7959	0.8194	0.8125	0.8014	0.8014	0.8151	0.8235	0.8370	0.8309	0.8129	0.8444	0.8261	0.8169	0.8169	0.8169	0.8239	0.8125	0.9063	0.9286	0.9600	0.9449	0.9000	0.8923	0.8881	0.9077	0.8881											
SJ9	0.8333	0.8194	0.8511	0.8392	0.8392	0.8151	0.8370	0.8235	0.8175	0.8394	0.8444	0.8667	0.8169	0.8169	0.8169	0.8239	0.8125	0.8915	0.9286	0.9444	0.9449	0.9297	0.9370	0.9167	0.9375	0.9023	0.9524										
SL1	0.8288	0.8531	0.8592	0.8345	0.8472	0.8483	0.8321	0.8321	0.8261	0.8214	0.8394	0.8478	0.8777	0.8777	0.8777	0.8849	0.8723	0.9603	0.8779	0.9225	0.9231	0.8657	0.8864	0.9394	0.9308	0.8963	0.9302	0.9302									
SL2	0.8493	0.8611	0.8671	0.8425	0.8552	0.8690	0.8540	0.8540	0.8478	0.8429	0.8613	0.8696	0.8993	0.8993	0.8993	0.9065	0.8936	0.9380	0.8722	0.9015	0.9023	0.8467	0.8667	0.9328	0.9242	0.8905	0.8947	0.9615									
SL3	0.8493	0.8611	0.8671	0.8425	0.8552	0.8690	0.8540	0.8540	0.8478	0.8429	0.8613	0.8696	0.8993	0.8993	0.8993	0.9065	0.8936	0.9380	0.8722	0.9015	0.9023	0.8467	0.8667	0.9328	0.9242	0.8905	0.8947	0.9615	1.0000								
SL4	0.8483	0.8601	0.8662	0.8414	0.8542	0.8552	0.8947	0.8881	0.8824	0.9023	0.8963	0.8582	0.8582	0.8582	0.8582	0.8652	0.8531	0.9077	0.8855	0.9008	0.8872	0.8593	0.8657	0.9179	0.9091	0.9179	0.8939	0.8939	0.9167	0.9542	0.9542						
SL5	0.8345	0.8462	0.8521	0.8276	0.8403	0.8414	0.8519	0.8519	0.8456	0.8406	0.8593	0.8540	0.8705	0.8705	0.8777	0.8652	0.9524	0.8992	0.9147	0.9008	0.8722	0.8788	0.9318	0.9231	0.9030	0.9077	0.9077	0.9609	0.9690	0.9690	0.9535						
SL7	0.8276	0.8521	0.8582	0.8333	0.8462	0.8345	0.8582	0.8582	0.8519	0.8467	0.8657	0.8603	0.8633	0.8705	0.8582	0.9600	0.8915	0.9070	0.8931	0.8647	0.8712	0.9389	0.9302	0.9098	0.9147	0.9147	0.9685	0.9612	0.9612	0.9457	0.9920						




Figure 2. Dispersion of 34 families of *Cucurbita maxima* half-sib families, based on the main axes (1 and 2) of the simple correspondence analysis on 246 discrete distribution characteristics (polymorphism in 81). Circled in blue are the families of half-sib of Iranduba (Lower Solimões River) and in red, those of Benjamin Constant (Upper Solimões River). Amazonas, 2015.

origin, are located nearby Benjamin Constant. This can be explained by the fact that the marker technique performs structural characterization, that is, the AFLP has a large genome coverage, however the observed polymorphism is related to the genetic structure. Different from the morphoagronomic techniques that they identify by the phenotypic expression.

Genetic resources are not only linked to genotypes, but everything that is linked to them, such as information, environment, socioeconomic and ecological [30]. It includes the set of environmental factors (pests, weeds) and cultural factors (associated traditional knowledge).

Molecular markers Random Amplified Polymorphic DNA (RAPD) [31], using Amplified Fragment Length Polymorphism (AFLP) [32] and using microsatellite markers contributed to describe and confirm the genetic variability of *Cucurbita* species, especially *C. moschata* and *C. maxima* for the conservation in agricultural crops of varieties by farmers [33].

The genetic variability of the half-sib families was detected by the similarity matrix generated by the *Jaccard* coefficient, with values in the range of 0.7786 and 1.00 (Table 4). The greatest similarity between half-sib families occurred with the comparison between samples NL1 and NL2, NL1 and NL3, NL2 and NL3, SL2 and SL3. The smallest similarities were observed among the NP6 and IB3 families (0.7786).

The calculated values for the class interval of the similarity parameters presented mean and variance of 0.87 and 0.002, respectively. It was found that 70.4% of the sample values are in the range of 0.8 to 0.89 and all values were grouped above 0.7, indicating the number of occurrences and the percentage of distances genetics organized by the interval classes.

The distribution of the different samples of half-sib families by the main axes of the correspondence analysis (**Figure 2**). Two groups according to geographic location were formed: the first one formed by the half-sib families of the local cultivars of Upper Solimões River (NL, NP, SJ and SL) and the second group bringing together the families of half-sib families IA and IB, geographically located in the municipality of Iranduba.

This result shows that the methodology used was able to separate the variability between the local cultivars (Upper Solimões *versus* Lower Solimões Rivers).

4. Conclusion

The analysis of the results obtained by methods of estimation of genetic variation by molecular markers shows that the forms of cultivation and management adopted by family farmers maintain the identities of the local/native cultivars and, at the same time, the levels of diversity for the assurance of adaptability macro-environmental.

References

- [1] Fonseca, M.A., Queiroz, M.A., Lopes, J.F., Assis, J.G.A., Romão, R.L., Ramos, S.R.R., Carmo, C.A.S., Moura, M.C.L., Dias, R.C.S., Silveira, L.M., Barbieri, R.L., Silva, D.J.H. and Noronha, S. (2015) Geographical Distribution and Conservation of Cucurbita in Brazil. *Magistra, Cruz das Almas, BA*, **27**, 432-442.
- [2] Esquinas-Alcazar, J.T. and Gulick, P.J. (1983) Genetic Resources of Cucurbitaceae. IPBGR, Rome, 101.
- [3] Saade, R.L. and Hernández, S.M. (1992) La Agricultura em Mesoamerica: *Cucurbitas*. In: Hernández Bermejo, J.E. and León, J., Eds., *Cultivos Marginados. Outra perspectiva de 1492*. Colección FAO: Producción y protección vegetal, No. 26, Roma.
- [4] Heiden, G.; Barbieri, R. L.; Neitzke, R. S. (2007) Key for the identification of species of pumpkins (*Cucurbita*, Cucurbitaceae) grown in Brazil. (Documents, 197). Embrapa Temperate Weather, Pelotas, 31.
- [5] Canul Ku, J. Vallejo, P.R., González, F.C. and Servia, J.L.C. (2005) Morphological Diversity of Squash Grown in the Center-East of Yucatan, Mexico. *Revista Fitotecnia Mexicana*, **28**, 339-349.
- [6] Carvalho, P.G.B., de Peixoto, A.A.P. and Ferreira, M.A.J.F. (2011) Characterization of Pumpkins as to the Levels of Total Carotenoids Alpha and Betacarotene (Research and Development Bulletin 78). Embrapa Hortaliças, Brasília, 20.
- [7] Lima, G.K.L. (2013) Rescue and Germplasm Study of *Cucurbita* spp. of Rio Grande do Norte. Thesis, Mossoró.
- [8] Romano, C.M., Stumpf, E.R.T., Barbieri, R.L., Bevilaqua, G.A.P. and Rodrigues, W.F. (2008) Manual Pollination in Pumpkins. Embrapa Temperate Weather (Documents, 225), Pelotas, 26.
- [9] OECD, Organization for Economic Co-operation and Development, Environment, Health and Safety Publications (2012) Consensus Document on the Biology of *Cucurbita* L. (Squashes, Pumpkins, Zucchini and Gourds). Series on Harmonization of Regulatory Oversight in Biotechnology. No. 53 ENV/JM/MONO31, 77.

- [10] Bisognin, D.A. (2002) Origin and Evolution of Cultivated Cucurbits. *Rural Science. Santa Maria*, **32**, 715-723. <https://doi.org/10.1590/S0103-84782002000400028>
- [11] Pasqual, M., Rezende, R.K.S., Villa, F. and Chagas, E.A. (2008) Biotechnology Applied to the Improvement of Fruit Trees. In: Bruckner, C.H., Ed., *Fundamentals of Fruit Tree Breeding*, UFV, Viçosa, 117-170.
- [12] Faleiro, F.G. (2007) Genetic-Molecular Markers Applied to Conservation and Use of Genetic Resources. Embrapa, Cerrados, Planaltina, 102.
- [13] Vos, P., Hogers, R., Bleeker, M., Reijans, M., van de Lee, M.T., Hornes, M., Frijters, A., Pot, J., Peleman, J., Kuiper, M. and Zabeau, M. (1995) AFLP: A New Technique for DNA Fingerprinting. *Nucleic Acids Research*, **23**, 4407-4414. <https://doi.org/10.1093/nar/23.21.4407>
- [14] Caixeta, E.T., de Oliveira, A.C.B., de Brito, G.G. and Sakiyama, N.S. (2009) Types of Markers. In: Borém, A. and Caixeta, E.T. (Eds), *Molecular Markers*, Folha de Viçosa, Viçosa, 11-93.
- [15] Milach, S.C. K. (1998) DNA Markers. Applications in Plant Breeding. *Biotechnology Science & Development*, **5**, 14-17.
- [16] Ferreira, M.E. and Grattapaglia, D. (1998) Introduction to the Use of Molecular Markers in Genetic Analysis. 3rd Edition, Document 20, Embrapa-Cenargen, Brasília, 220 p.
- [17] Brammer, S.P. (2000) Molecular Markers: Basic Principles and Use in Plant Breeding Programs. http://www.cnpt.embrapa.br/biblio/p_do03.htm
- [18] Milach, S.C.K. (1999) Molecular Markers on Genetic Resources and Plant Breeding. In: Queiroz, M.A., Goedert, C.O. and Ramos, S.R.R., Eds., *Genetic Resources and Plant Breeding for the Brazilian Northeast*, Embrapa Semiárido, Brasília, DF, Embrapa Genetic Resources and Biotechnology, Petrolina. <http://www.cpsa.embrapa.br:8080/catalogo/livroorg/index.html>
- [19] Vencovsky, R. (1987) Effective Population Size in the Collection and Preservation of Germplasm Outcrossing Species. *IPEF*, **35**, 79-84.
- [20] Martins, L.H.P. (2015) Genetic Variability and Conservation of *Cucurbita maxima* Duchesne by Family Agriculture in the Western Central Amazon. Thesis, Federal University of Amazonas, 151 p.
- [21] Couch, J.A. and Fritz, P.J. (1990) Isolation of DNA from Plants High in Polyphenolics. *Plant Molecular Biology Reporter*, **8**, 8-12. <https://doi.org/10.1007/BF02668875>
- [22] Lopes, R., Lopes, M.T.G., Carneiro, M.S., Matta, F.P., Camargo, L.E.A. and Vieira, M.L.C. (2006) AFLP Linkage Analysis and Mapping of Resistance Genes to *Xanthomonas axonopodis* pv. *passiflorae* n Yellow Passion Fruit. *Genome*, **49**, 17-29. <https://doi.org/10.1139/g05-081>
- [23] Creste, S., Tulmann Neto, A. and Figueira, A. (2001) Detection of Single Sequence Repeat Polymorphisms in Denaturing Polyacrylamide Sequencing Gels by Silver Staining. *Plant Molecular Biology Reporter*, **19**, 299-306. <https://doi.org/10.1007/BF02772828>
- [24] Cruz, C.D. (2008) GENES Program: Genetic Diversity. Viçosa. Ed. UFV. 278 p.
- [25] Rohlf, F.J. (2000) NTSYS-pc, Version 2.1. Numerical Taxonomy and Multivariate Analysis System. Exeter Software, New York.
- [26] Hammer, Harper, D.A.T. and Ryan, P.D. (2012) PAST: Paleontological Statistics Software Package for Education and Data Analysis. *Electronic Paleontology*, **4**, 187 .

- [27] Priori, D., Barbieri, R.L., Castro, C.M., Oliveira, A.C., Vilella, J.C.B. and Mistura, C.C. (2012) Molecular Characterization of Creole Varieties of Pumpkins with Microsatellite Markers. *Brazilian Horticulture*, **30**, 499-506. <https://doi.org/10.1590/S0102-05362012000300024>
- [28] Restrepo, J.A. and Vallejo, C.F.A. (2008) Molecular Characterization of Colombian Introductions of Zapallo *Cucurbita moschata*. *Acta Agronómica (Palmira)*, **57**, 9-17.
- [29] Mingoti, S.A. (2005) Data Analysis through Multivariate Statistical Methods: An Applied Approach. Belo Horizonte. UFMG.
- [30] Clement, C.R., Rizzi Rocha, S.F., Cole, D.M. and Vivan, J.L. (2007) Conservation on Farm. In: Nass, L., Ed., *Plant Genetic Resources*, Embrapa Genetic Resources and Biotechnology, Brasília, 511-544.
- [31] Gwanama, C., Labuschagne, M.T. and Botha, A.M. (2000) Analysis of Genetic Variation in *Cucurbita moschata* by Random Amplified Polymorphic DNA (RAPD) Markers. *Euphytica*, **113**, 19-24. <https://doi.org/10.1023/A:1003936019095>
- [32] Ferriol, M.B.P., Cordova, P.F. and Nuez, F. (2004) Molecular Diversity of a Germplasm Collection of Squash (*Cucurbita moschata*) Determined by SRAP and AFLP Markers. *Crop Science*, **44**, 654-663.
- [33] Watcharawongpaiboon, N. and Chunwongse, J. (2007) Development of Microsatellite Markers from an Enriched Genomic Library of Pumpkin (*Cucurbita moschata* L.). *Songklanakarin Journal of Science and Technology*, **29**, 1217-1223.