

# Detection of *Salmonella enterica* Serovar Rissen in Slaughter Pigs in Northern Ireland

David A. Egan, Violetta Naughton\*, James S. Dooley, Patrick J. Naughton

Biomedical Sciences Research Institute, School of Biomedical Sciences, Ulster University, Coleraine, UK

Email: \*v.naughton@ulster.ac.uk

**How to cite this paper:** Egan, D.A., Naughton, V., Dooley, J.S. and Naughton, P.J. (2017) Detection of *Salmonella enterica* Serovar Rissen in Slaughter Pigs in Northern Ireland. *Advances in Microbiology*, 7, 513-522.

<https://doi.org/10.4236/aim.2017.76040>

**Received:** May 11, 2017

**Accepted:** June 23, 2017

**Published:** June 26, 2017

Copyright © 2017 by authors and  
Scientific Research Publishing Inc.

This work is licensed under the Creative  
Commons Attribution International  
License (CC BY 4.0).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

## Abstract

*Salmonella enterica* serovar Rissen has been recognised as a common serovar in humans and pigs around the world. This study investigated *S. Rissen* prevalence in pigs slaughtered in Northern Ireland additionally looking at antibiotic susceptibility profiles, genetic profiles and plasmid profiles to provide information on an emerging non-typhoid *Salmonella* serotype with the potential to cause disease in humans. *S. Rissen* were isolated on five separate sampling occasions from both the boning hall and slaughter line of a randomly selected single pig abattoir in Northern Ireland (NI). Following antibiotic susceptibility testing against 16 antibiotics, all *S. Rissen* isolates were identified as susceptible to 15 antibiotics but resistant to tetracycline (R-type: Te). Of the 29 *S. Rissen*, 27 were isolated from pigs originating in NI and two *S. Rissen* were isolated from pigs originating in the Republic of Ireland (RoI). The combined results of the PFGE and plasmid profiling analyses were capable of subdividing the *S. Rissen* isolates into three distinct groups. The data suggests that *S. Rissen* is an emerging serovar in Northern Ireland and continued surveillance of this serovar is warranted as it has the potential to cause disease in humans.

## Keywords

Antibiotic Susceptibility Testing, Pulsed Field Gel Electrophoresis, Plasmid Profiles

## 1. Introduction

The most recent reports identify *Salmonella* as the most common causative agent in food-borne outbreaks with known origin in the European Union (EU) [1]. The most recently published European Food Safety Authority and European Centre for Disease Prevention and Control (EFSA and ECDC) report identified *Salmonella enterica* serovar Rissen amongst the 20 most frequent serovars asso-

ciated with cases of human Salmonellosis and as one of the top 10 serovars associated with pig meat in the EU [2].

*S. Rissen* is a commonly reported serotype around the world and it is amongst the top three serotypes found in pigs and pork products in Europe and Southeast Asia [3]. Numerous studies have identified *S. Rissen* in slaughtered pigs in Portugal [4] and within pig fattening units and from slaughter-age pigs and Spain [5] [6] [7] [8] [9]. There have been further reports of *S. Rissen* in pig abattoir studies in Italy [10], Belgium [11], the Netherlands [12], Portugal [13] as well as in breeding herds in the UK [14] and in finishing pigs in Northern Ireland [15], thus demonstrating its' increase in European pig herds.

Worldwide *S. Rissen* is considered to be to be one of the most prevalent causes of human salmonellosis. In the USA in 2009 *S. Rissen* was responsible for a major *Salmonella* outbreak infecting more than 80 people over four states [16] and [17]. *S. Rissen* infections in humans have been reported in Ireland, Demark and the UK [18] [19] [20]. This clearly demonstrates that pigs continue to act as a carrier/host of *S. Rissen* infection and hence a potential reservoir of human infection. The risk of salmonellosis in humans coupled with the rise of multi-drug resistance in *Salmonella* highlights the need for the continued surveillance of emerging *Salmonella* serovars including *S. Rissen*. This study aims to add to previous research within the pig industry whilst adding to the continued assessment of the *Salmonella* serovars in pigs. This study investigated *S. Rissen* isolated from pigs slaughtered in Northern Ireland additionally looking at antibiotic susceptibility profiles, genetic profiles and plasmid profiles to provide information on an emerging non-typhoid *Salmonella* serotype with the potential to cause disease in humans.

## 2. Materials and Methods

### 2.1. Sampling

Sampling was carried out over an 18 month period. In all 405 samples were taken at the randomly selected single (out of seven) abattoir. For the oyster cut 25 - 50 g trimming was removed from the 50:50 product tray (50:50 is a half fat half pork trimming from the oyster cut) and transferred to a sterile jar (Trafalgar Scientific (CON7764) Leicester, UK). For faecal and caecal sampling the viscera was removed from the pig's abdominal cavity and following separate incisions into the rectum and caecum approximately 30 - 50 g of sample was removed and placed into sterile jars. Carcass swabs were taken from four EU approved swabbing points identified on the pig's carcass, jowl, belly, back and ham (EU directive 471/2001/EU) using EnviroSponge™ USDA approved cellulose sponges [Biotrace International (BP-133ES) Bridgend, UK] and placed in sterile bags provided with the sponge and labelled.

All samples were examined for the presence of *Salmonella* according to ISO 6579. Briefly 25 g of sample was transferred into 225 ml ISO Buffered peptone Water [BPW, Oxoid (CM1049) Basingstoke, UK], The sample and ISO BPW was transferred to a stomacher bag [Seward (BA6141) Worthing, West Sussex,

UK] and stomached for 1 min (Seward Tekmar Stomacher 400, Seward, Worthing, West Sussex, UK). The sample was incubated at 37° for 18 - 20 hours. Controls comprised *S. Nottingham* NCTC 7832 (positive control) and *Escherichia coli* NCTC10418 (negative control). There followed selective enrichment in Rappaport-Vassiliadis Soya (RVS) broth [Oxoid (CM0866) Basingstoke, UK], plating onto selective Xylose Lysine Decarboxylase [Oxoid (CM0469) Basingstoke, UK] and Brilliant Green Agar [Oxoid (CM0263) Basingstoke, UK] followed by performance of confirmatory tests on suspect colonies.

All suspect *Salmonella* were sent to Agri-Food & Biosciences Institute (AFBNI), Newforge Lane, Belfast, Northern Ireland for serotyping. All resulting *Salmonella* were subsequently sent to The Laboratory of Enteric Pathogens, HPA, Collindale, UK to be phage typed using a standard set of the typing phage and following the Collindale scheme.

## 2.2. Antibiotic Susceptibility Testing

Resistance profiling used BSAC's disc diffusion method

(<http://www.bsac.org.uk/wp-content/uploads/2012/02/Version-8-January-2009.pdf>)

according to established protocols [21]. All *S. Rissen*, were tested for their *in vitro* sensitivity to 16 different antibiotics. The antibiotics and their concentrations were as follows; Amikacin 30 µg (AK), Ampicillin 10 µg (AMP), Amoxycillin & Clavulanic Acid 30 µg (AMC), Apramycin 15 µg (APR), Cefotaxime 30 µg (CTX), Ceftazidime 30 µg (CAZ), Chloramphenicol 10 µg (C), Ciprofloxacin 1 µg (CIP), Compound Sulphonamides 300 µg (Su), Furazolidone 15 µg (FR), Gentamicin 10 µg (Cn), Nalidixic Acid 30 µg (NA), Neomycin 10 µg (N), Streptomycin 25 µg (S), Sulphamethoxazole/trimethoprim 25 µg (SXT) and Tetracycline 10 µg (T). Minimum inhibitory concentration (MIC) tests were carried using M.I.C. Evaluators (Oxoid, Basingstoke, UK) where resistance was identified. Multidrug resistance was defined as having a resistance pattern to five or more antibiotics.

## 2.3. Pulsed Field Gel Electrophoresis (PFGE)

PFGE was performed according to the CDC PulseNet protocol [22] with some minor modifications. *Salmonella* isolates were streaked onto Tryptone Soya Agar [TSA, Oxoid (CM0131) Basingstoke, UK] and incubated at 37°C for 18 - 20 hours. *Salmonella* isolates from TSA were resuspended in 2 ml of cell suspension buffer (CSB) (100 mM Tris: 100mM EDTA, pH 8.0) creating at attenuation of approximately 1.35 at  $\lambda$  610 using a spectrophotometer. 10 µl of Proteinase K (20 mg/ml stock) [Sigma-Aldrich (P6556) Pole, Dorset, UK] was added to 200 µl of *Salmonella* suspension and gently mixed. 200 µl of molten 2% SeaKem Gold agarose [Cambrex (BMA 50150) Wokingham, Berkshire, UK] was then added to this suspension. *S. Branderup* H9812 was used as the reference strain. The bacterial/agarose mixture was then dispensed into PFGE disposable moulds [BioRad (170-3713) Hemel Hempstead Hertfordshire, UK] and allowed to solidify for approximately 10 minutes. The freshly moulded plugs were then transferred to

tubes containing 1.5 ml of CSB and stored at 4°C. For cell lysis the plugs were placed in a 1.5 ml eppendorf containing 1 ml Cell Lysis Buffer-CLB (50 mM EDTA, pH 8, 1% sarcosyl) and 40 µl of Proteinase K solution (20 mg/ml) and incubated in a shaking incubator at 54°C for 2 hours. The plugs were then washed in series of steps. Firstly, in a Polypropylene Super Clear™ 50 ml sterile tube, [Sarstedt (SLS8106) Leicester, UK] containing 10 - 15 ml of deionised water in a shaking incubator at 50°C for 10 minutes and following subsequent washing steps were placed in fresh TE buffer and stored at 4°C until restriction digestion. The restriction enzyme *Xba*I [Sigma-Aldrich (R7260) Poole, Dorset, UK] was first diluted 1:10 with dilute SuRE Cut buffer H [11417991001; Roche, UK] and 100 µl of this restriction mixture were added the micro-centrifuge tube containing the agarose plugs. Both sample and control plug slices were gently mixed and incubated for two hours at 37°C.

A 1% agarose gel was prepared using SeaKem® Gold agarose [Cambrex (BMA50150) Woking, Berkshire, UK]. The gel plugs were then overlaid with a small quantity of molten SeaKem® Gold agarose [Cambrex (BMA50150) Woking, Berkshire, UK] and allowed to solidify. PFGE was conducted in a CHEF DR II chamber (BioRad, Hertfordshire, UK) containing fresh 0.5 M Tris Borate EDTA (TBE) running buffer [Gibco, (15581-044) Paisley, UK] under the following conditions: Initial switch time 2.2 s; Final switch time 63.8 s; Start Ratio 1.0; Voltage 200 V; Run time 19 - 20 hours at 14°C. After staining for 20-30 min (in 40 µl ethidium bromide (EtBr) [Sigma-Aldrich (160539) Poole, Dorset, UK] and de-staining in sterile water for 60 min the gel image was then visualised and captured using a transillumination table (AlphaImager™ 2200, Alpha Innotech, CA, USA). All PFGE gel images were converted to TIFF files prior to data analysis. Dendrograms and cluster analysis were created using Bionumerics 5.1 software (Applied Maths, Belgium). Similarity analysis was performed using Dice coefficient, and clustering was created using the unweighted pair group method with arithmetic means (UPGMA). When isolates were found to be indistinguishable upon *XBa*I digestion and additional enzyme *Spe*I, was used to improve discrimination.

## 2.4. Plasmid Profiling

Plasmid profiling was carried out using a modified version of the Kado and Liu method [23] [24]. A single colony was inoculated into 3 ml of Brain Heart Infusion [Oxoid (CM1032) Basingstoke, UK] and following incubation at 37°C for 18 - 20 hours 1ml of the culture centrifuged at 6600 G for two minutes. Cells were lysed by addition of 100 µl of lysis buffer (3% SDS in 50 mM Tris, pH 12.6). The sample was incubated at 56°C for 30 minutes in a pre-heated heat block, followed by a chloroform extraction. The upper aqueous layer was transferred to a 1ml eppendorf. Loading dye (6 - 8 µl) was added to each sample and vortexed (sample stored at -20° for later analysis). Each sample (12 µl) was loaded onto a 0.7% agarose gel with a super-coiled ladder [Sigma-Aldrich (D5292) Poole, Dorset, UK] as molecular marker. Electrophoresis run-25 volts for four hours

using a RunOne electrophoresis system (EmbiTec, CA, USA). The gel matrix was then stained using EtBr [Sigma-Aldrich (160539) Poole, Dorset, UK]. The resulting bands were viewed using a transilluminator (AlphaImager™ 2200, Alpha Innotech Corp., USA).

### 3. Results

In all 29 *S. Rissen*, were isolated from the abattoir, 21 of which were isolated from pork cuts in the boning hall, while four came from the rectal contents of pigs (post-slaughter), two were isolated from caecum contents of pigs (post-slaughter) and one was from a carcass swab (post-chill). The *Salmonella* were isolated over an 18-month period on five separate sampling occasions. Two of the pigs positive for *S. Rissen* in terms of rectal samples were of Republic of Ireland (RoI) origin, the remaining *S. Rissen* were isolated from pigs originating in Northern Ireland. *S. Rissen* accounted for 59% of the *Salmonella* identified, followed by *S. typhimurium* (21%), *S. panama* (17%) and *S. meleagridis* (3%).

#### 3.1. Antibiotic Susceptibility

All 29 *S. Rissen* isolates exhibited resistance to only a single antibiotic-tetracycline at a concentration of 10 µg/l. The MIC for Tetracycline was determined to be >256 µg/ml. No resistance to the fluoroquinolone antibiotic ciprofloxacin or to the third generation cephalosporin antibiotics cefotaxime and ceftazidime was observed.


#### 3.2. PFGE Analysis Using Xba1 and Spe1

Consistent and unambiguous Xba1-PFGE profiles were generated from 23 of the 29 isolates and analysis of these identified 19 that displayed a common profile consisting of 15 bands which ranged in size from 20 kb to 668 kb approximately. These 19 *S. Rissen* were designated cluster A and all presented a similarity higher than 96% - 97% using the DICE co-efficient. The remaining four *S. Rissen* displayed a common profile consisting of 16 bands which ranged in size from approximately 20 kb to 668 kb. These were designated cluster B and represented a similarity of approximately 97%. As both cluster A and B generated such similar Xba1 profiles further PGFE analysis using the restriction enzyme Spe1, as an alternative to Xba1, was conducted on representative *S. Rissen* from both cluster A and cluster B (**Figure 1**). Spe1 generated RFLP profiles that were identical for all *S. Rissen* typed. This analysis when combined with that of the Xba1 profiles and analysed using the BioNumerics 5.1 software (Applied Maths, Belgium) indicated that all the *S. Rissen* isolates from the pig abattoir were a clonal match. No further genetic differences were observed.


#### 3.3. Plasmid Profiling

All 29 *S. Rissen* isolates were observed to have a low molecular weight band plasmid of approximately 4.650 kbp. A second low molecular weight plasmid of approximately 2.750 kb in size was found to be present in three out of the 29

isolates (Figure 2). All three of these *S. Rissen* that harboured the 2.750 kbp plasmid were recovered on a single sampling visit and were isolated from pigs originating in NI. Three other *S. Rissen* isolates were recovered that did not harbour the 2.750 kbp low molecular weight plasmid (Figure 2). The three *S.*



**Figure 1.** Pulse Field Gel Electrophoresis analysis of 7 of the *S. Rissen* isolated from a NI abattoir restricted with both *Spe*I and *Xba*I and the dendrogram comparing the cluster analysis based on the combined macro-restriction profiles is shown. The 7 *S. Rissen* isolates: B70 (Pig faeces), B85 (Pig Faeces), B92 (Pig Faeces), C96 (Carcass Post Chill), E300 (Boning Hall pork), E63 (Boning Hall Pork), T72 (Pig Caecum). Two *S. typhimurium* isolates: H061 (laboratory isolate) and T108 (Pig Caecum) have been included for comparison purposes. The numbers on the horizontal axis represent the percent similarly between isolates based on dice coefficient and UGPMA clustering.



**Figure 2.** Plasmid profiles of *S. Rissen* isolates. Lanes 1 - 5 (*S. Rissen* isolates T70, T71, T72 from Pig Caecum). Lanes 6 - 7 (*S. Rissen* isolates B70, B71 from Pig Faeces) and lane 8 (*S. Rissen* C96 from a Carcass Swab). A low molecular weight plasmid (4650 KB) is observed in lanes 1, 2 and 5.



Rissen isolates displaying two plasmids were not the same isolates as the three *S. Rissen* observed to contain the extra band Xba1-PFGE in cluster B. The identification of two plasmid profile types significantly enhanced the differentiation of *S. Rissen* that were genetically dissimilar. When the genotyping data obtained through plasmid profiling is combined with that obtained by PFGE three different *S. Rissen* genetic profiles can be observed.

#### 4. Discussion

All 29 *S. Rissen* exhibited resistance to a single antibiotic tetracycline (T) while displaying no resistance to the other 15 antibiotics at the concentrations tested in this study. Of the *S. Rissen* isolated from pigs in Europe, tetracycline is the most common resistance phenotype [7] [19] [25]. Garcia-Feliz [26] investigating the antibiotic resistance of a panel of 114 *S. Rissen* isolates, all originating from pigs, recorded that over 50% of isolates were resistant to tetracycline alone. This study supports other research that multidrug resistance is less common amongst *S. Rissen* isolates when compared to the antibiotic resistance characteristics of *S. Typhimurium* isolates [19] [26]. However, *S. Rissen* displaying multidrug resistance has been recorded including a study conducted in Spain which identified 48 of 114 *S. Rissen* isolates as MDR ranging from four to nine antimicrobials [26] and Thailand where *S. Rissen* displayed resistance from one to five antibiotics [27].

PFGE utilizing the restriction enzyme Xba1 resulted in two distinctive PFGE profiles among the 23 *S. Rissen* which were designated cluster A and cluster B. The remaining profiles were identical to the profiles observed for Cluster A and the *S. Rissen* isolates in cluster B that contained the extra band were still categorised as closely related [28]. The three *S. Rissen* in cluster B were isolated from pigs that were exported from the Republic of Ireland (RoI) for slaughter in NI. Two of the three *S. Rissen* isolated from these RoI reared pigs were recovered from faecal samples suggesting that these pigs may have been harbouring this *S. Rissen* clone on entering the abattoir.

PFGE and Plasmid profiling were used to establish the genetic diversity amongst the *S. Rissen* and suggests that the 23 *S. Rissen*, although not 100% identical, were all closely related. Xba1 generated PFGE images for *S. Rissen* from pigs and pork published by [4] [10] appear to have a common primary set of Xba1 generated restriction fragments consisting of approximately 15 restriction fragments similar to this study suggests that a common Xba1 pulsotype may exist for *S. Rissen* throughout Europe but further investigations would be necessary to confirm this.

#### 5. Conclusion

PFGE identified two distinct profiles while plasmid profiling determined that three of the *S. Rissen* categorised as 100% clonal by PFGE actually carried an extra low molecular weight plasmid and of the 16 antibiotics tested against all 29 *S. Rissen* isolates, resistance to tetracycline was the only resistance phenotype ob-

served. The prevalence of this serovar is of some concern given that *S. Rissen* is capable of plasmid insertions and there may be a propensity for increases in *Salmonella* antibiotic resistance phenotypes. The literature would suggest that pigs are one of the main reservoirs of *S. Rissen* and it is likely that increased multidrug resistance in this serovar will see an increasing number of human cases. The continued surveillance of *S. Rissen* in pig herds is warranted.

## Acknowledgements

We are grateful to the School of Biomedical Sciences, University of Ulster for supporting this project.

## Conflict of Interest

The authors declare that they have no conflict of interest.

## References

- [1] Eurosurveillance Editorial Team (2015) The 2013 Joint ECDC/EFSA Report on Trends and Sources of Zoonoses, Zoonotic Agents and Food-Borne Outbreaks Published. *Eurosurveillance*, **20**, 21021.  
<http://www.eurosurveillance.org/ViewArticle.aspx?ArticleId=21021>
- [2] EFSA and ECDC (European Food Safety Authority and European Centre for Disease Prevention and Control) (2015) The European Union Summary Report on Trends and Sources of Zoonoses, Zoonotic Agents and Food-borne Outbreaks in 2013. *EFSA Journal*, **13**, 3991.  
<http://www.efsa.europa.eu/en/efsajournal/doc/3991.pdf>
- [3] Schmidt, J.W., Brichta-Harhay, D.M., Kalchayanand, N., Bosilevac, J.M., Shackelford, S.D., Wheeler, T.L. and Koohmaraie, M. (2012) Prevalence, Enumeration, Serotypes, and Antimicrobial Resistance Phenotypes of *Salmonella enterica* Isolates from Carcasses at Two Large United States Pork Processing Plants. *Applied and Environmental Microbiology*, **78**, 2716-2726.  
<https://doi.org/10.1128/AEM.07015-11>
- [4] Vieira-Pinto, M., Tenreiro, R. and Martins, C. (2006) Unveiling Contamination Sources and Dissemination Routes of *Salmonella* sp. in Pigs at a Portuguese Slaughterhouse through Macrorestriction Profiling by Pulsed-Field Gel Electrophoresis. *International Journal of Food Microbiology*, **110**, 77-84.  
<https://doi.org/10.1016/j.ijfoodmicro.2006.01.046>
- [5] García-Feliz, C., Collazos, J.A., Carvajal, A., Vidal, A.B., Aladueña, A., Ramiro, R., De La Fuente, M., Echeita, M.A. and Rubio, P. (2007) *Salmonella enterica* Infections in Spanish Swine Fattening Units. *Zoonoses and Public Health*, **54**, 294-300.  
<https://doi.org/10.1111/j.1863-2378.2007.01065.x>
- [6] García-Feliz, C., Carvajal, A., Collazos, J.A. and Rubio, P. (2007) Herd-Level Risk Factors for Faecal Shedding of *Salmonella enterica* in Spanish Fattening Pigs. *Preventive Veterinary Medicine*, **91**, 130-136.  
<https://doi.org/10.1016/j.prevetmed.2009.05.011>
- [7] Márquez, R.J.A., Echeita-Salaberria, A., Maldonado García, A., Valdezate-Jimenez, S., Carbonero-Martinez, A., Aladueña-García, A. and Arenas-Casas, A. (2007) Surveillance and Antimicrobial Resistance of *Salmonella* Strains Isolated from Slaughtered Pigs in Spain. *Journal of Food Protection*, **70**, 1502-1506.  
<https://doi.org/10.4315/0362-028X-70.6.1502>



- [8] Mejia, W., Casel, J., Zapata, D., Sánchez, G.J., Martin, M. and Mateu, E. (2006) Epidemiology of *Salmonella* Infections in Pig Units and Antimicrobial Susceptibility Profiles of the Strains of *Salmonella* Species Isolated. *Veterinary Record*, **159**, 271-276. <https://doi.org/10.1136/vr.159.9.271>
- [9] Arguello, H., Carvajal, A., Collazos, J.A., García-Feliz, C. and Rubio, P. (2012) Prevalence and Serovars of *Salmonella enterica* on Pig Carcasses, Slaughtered Pigs and the Environment of Four Spanish Slaughterhouses. *Food Research International*, **45**, 905-912. <https://doi.org/10.1016/j.foodres.2011.04.017>
- [10] Piras, F., Brown, D.J., Meloni, D., Mureddu, A. and Mazzette, R. (2011) Investigation of *Salmonella enterica* in Sardinian Slaughter Pigs: Prevalence, Serotype and Genotype Characterization. *International Journal of Food Microbiology*, **151**, 201-209. <https://doi.org/10.1016/j.ijfoodmicro.2011.08.025>
- [11] De Busser, E.V., Maes, D., Houf, K., Dewulf, J., Imberechts, H., Bertrand, S. and De Zutter, L. (2011) Detection and Characterization of *Salmonella* in Lairage, on Pig Carcasses and Intestines in Five Slaughterhouses. *International Journal of Food Microbiology*, **145**, 279-286. <https://doi.org/10.1016/j.ijfoodmicro.2011.01.009>
- [12] van Hoek, A.H.M., de Jonge, R., van Overbeek, W.M., Bouw, E., Pielaat, A., Smid, J.H., Malorny, B., Junker, E., Löfström, C., Pedersen, K., Aarts, H.J.M. and Heres, L. (2012) A Quantitative Approach towards a Better Understanding of the Dynamics of *Salmonella* spp. in a Pork Abattoir. *International Journal of Food Microbiology*, **153**, 45-52. <https://doi.org/10.1016/j.ijfoodmicro.2011.10.013>
- [13] Gomes-Neves, E., Antunes, P., Manageiro, V., Gärtner, F., Caniça, M., da Costa, J.M. and Peixe L. (2014) Clinically Relevant Multidrug Resistant *Salmonella enterica* in Swine and Meat Handlers at the Abattoir. *Veterinary Microbiology*, **168**, 229-233. <https://doi.org/10.1016/j.vetmic.2013.10.017>
- [14] Wales, A.D., McLaren, I.M., Bedford, S., Carrique-Mas, J.J., Cook, A.J. and Davies, R.H. (2009) Longitudinal Survey of the Occurrence of *Salmonella* in Pigs and the Environment of Nucleus Breeder and Multiplier Pig Herds in England. *Veterinary Record*, **165**, 648-657. <https://doi.org/10.1136/vr.165.22.648>
- [15] Belsué, J.B., Alujas, A.M. and Porter, R. (2011) Detection of High Serological Prevalence and Comparison of Different Tests for *Salmonella* in Pigs in Northern Ireland. *Veterinary Record*, **169**, 153. <https://doi.org/10.1136/vr.d4382>
- [16] Flynn, D. (2010) How Salmonella Rissen Came to America. *Food Safety News*. <http://www.foodsafetynews.com/2010/08/how-salmonella-rissen-came-to-america/#.UjHt-mTwJSY>
- [17] Higa, J. (2011) Outbreak of *Salmonella* Rissen Associated with Ground White Pepper: The Epi Investigation. [http://www.cdph.ca.gov/programs/DFDRS/Documents/QSS\\_Presentation\\_SRissen\\_and\\_%20white%20pepper\\_010611.pdf](http://www.cdph.ca.gov/programs/DFDRS/Documents/QSS_Presentation_SRissen_and_%20white%20pepper_010611.pdf)
- [18] Irvine, N. (2009) Communicable Diseases Monthly Report, Northern Ireland Edition. The Health Protection Agency. <http://www.publichealth.hscni.net/directorate-public-health/health-protection/surveillance-data>
- [19] Hendriksen, R.S., Bangtrakulnonth, A., Pulsrikarn, C., Pornreongwong, S., Hasman, H., Song, S.W. and Aarestrup, F.M. (2008) Antimicrobial Resistance and Molecular Epidemiology of *Salmonella* Rissen from Animals, Food Products, and Patients in Thailand and Denmark. *Foodborne Pathogens and Disease*, **5**, 605-619. <https://doi.org/10.1089/fpd.2007.0075>
- [20] Foley, B., McKeown, P. and Cormican, M. (2005) Epidemiology of Human Salmonellosis in Ireland, 2003. *Epi-Insight*, **6**, 2-3.

- [21] Animal Health and Veterinary Laboratories Agency (AHVLA) (2014) Chapter 12: Antimicrobial Susceptibility in *Salmonella*: *Salmonella* in Livestock Production in Great Britain 2013. [http://vla.defra.gov.uk/reports/rep\\_salm\\_rep11.htm](http://vla.defra.gov.uk/reports/rep_salm_rep11.htm)
- [22] Ribot, E.M., Fair, M.A., Gautom, R., Cameron, D.N., Hunter, S.B., Swaminathan, B. and Barrett, T.J. (2006) Standardization of Pulsed-Field Gel Electrophoresis Protocols for the Subtyping of *Escherichia coli* O157:H7, *Salmonella*, and *Shigella* for PulseNet. *Foodborne Pathogens and Disease*, **3**, 59-67. <https://doi.org/10.1089/fpd.2006.3.59>
- [23] Brown, D. (2004) Plasmid Profile Analysis by the Method of Kado and Liu. Scottish Salmonella Reference Laboratory, Laboratory Procedure, Scotland.
- [24] Kado, C.I. and Liu, S.T. (1981) Rapid Procedure for Detection and Isolation of Large and Small Plasmids. *Journal of Bacteriology*, **145**, 1365-1373.
- [25] Botteldoorn, N., Herman, L., Rijpens, N. and Heyndrickx, M. (2004) Phenotypic and Molecular Typing of *Salmonella* Strains Reveals Different Contamination Sources in Two Commercial Pig Slaughterhouses. *Applied and Environmental Microbiology*, **70**, 5305-5314. <https://doi.org/10.1128/AEM.70.9.5305-5314.2004>
- [26] García-Feliz, C., Collazos, J.A., Carvajal, A., Herrera, S., Echeita, M.A. and Rubio, P. (2008) Antimicrobial Resistance of *Salmonella enterica* Isolates from Apparently Healthy and Clinically Ill Finishing Pigs in Spain. *Zoonoses and Public Health*, **55**, 195-205. <https://doi.org/10.1111/j.1863-2378.2008.01110.x>
- [27] Tadee, P., Boonkhot, P., Pornruangwong, S. and Patchanee, P. (2015) Comparative Phenotypic and Genotypic Characterization of *Salmonella* spp. in Pig Farms and Slaughterhouses in Two Provinces in Northern Thailand. *PLoS ONE*, **10**, e0116581.
- [28] Tenover, F.C., Arbeit, R.D., Goering, R.V., Mickelsen, P.A., Murray, B.E., Persing, D.H. and Swaminathan, B. (1995) Interpreting Chromosomal DNA Restriction Patterns Produced by Pulsed-Field Gel-Electrophoresis—Criteria for Bacterial Strain Typing. *Journal of Clinical Microbiology*, **33**, 2233-2239.



Scientific Research Publishing

**Submit or recommend next manuscript to SCIRP and we will provide best service for you:**

Accepting pre-submission inquiries through Email, Facebook, LinkedIn, Twitter, etc.

A wide selection of journals (inclusive of 9 subjects, more than 200 journals)

Providing 24-hour high-quality service

User-friendly online submission system

Fair and swift peer-review system

Efficient typesetting and proofreading procedure

Display of the result of downloads and visits, as well as the number of cited articles

Maximum dissemination of your research work

Submit your manuscript at: <http://papersubmission.scirp.org/>

Or contact [aim@scirp.org](mailto:aim@scirp.org)

