

Analysis of the Alkane Hydroxylase Gene and Long-Chain Cyclic Alkane Degradation in *Rhodococcus*

Taiki Kawagoe, Kenzo Kubota, Kiwako S. Araki, Motoki Kubo

Department of Biotechnology, Faculty of Life Sciences, Ritsumeikan University, Shiga, Japan

Email: kubo@sk.ritsumei.ac.jp

How to cite this paper: Kawagoe, T., Kubota, K., Araki, K.S. and Kubo, M. (2019) Analysis of the Alkane Hydroxylase Gene and Long-Chain Cyclic Alkane Degradation in *Rhodococcus*. *Advances in Microbiology*, 9, 151-163.

<https://doi.org/10.4236/aim.2019.93012>

Received: January 30, 2019

Accepted: March 2, 2019

Published: March 5, 2019

Copyright © 2019 by author(s) and Scientific Research Publishing Inc.

This work is licensed under the Creative Commons Attribution International License (CC BY 4.0).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

Abstract

To characterize the long-chain cyclic alkane (*c*-alkane) degradation of bacteria in *Rhodococcus*, we analyzed the relationship between the alkane hydroxylase gene (*alkB*) and long-chain *c*-alkane degradation in 19 species. Eleven strains which were isolated from nature using long-chain *c*-alkane as a substrate were identified as *R. erythropolis*, and all were shown to carry the *alkB* [*alkB* R2 type]. This gene type was also carried by two other species, *R. rhodochrous* and *R. baikonurensis*. In total, 17 species of the genus *Rhodococcus* carried *alkB*, but the gene types differed from each other. The two species *R. rhodnii* and *R. coprophilus* did not carry *alkB*, and their long-chain *c*-alkane degradation levels were low.

Keywords

Alkane Hydroxylase, *alkB*, Bioremediation, Long-Chain *c*-Alkane, *Rhodococcus*

1. Introduction

In recent years, environmental pollution by petroleum hydrocarbons has been a serious problem worldwide. To clean up hydrocarbon-contaminated sites, a range of treatments can be performed, including incineration, chemical oxidation (Fenton reaction), washing, and evaporation. However, a lot of energy is required for these treatments [1] [2] [3] [4]. Biological treatments such as bioremediation have been investigated as an effective system of hydrocarbon degradation [5] [6] [7] [8] [9], and many microorganisms with the ability to degrade petroleum hydrocarbons have been isolated from their natural habitat and characterized [10] [11] [12] [13] [14]. However, it is difficult for bacteria to degrade long-chain hydrocarbons, especially long-chain cyclic alkanes (*c*-alkanes) [15] [16]. Therefore,

the isolation of bacteria capable of degrading long-chain hydrocarbons is important for the bioremediation of hydrocarbons [17] [18] [19].

Kubota *et al.* previously isolated hydrocarbon-degrading bacteria (HDB) from their natural habitats, and determined their phylogenetic relationships based on partial sequences of the 16S rRNA gene [20]. This identified many isolated species as belonging to the genera *Pseudomonas*, *Rhodococcus*, *Gordonia*, and *Acinetobacter* [20]. The strains belonging to the genera *Rhodococcus* and *Gordonia* degraded not only long-chain normal alkanes (*n*-alkanes) but also *c*-alkanes as the sole carbon and energy source [20] [21] [22] [23]. This indicated that the strains degrading long-chain *c*-alkanes have potential for cleaning petroleum hydrocarbon pollutants, because long-chain *n*-alkanes and *c*-alkanes remain in the soil for long periods.

The first step of degrading long-chain *c*-alkanes, the oxidation of hydrocarbons, is catalyzed by alkane hydroxylases (Alk) [24] [25]. For many HDB, *alkB* genes encoding Alk have been well analyzed [26] [27] [28] [29]. *alkB* genes are categorized as seven types (*alkB1* to *alkB7*) [30] [31], and many strains of the genus *Rhodococcus* are known to carry *alkB2*. However, the relationship between different types of *alk* genes and the *c*-alkane degradation ability has not been characterized for each species. Moreover, *alkB* varies among species of the genus *Rhodococcus*, and oxidizable substrates differ depending on the *alkB* gene type [32]. For example, Fukuhara *et al.* [33] revealed that the *R. erythropolis* strain NDKK6, carrying the *alkB* R2 type, showed a high *c*-alkane degradation ability. Furthermore, the carbon number of the alkyl side chain seemed to influence *c*-alkane degradation [34] [35] [36] [37]. A previous study isolated 11 strains of HDB from the genus *Rhodococcus* [20], but they have not been characterized yet.

In the present study, we investigated the relationship between *alkB* genes and long-chain *c*-alkane degradation for 11 isolated strains of *Rhodococcus* and 19 strains obtained from the NITE Biological Resource Center (NBRC), Japan.

2. Materials and Methods

2.1. Bacterial Strains

Thirty bacterial strains were used as long-chain *c*-alkane-degrading bacteria. Eleven strains (*R. erythropolis* NDKK1, *R. erythropolis* NDKK2, *R. erythropolis* NDKK5, *R. erythropolis* NDKK6, *R. erythropolis* NDKK7, *R. erythropolis* NDKK48, *R. erythropolis* ODNM1C, *R. erythropolis* NDKY82A, *R. erythropolis* ODMI54, *R. erythropolis* ODNM2B, and *R. erythropolis* NDMI144) were isolated in our previous study [20], and 19 species (*R. erythropolis* NBRC15567, *R. rhodochrous* NBRC15564, *R. baikonurensis* NBRC100611, *R. wratislaviensis* NBRC100605, *R. opacus* NBRC100624, *R. ruber* NBRC15591, *R. equi* NBRC101255, *R. percoletus* NBRC100626, *R. jostii* NBRC16295, *R. triatomae* NBRC103116, *R. koreensis* NBRC100607, *R. corynebacterioides* NBRC14404, *R. zopfii* NBRC100606, *R. tukisamuensis* NBRC100609, *R. maanshanensis* NBRC100610, *R. pyridinivorans*

NBRC100608, *R. kroppenstedtii* NBRC103113, *R. rhodnii* NBRC100604, and *R. coprophilus* NBRC100603) were obtained from the NBRC.

2.2. Identification of Bacterial Strains

The 11 strains from our earlier study were identified based on the full length 16S rRNA gene sequence. These strains were cultivated in Luria-Bertani (LB) medium (1% peptone, 0.5% yeast extract, and 0.5% NaCl) at 30°C with shaking at 200 rpm [36]. Total DNA was extracted from the culture, and the 16S rRNA gene sequence was determined [38] [39]. Sequence data were deposited in the DNA Data Bank of Japan (DDBJ) database (<https://www.ddbj.nig.ac.jp/index-e.html>) under accession numbers LC107434 to LC107444. The sequences were also compared with data in the GenBank database using BLAST+ 2.2.31 which is available at <https://www.ncbi.nlm.nih.gov/blast/>. Phylogenetic analyses were carried out for the 11 strains and the additional 19 species whose sequence data were obtained from the GenBank database using ClustalW (version 2.1). The phylogenetic tree was constructed by the neighbor-joining method and bootstrap values were calculated by 1000 replications.

2.3. Sequences of *alkB* and *alkB* [*alkB* R2 Type] Genes

Sequences of *alkB* were also determined for 30 strains. PCR was conducted using primers *alkB*-F (5'-AACTAYMTCGARCAYTAYGG-3')/*alkB*-R (5'-TGRTCKSTCGYTGVARGTG-3') and *alkB* R2-F (5'-CGGTTGTGTGTCGCAGGA-TC-3)/*alkB* R2-R (5'-AACGACTGCGCCAGAGTGAT-3') [31] to confirm the presence of *alkB* and *alkB* genes [*alkB* R2 type], respectively. *alkB* and *alkB* gene [*alkB* R2 type] PCR products were 140 bp and 100 bp, respectively. *alkB* primers were designed for the consensus region of HDB. *alkB* sequences were determined by the ABI PRISM 3100 Genetic Analyzer (Applied Biosystems, Foster City, CA, USA), and sequence data were deposited in the DDBJ database under accession numbers LC107445 to LC107470. Phylogenetic analysis was performed for *alkB* sequences as described above.

2.4. Analysis of *c*-Alkane Degradation by *Rhodococcus* Strains

To evaluate bacterial long-chain *c*-alkane degrading abilities, 30 strains were cultivated in modified SW medium with undecylcyclohexane (UDC), dodecylcyclohexane (DDC), and tridecylcyclohexane (TDC) substrates. Pre-culture of each strain was prepared in LB medium. Substrate hydrocarbon (0.10 g) and pre-culture (1 ml) were inoculated in 100 ml of modified SW medium (1% v/w; per liter: 1.21 g NH₄NO₃, 14.3 g Na₂HPO₄·12H₂O, 5.44 g KH₂PO₄, 0.5 g NaCl, 0.247 g MgSO₄, 2.78 mg FeSO₄·7H₂O, 14.7 mg CaCl₂·2H₂O, 2.01 mg ZnSO₄·7H₂O, 0.15 mg [NH₄]₆Mo₇O₂₄·4H₂O, 2 mg CuSO₄·5H₂O, 0.4 mg CoCl₂·6H₂O, 1.49 mg MnSO₄·5H₂O, 0.5 g polypeptone, and 0.25 g yeast extract) in a baffle flask before incubating at 30°C with shaking at 120 rpm.

After 3 days of cultivation, 30 ml of chloroform-methanol (3:1) mixture was added to 100 ml medium and mixed well before centrifuging at 4000× g for 30 min. The separated organic layer was taken for gas-chromatographic analysis using a flame ionized detector (GC-FID) (GC-2010, Shimadzu, Kyoto, Japan). The degradation test was carried out in triplicate. The degradation ratio was calculated from the rate of decrease of the peak area of the gas-chromatogram as follows:

$$1 - (\text{Peak area after cultivation} / \text{Peak area before cultivation}) \times 100 (\%)$$

The concentration of hydrocarbon in the soil was measured using the OCMA-355 oil content analyzer (Horiba, Kyoto, Japan).

3. Results and Discussion

3.1. Identification of Long Chain *c*-Alkane-Degrading Bacteria and Analysis of Alkane Hydroxylase Genes in *Rhodococcus*

To identify *Rhodococcus* strains isolated in our previous study [20], we determined the 16S rRNA gene sequences (1429 - 1472 bp). Phylogenetic analysis showed that all strains belonged to *R. erythropolis*, based on similarity with the reference strain *R. erythropolis* MPU33 (DNA databank accession number: AB334770) (Figure 1). These strains were isolated from a culture medium containing long chain *c*-alkane as the sole carbon source [20], indicating that the strains carried *alkB*.

The presence and type of *alkB* genes in the strains were next examined. All strains carried *alkB* genes, with 13 *Rhodococcus* species of the 29 strains used in this study shown to carry *alkB* (Table 1 and Table 2). *alkB* sequences of *R. erythropolis* isolated from nature were 98.1% - 100% homologous with the *alkB* R2 type of *R. erythropolis* NDKK6 that has high *c*-alkane degradation ability (Table 1). The *alkB* sequence from *R. erythropolis* NBRC15567 also showed high similarity with *alkB* R2 type (95.7%). Additionally, two other strains (*R. rhodochrous* and *R. baikonurensis*) obtained from NBRC carried *alkB* genes with approximately 100% homology to *alkB* R2 type (Table 2).

Ten strains also carried *alkB* genes, but homology analysis showed that their sequences differed from *alkB* R2 type (21.9% to 78.3% homologous) so original *alkB* names were given to each *alkB* gene type (Table 2). The similarity of *R. opacus* and *R. percolatus* *alkB* genes was 99.7% (Figure 2), so they were named *alkB* ROP type. The remaining 6 species lacked *alkB* genes. Phylogenetic analysis based on *alkB* revealed separate clusters of *alkB* [*alkB* R2 type] and *alkB* [non-*alkB* R2 type] groups (Figure 2). In the *Rhodococcus* genus, 3 groups were categorized as follows: first group included species such as *R. baikonurensis* and *R. rhodochrous* carrying *alkB* R2 type, another group carried other types of *alkB* genes, and the third group lacked any *alkB* genes.

3.2. Degradation of Long Chain *c*-Alkane by *Rhodococcus* Species Carrying *alkB* R2 Type

The degradation rates of long chain *c*-alkanes (UDC, DDC, and TDC) by 14




Figure 1. Phylogenetic tree based on 16S rRNA gene sequences of 26 strains of genus *Rhodococcus*. Sequences of the strain shown in bold were determined in this study and the other strains were taken from the GenBank database (<https://www.ncbi.nlm.nih.gov/genbank/>). The tree was constructed by the neighbor-joining method. The scale bar indicates the number of substitutions per site. Bootstrap values (1000 replicates) greater than 50% are indicated in the corresponding nodes.

Table 1. The presence of *alkB* genes and similarity to *alkB* [*alkB*R2 type] genes.

Strain	<i>alkB</i> Gene	Similarity to <i>alkB</i> Gene [<i>alkB</i> R2 Type] (%)*
<i>R. erythropolis</i> NDKK1	+	99.6
<i>R. erythropolis</i> NDKK2	+	100
<i>R. erythropolis</i> NDKK5	+	99.9
<i>R. erythropolis</i> NDKK7	+	99.8
<i>R. erythropolis</i> NDKK48	+	99.7
<i>R. erythropolis</i> ODNM1C	+	99.8
<i>R. erythropolis</i> NDKY82A	+	98.1
<i>R. erythropolis</i> ODMI54	+	99.7
<i>R. erythropolis</i> ODNM2B	+	99.2
<i>R. erythropolis</i> NDMI144	+	99.7
<i>R. erythropolis</i> NBRC15567	+	95.7

* *alkB* gene [*alkB*R2 type] from *Rhodococcus erythropolis* NDKK6 [1].

Table 2. The presence of *alkB* genes similarity to *alkB* genes [*alkB* R2 type], and types of *alkB* gene.

Strain	<i>alkB</i> Gene	Similarity to <i>alkB</i> Gene [<i>alkB</i> R2 Type] (%)*	Type of <i>alkB</i> gene
<i>R. rhodochrous</i> NBRC15564	+	100	<i>alkB</i> R2
<i>R. baikonurensis</i> NBRC100611	+	99.7	<i>alkB</i> R2
<i>R. wratislaviensis</i> NBRC100605	+	22.6	<i>alkB</i> RW
<i>R. opacus</i> NBRC100624	+	23.4	<i>alkB</i> ROP
<i>R. ruber</i> NBRC15591	+	30.5	<i>alkB</i> RR
<i>R. equi</i> NBRC101255	+	78.3	<i>alkB</i> RE
<i>R. percolatus</i> NBRC100626	+	21.9	<i>alkB</i> ROP
<i>R. jostii</i> NBRC16295	+	30.1	<i>alkB</i> RJ
<i>R. triatoma</i> NBRC103116	+	30.1	<i>alkB</i> R1
<i>R. koreensis</i> NBRC100607	+	30.5	<i>alkB</i> RK
<i>R. corynebacterioides</i> NBRC14404	+	24.1	<i>alkB</i> RC
<i>R. zopfii</i> NBRC100606	+	30.7	<i>alkB</i> RZ
<i>R. tukisamuensis</i> NBRC100609	–		
<i>R. maanshanensis</i> NBRC100610	–		
<i>R. pyridinivorans</i> NBRC100608	–		
<i>R. kroppenstedtii</i> NBRC103113	–		
<i>R. rhodnii</i> NBRC100604	–		
<i>R. coprophilus</i> NBRC100603	–		

* *alkB* gene [*alkB* R2 type] from *Rhodococcus erythropolis* NDKK6 [1].

Rhodococcus strains carrying *alkB* R2 type are shown in **Table 3**. The average degradation rates of UDC, DDC, and TDC were 55.5%, 31.8%, and 55.2%, respectively. Strain ODNM2B showed the highest degradation rate (98.9%) of UDC. Similarly, strains NDKK2 and ODNM2B showed superior degradation of DDC and TDC (42.9% and 94.6%, respectively). **Figure 3** shows the degradation of UDC by *R. erythropolis* NDKK6. These observations of *R. erythropolis* strains including our identified strains and one from NBRC indicate that *alkB* R2 type plays an important role in the degradation of long chain *c*-alkanes, and that *Rhodococcus* carrying *alkB* R2 type can utilize *n*-hexadecane [40].

3.3. Degradation of Long Chain *c*-Alkane by Strains Carrying Non-*alkB* R2 Type

The degradation rates of long chain *c*-alkanes (UDC, DDC, and TDC) by 10 *Rhodococcus* species carrying non-*alkB* R2 type are shown in **Table 4**. Average degradation rates of UDC, DDC, and TDC were 49.4%, 29.4%, and 53.9%, respectively. Although strains of the genus *Rhodococcus* carrying *alkB* genes can degrade hydrocarbons such as long chain *c*-alkanes, the degradation rates of strains carrying non-*alkB* R2 type were lower than those carrying *alkB* R2 type




Figure 2. Phylogenetic tree based on alkane hydroxylase (*alkB*) gene sequences of 24 strains of the genus *Rhodococcus*. The scale bar indicates the number of substitutions per site. Corresponding gene names of each strain and group of strains are followed by names in parenthesis and braces, respectively. The tree was constructed by the neighbor-joining method. Bootstrap values (1,000 replicates) greater than 50% are indicated in the corresponding nodes.

(Table 3 and Table 4). This indicates that *alkB* R2 type was more suitable for the degradation of long chain *c*-alkanes. Moreover, the higher degradation rates of UDC and TDC compared with DDC suggest that long chain *c*-alkanes with an odd number of carbon atoms are easier to biodegrade by *Rhodococcus* species [21].

3.4. Degradation of Long Chain *c*-Alkane by Strains Carrying No *alkB* Genes


The degradation rates of long chain *c*-alkanes (UDC, DDC, and TDC) by 6 strains carrying no *alkB* genes are shown in Table 5. *R. rhodnii* NBRC100604 had the highest degradation rate of UDC, DDC, and TDC among all 6 strains. The average degradation rates of UDC, DDC, and TDC by these 6 strains were 19.3%, 13.3%, and 19.1%, respectively. The average degradation ratios of UDC, DDC, and TDC by strains carrying no *alkB* genes were lower than in these by strains carrying *alkB* [*alkB* R2 type] and *alkB* [non-*alkB* R2 type]. Strains of genus

Rhodococcus carrying *alkB* genes had higher degradation abilities of long-chain *c*-alkane than those of strains carrying no *alkB* genes (Table 3 and Table 5). However, *R. rhodnii* NBRC100604 carrying no *alkB* gene showed a degradation rate of above 30%. The ability of long-chain *c*-alkane degradation might be regulated

Table 3. The degradation of long-chain *c*-alkanes by stains harboring *alkB* [*alkB* R2 type] in the genus *Rhodococcus*.

Strain	Degradation Ratio (%)*		
	Undecylcyclohexane (UDC)	Dodecylcyclohexane (DDC)	Tridecylcyclohexane (TDC)
<i>R. erythropolis</i> NDKK1	16.0 ± 7.5	31.7 ± 8.7	17.9 ± 7.3
<i>R. erythropolis</i> NDKK2	6.7 ± 10.7	42.9 ± 10.2	23.5 ± 3.2
<i>R. erythropolis</i> NDKK5	66.7 ± 0.2	42.2 ± 1.8	40.6 ± 10.0
<i>R. erythropolis</i> NDKK6	38.8 ± 1.0	18.9 ± 6.2	75.7 ± 6.1
<i>R. erythropolis</i> NDKK7	53.5 ± 3.6	42.1 ± 4.0	46.7 ± 9.1
<i>R. erythropolis</i> NDKK48	40.4 ± 2.5	33.1 ± 12.6	81.4 ± 0.1
<i>R. erythropolis</i> ODNM1C	78.2 ± 2.4	41.8 ± 2.9	91.0 ± 1.1
<i>R. erythropolis</i> NDKY82A	52.7 ± 4.9	39.1 ± 9.2	78.4 ± 1.5
<i>R. erythropolis</i> ODMI54	68.2 ± 5.0	37.8 ± 8.2	46.8 ± 2.6
<i>R. erythropolis</i> ODNM2B	98.9 ± 0.2	30.3 ± 3.5	94.6 ± 6.7
<i>R. erythropolis</i> NDMI144	91.4 ± 3.9	27.7 ± 10.3	39.9 ± 8.3
<i>R. erythropolis</i> NBRC15567	64.5 ± 9.4	0.0 ± 1.5	75.4 ± 1.5
<i>R. erythropolis</i> NBRC15564	50.6 ± 3.9	41.6 ± 9.1	39.6 ± 7.1
<i>R. baikonurensis</i> NBRC100611	50.4 ± 3.8	16.0 ± 3.0	20.6 ± 5.7
Average	55.5	31.8	55.2

*n = 3.






Figure 3. Gas chromatograms of UDC degraded by *R. erythropolis* NDKK6. Arrows indicate the peak of UDC. Images of gas chromatograms of LB medium with UDC substrate cultivated without (a) and with (b) *R. erythropolis* NDKK6.

Table 4. The degradation of long-chain *c*-alkanes by stains harboring *alkB* genes [non-*alkB* R2 type] in the genus *Rhodococcus*.

Strain	Type of <i>alkB</i> Gene	Degradation Ratio (%)*		
		Undecylcyclohexane (UDC)	Dodecylcyclohexane (UDC)	Tridecylcyclo-hexane (UDC)
<i>R. wratislaviensis</i> NBRC100605	<i>alkB</i> RW	39.7 ± 5.2	58.6 ± 4.0	25.1 ± 2.1
<i>R. opacus</i> NBRC100624	<i>alkB</i> ROP	94.9 ± 1.4	56.8 ± 5.8	43.1 ± 0.4
<i>R. percoletus</i> NBRC100626	<i>alkB</i> ROP	47.1 ± 13.5	35.7 ± 2.3	60.6 ± 11.1
<i>R. ruber</i> NBRC15591	<i>alkB</i> RR	98.2 ± 1.6	55.9 ± 3.6	43.2 ± 3.4
<i>R. equi</i> NBRC101255	<i>alkB</i> RE	48.9 ± 2.1	39.3 ± 3.4	98.4 ± 1.6
<i>R. jostii</i> NBRC16295	<i>alkB</i> RJ	29.8 ± 4.9	27.8 ± 10.7	65.0 ± 8.2
<i>R. triatoma</i> NBRC103116	<i>alkB</i> R1	34.0 ± 14.2	15.9 ± 7.4	75.3 ± 1.7
<i>R. koreensis</i> NBRC100607	<i>alkB</i> RK	6.0 ± 10.6	4.2 ± 5.4	5.5 ± 2.3
<i>R. corynebacterioides</i> NBRC14404	<i>alkB</i> RC	41.4 ± 6.5	0.0 ± 10.2	62.3 ± 0.6
<i>R. zopfii</i> NBRC100606	<i>alkB</i> RZ	53.9 ± 9.1	0.0 ± 9.7	60.3 ± 5.3
Average		49.4	29.4	53.9

*n = 3.

Table 5. The degradation of long-chain *c*-alkanes by stains harboring no *alkB* genes in the genus *Rhodococcus*.

Strain	Degradation Ratio (%)*		
	Undecylcyclohexane (UDC)	Dodecylcyclohexane (UDC)	Tridecylcyclo-hexane (UDC)
<i>R. tukisamuensis</i> NBRC100609	26.0 ± 1.4	4.1 ± 11.0	20.1 ± 8.0
<i>R. maanshanensis</i> NBRC100610	18.3 ± 7.4	13.4 ± 4.2	18.6 ± 3.4
<i>R. pyridinivorans</i> NBRC100608	2.0 ± 1.4	11.2 ± 13.1	19.6 ± 10.2
<i>R. kroppenstedtii</i> NBRC103113	12.3 ± 5.2	1.7 ± 1.0	15.9 ± 4.2
<i>R. rhodnii</i> NBRC100604	34.7 ± 11.5	34.2 ± 3.8	34.8 ± 3.3
<i>R. coprophilus</i> NBRC100603	22.2 ± 3.3	15.4 ± 6.7	5.6 ± 7.1
Average	19.3	13.3	19.1

*n = 3.

by the alkane hydroxylase gene, while genus *Rhodococcus* spp. carrying no *alkB* genes may degrade long-chain *c*-alkanes by other degradation systems such as those involving the cytochrome gene [41].

3.5. Alkane Hydroxylase Gene-Carrying Bacteria and Their Roles in Material Circulation in Nature

c-alkane-degrading bacteria belonging to the genus *Rhodococcus* isolated from their natural habitats in our previous study were closely related to *R. erythropolis* and strains harboring *alkB* R2 type [33]. Although many *c*-alkane-degrading bacteria of the genus *Rhodococcus* are widely distributed, long-chain *c*-alkanes are limited in the natural environment. These bacteria may utilize hydrocarbons such as the wax of leaves and aromatic compounds, which include linear or cyclic hydrocarbon components. Thus, these bacteria harboring alkane hydroxylase genes might contribute to the material circulation of several kinds of hydrocarbon compounds in nature [42].

4. Conclusion

To characterize the long-chain cyclic alkane (*c*-alkane) degradation of bacteria in *Rhodococcus*, 11 strains were isolated from nature using long-chain *c*-alkane as a substrate. These strains were identified as *R. erythropolis*, and all were shown to carry the *alkB* R2 type. It was also confirmed that the degradation ratios of long-chain *c*-alkanes by *Rhodococcus* carrying the *alkB* R2 type of gene (55.5% of UDC, 31.8% of DDC, and 55.2% of TDC) were higher than in those carrying the non-*alkB* R2 type of gene (49.4% of UDC, 29.4% of DDC, and 53.9% of TDC) and no *alkB* gene (19.3% of UDC, 13.3% of DDC, and 19.1% of TDC).

Conflicts of Interest

The authors declare no conflicts of interest regarding the publication of this paper.

References

- [1] Usman, M., Faure, P., Hanna, K., Abdelmoula, M. and Ruby, C. (2012) Application of Magnetite Catalyzed Chemical Oxidation (Fenton-Like and Persulfate) for the Remediation of Oil Hydrocarbon Contamination. *Fuel*, **96**, 270-276. <https://doi.org/10.1016/j.fuel.2012.01.017>
- [2] Hatayama, K., Sakihama, Y., Matsumiya, Y. and Kubo, M. (2008) Construction of an Efficient Bioremediation System for Petroleum Hydrocarbon-Contaminated Soils Using Specific Hydrocarbon-Degrading Bacteria and Bacterial Biomass Monitoring. In: Domínguez, J.B., Ed., *Soil Contamination Research Trends*, Nova Science Publishers, NY, 143-160.
- [3] Dua, M., Johri, A.K., Singh, A. and Sethunathan, N. (2002) Biotechnology and Bioremediation: Successes and Limitations. *Applied Microbiology and Biotechnology*, **59**, 143-152. <https://doi.org/10.1007/s00253-002-1024-6>
- [4] Vidali, M. (2001) Bioremediation. An Overview. *Pure and Applied Chemistry*, **73**, 1163-1172. <https://doi.org/10.1351/pac200173071163>

- [5] Stroud, J.L., Paton, G.I. and Semple, K.T. (2007) Microbe-Aliphatic Hydrocarbon Interactions in Soil: Implications for Biodegradation and Bioremediation. *Journal of Applied Microbiology*, **102**, 1239-1253. <https://doi.org/10.1111/j.1365-2672.2007.03401.x>
- [6] Kissin, Y.V. (1990) Catagenesis of Light Cycloalkanes in Petroleum. *Organic Geochemistry*, **15**, 575-594. [https://doi.org/10.1016/0146-6380\(90\)90103-7](https://doi.org/10.1016/0146-6380(90)90103-7)
- [7] Korda, A., Santas, P., Tenente, A. and Santas, R. (1997) Petroleum Hydrocarbon Bioremediation: Sampling Analytical Techniques, *in Situ* Treatment and Commercial Microorganisms Currently Used. *Applied Microbiology and Biotechnology*, **48**, 677-686. <https://doi.org/10.1007/s002530051115>
- [8] Pritchard, P.H., Mueller, J.G., Rogers, J.C., Kremer, F.V. and Glaser, J.A. (1992) Oil Spill Bioremediation: Experiences, Lessons and Results from the *Exxon Valdez* Oil Spill in Alaska. *Biodegradation*, **3**, 315-335. <https://doi.org/10.1007/BF00129091>
- [9] Adhikari, D., Perwira, I.Y., Araki, K.S. and Kubo, M. (2016) Stimulation of Soil Microorganisms in Pesticide-Contaminated Soil Using Organic Materials. *AIMS Bioengineering*, **3**, 379-388. <https://doi.org/10.3934/bioeng.2016.3.379>
- [10] Blakley, E.R. (1978) The Microbial Degradation of Cyclohexanecarboxylic Acid by a β -Oxidation Pathway with Simultaneous Induction to the Utilization of Benzoate. *Canadian Journal of Microbiology*, **24**, 847-855. <https://doi.org/10.1139/m78-141>
- [11] Blakley, E.R. and Papish, B. (1982) The Metabolism of Cyclohexanecarboxylic Acid and 3-Cyclohexenecarboxylic Acid by *Pseudomonas putida*. *Canadian Journal of Microbiology*, **28**, 1324-1329. <https://doi.org/10.1139/m82-198>
- [12] Komukai-Nakamura, S., Sugiura, K., Yamauchi-Inomata, Y., Toki, H., Venkateswaran, K., Yamamoto, S., Tanaka, H. and Harayama, S. (1996) Construction of Bacterial Consortia That Degrade Arabian Light Crude Oil. *Journal of Fermentation and Bioengineering*, **82**, 570-574. [https://doi.org/10.1016/S0922-338X\(97\)81254-8](https://doi.org/10.1016/S0922-338X(97)81254-8)
- [13] Aislabie, J., Saul, D.J. and Foght, J.M. (2006) Bioremediation of Hydrocarbon-Contaminated Polar Soils. *Extremophiles*, **10**, 171-179. <https://doi.org/10.1007/s00792-005-0498-4>
- [14] Zampolli, J., Collina, E., Lasagni, M. and Gennaro, P.G. (2014) Biodegradation of Variable-Chain-Length *n*-alkanes in *Rhodococcus opacus* R7 and the Involvement of an Alkane Hydroxylase System in the Metabolism. *AMB Express*, **4**, 73. <https://doi.org/10.1186/s13568-014-0073-4>
- [15] Perry, J.J. (1979) Microbial Cooxidations Involving Hydrocarbons. *Microbiological Reviews*, **43**, 59-72.
- [16] Throne-Holst, M., Wentzel, A., Ellingsen, T.E., Kotlar, H.K. and Zotchev, S.B. (2007) Identification of Novel Genes Involved in Long-Chain *n*-Alkane Degradation by *Acinetobacter* sp. Strain DSM 17874. *Applied and Environmental Microbiology*, **73**, 3327-3332. <https://doi.org/10.1128/AEM.00064-07>
- [17] Lal, B. and Khanna, S. (1996) Degradation of Crude Oil by *Acinetobacter calcoaceticus* and *Alcaligenes odorans*. *Journal of Applied Microbiology*, **81**, 355-362. <https://doi.org/10.1111/j.1365-2672.1996.tb01926.x>
- [18] Sakai, Y., Maeng, J.H., Tani, Y. and Kato, N. (1994) Use of Long-Chain *n*-alkanes (C13-C44) by an Isolate, *Acinetobacter* sp. M-1. *Bioscience, Biotechnology and Biochemistry*, **58**, 2128-2130. <https://doi.org/10.1271/bbb.58.2128>
- [19] Táncsics, A., Benedek, T., Szoboszlai, S., Veres, P.G., Farkas, M., Máthé, I., Mária-ligeti, K., Kukolya, J., Lányi, S. and Kriszt, B. (2015) The Detection and Phylogenetic Analysis of the Alkane 1-Monooxygenase Gene of Members of the Genus *Rhodococcus*. *Systematic and Applied Microbiology*, **38**, 1-7.

- <https://doi.org/10.1016/j.syapm.2014.10.010>
- [20] Kubota, K., Koma, D., Matsumiya, Y., Chung, S.Y. and Kubo, M. (2008) Phylogenetic Analysis of Long-Chain Hydrocarbon-Degrading Bacteria and Evaluation of Their Hydrocarbon-Degradation by the 2,6-DCPIP Assay. *Biodegradation*, **19**, 749-757. <https://doi.org/10.1007/s10532-008-9179-1>
- [21] Koma, D., Sakashita, Y., Kubota, K., Fujii, Y., Hasumi, F., Chung, S.Y. and Kubo, M. (2004) Degradation Pathways of Cyclic Alkanes in *Rhodococcus* sp. NDKK48. *Applied Microbiology and Biotechnology*, **66**, 92-99. <https://doi.org/10.1007/s00253-004-1623-5>
- [22] Kostichka, K., Thomas, S.M., Gibson, K.J., Nagarajan, V. and Cheng, Q. (2001) Cloning and Characterization of a Gene Cluster for Cyclododecanone Oxidation in *Rhodococcus ruber* SC1. *Journal of Bacteriology*, **183**, 6478-6486. <https://doi.org/10.1128/JB.183.21.6478-6486.2001>
- [23] van Hamme, J.D., Odumeru, J.A. and Ward, O.P. (2000) Community Dynamics of a Mixed-Bacterial Culture Growing on Petroleum Hydrocarbons in Batch Culture. *Canadian Journal of Microbiology*, **46**, 441-450. <https://doi.org/10.1139/w00-013>
- [24] van Beilen, J.B., Smits, T.H.M., Roos, F.F., Brunner, T., Balada, S.B., Rothlisberger, M. and Witholt, B. (2005) Identification of an Aminic Acid Position That Determines the Substrate Range of Integral Membrane Alkane Hydroxylases. *Journal of Bacteriology*, **187**, 85-91. <https://doi.org/10.1128/JB.187.1.85-91.2005>
- [25] van Beilen, J.B. and Funhoff, E.G. (2007) Alkane Hydroxylases Involved in Microbial Alkane Degradation. *Applied Microbiology and Biotechnology*, **74**, 13-21. <https://doi.org/10.1007/s00253-006-0748-0>
- [26] Whyte, L.G., Smits, T.H.M., Labbé, D., Witholt, B., Greer, C.W. and van Beilen, J.B. (2002) Gene Cloning and Characterization of Multiple Alkane Hydroxylase Systems in *Rhodococcus* Strains Q15 and NRRL B-16531. *Applied and Environmental Microbiology*, **68**, 5933-5942. <https://doi.org/10.1128/AEM.68.12.5933-5942.2002>
- [27] Fujii, T., Narikawa, T., Takeda, K. and Kato, J. (2004) Biotransformation of Various Alkanes Using the *Escherichia coli* Expressing an Alkane Hydroxylase System from *Gordonia* sp. TF6. *Bioscience, Biotechnology and Biochemistry*, **68**, 2171-2177. <https://doi.org/10.1271/bbb.68.2171>
- [28] Vomberg, A. and Klinner, U. (2000) Distribution of *alkB* Genes within *n*-Alkane-Degrading Bacteria. *Journal of Applied Microbiology*, **89**, 339-348. <https://doi.org/10.1046/j.1365-2672.2000.01121.x>
- [29] Whyte, L.G., Hawari, J., Zhou, E., Bourbonnière, L., Inniss, W. and Greer, C.W. (1998) Biodegradation of Variable-Chain-Length Alkanes at Low Temperatures by a Psychrotrophic *Rhodococcus* sp. *Applied and Environmental Microbiology*, **64**, 2578-2584.
- [30] van Beilen, J.B., Smits, T.H.M., Whyte, L.G., Schorcht, S., Röthlisberger, M., Plagemeier, T., Engesser, H. and Witholt, B. (2002) Alkane Hydroxylase Homologues in Gram-Positive Strains. *Environmental Microbiology*, **4**, 676-682. <https://doi.org/10.1046/j.1462-2920.2002.00355.x>
- [31] Hara, A., Baik, S.H., Syutsubo, K., Misawa, N., Smits, T.H.M., van Beilen, J.B. and Harayama, S. (2004) Cloning and Functional Analysis of *alkB* Genes in *Alcanivorax borkumensis* SK2. *Environmental Microbiology*, **6**, 191-197. <https://doi.org/10.1046/j.1462-2920.2003.00550.x>
- [32] Rehm, H.J. and Reiff, I. (1981) Mechanisms and Occurrence of Microbial Oxidation of Long-Chain Alkanes. In: Fiechter, A., Ed., *Reactors and Reactions, Vol. 19: Advances in Biochemical Engineering*, Springer, Berlin, 175-215.

https://doi.org/10.1007/3-540-10464-X_18

- [33] Fukuhara, Y., Horii, S., Matsuno, T., Matsumiya, Y., Mukai, M. and Kubo, M. (2013) Distribution of Hydrocarbon-Degrading Bacteria in the Soil Environment and Their Contribution to Bioremediation. *Applied Biochemistry and Biotechnology*, **170**, 329-339. <https://doi.org/10.1007/s12010-013-0170-x>
- [34] Perry, J.J. (1984) Microbial Metabolism of Cyclic Alkanes. In: Atlas, R.M., Ed., *Petroleum Microbiology*, Macmillan, 61-98.
- [35] Koma, D., Sakashita, Y., Kubota, K., Fujii, Y., Hasumi, F., Chung, S.Y. and Kubo, M. (2003) Degradation of Car Engine Base Oil by *Rhodococcus* sp. NDKK48 and *Gordonia* sp. NDKY76A. *Bioscience, Biotechnology and Biochemistry*, **67**, 1590-1593. <https://doi.org/10.1271/bbb.67.1590>
- [36] Koma, D., Hasumi, F., Yamamoto, E., Ohta, T., Chung, S.Y. and Kubo, M. (2001) Biodegradation of Long-Chain *n*-Paraffins from Waste Oil of Car Engine by *Acinetobacter* sp. *Journal of Bioscience and Bioengineering*, **91**, 94-96. [https://doi.org/10.1016/S1389-1723\(01\)80120-1](https://doi.org/10.1016/S1389-1723(01)80120-1)
- [37] Koma, D., Hasumi, F., Chung, S.Y. and Kubo, M. (2003) Biodegradation of *n*-Alkylcyclohexanes by Co-Oxidation via Multiple Pathways in *Acinetobacter* sp. ODDK71. *Journal of Bioscience and Bioengineering*, **95**, 641-644. [https://doi.org/10.1016/S1389-1723\(03\)80178-0](https://doi.org/10.1016/S1389-1723(03)80178-0)
- [38] Iwamoto, T. and Nasu, M. (2001) Current Bioremediation Practice and Perspective. *Journal of Bioscience and Bioengineering*, **92**, 1-8. [https://doi.org/10.1016/S1389-1723\(01\)80190-0](https://doi.org/10.1016/S1389-1723(01)80190-0)
- [39] Sanpa, S., Sumiyoshi, S., Kujira, T., Matsumiya, Y. and Kubo, M. (2006) Isolation and Characterization of a Bluegill Degrading Microorganism, and Analysis of the Root Hair-Promoting Effect of the Degraded Products. *Bioscience, Biotechnology and Biochemistry*, **70**, 340-347. <https://doi.org/10.1271/bbb.70.340>
- [40] van Beilen, J.B., Li, Z., Duetz, W.A. and Witholt, B. (2003) Diversity of Alkane Hydroxylase Systems in the Environment. *Oil Gas Science and Technology*, **58**, 427-440. <https://doi.org/10.2516/ogst:2003026>
- [41] van Beilen, J.B., Funhoff, E.G., van Loon, A., Just, A., Kaysser, L., Bouza, M., Holtackers, R., Röthlisberger, M., Li, Z. and Witholt, B. (2006) Cytochrome P450 Alkane Hydroxylases of the CYP153 Family Are Common in Alkane-Degrading Eubacteria Lacking Integral Membrane Alkane Hydroxylases. *Applied and Environmental Microbiology*, **72**, 59-65. <https://doi.org/10.1128/AEM.72.1.59-65.2006>
- [42] Nie, Y., Chi, C.Q., Fang, H., Liang, J.L., Lu, S.L., Lai, G.L., Tang, Y.Q. and Wu, X.L. (2014) Diverse Alkane Hydroxylase Genes in Microorganisms and Environments. *Scientific Reports*, **4**, Article No. 4968. <https://doi.org/10.1038/srep04968>