

Exact Quasi-Classical Asymptotic beyond Maslov Canonical Operator and Quantum Jumps Nature

Jaykov Foukzon¹, Alex Potapov², Stanislav Podosenov³

¹Israel Institute of Technology, Haifa, Israel

²The Institute of Radioengineering and Electronics (IRE) of Russian Academy of Sciences, Moscow, Russian

³All-Russian Scientific-Research Institute, Moscow, Russian

Email: jaykovfoukzon@list.ru

Received 12 February 2015; accepted 26 May 2015; published 29 May 2015

Copyright © 2015 by authors and Scientific Research Publishing Inc.

This work is licensed under the Creative Commons Attribution International License (CC BY).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

Abstract

Exact quasi-classical asymptotic beyond WKB-theory and beyond Maslov canonical operator to the Colombeau solutions of the n -dimensional Schrodinger equation is presented. Quantum jumps nature is considered successfully. We pointed out that an explanation of quantum jumps can be found to result from Colombeau solutions of the Schrödinger equation alone without additional postulates.

Keywords

Quantum Jumps, Quantum Measurements Theory, Quantum Averages, Limiting Quantum Trajectory, Schrodinger Equation, Stochastic Quantum Jump Equation, Colombeau Solution, Feynman Path Integral, Maslov Canonical Operator, Feynman-Colombeau Propagator

1. Introduction

A number of experiments on trapped single ions or atoms have been performed in recent years [1]-[4]. Monitoring the intensity of scattered laser light off of such systems has shown abrupt changes that have been cited as evidence of “quantum jumps” between states of the scattered ion or atom. The existence of such jumps was required by Bohr in his theory of the atom. Bohr’s quantum jumps between atomic states [5] were the first form of quantum dynamics to be postulated. He assumed that an atom remained in an atomic eigenstate until it made an instantaneous jump to another state with the emission or absorption of a photon. Since these jumps do not appear

to occur in solutions of the Schrodinger equation, something similar to Bohr’s idea has been added as an extra postulate in modern quantum mechanics.

Stochastic quantum jump equations [6]-[8] were introduced as a tool for simulating the dynamics of a dissipative system with a large Hilbert space and their links with quantum measurement theory were also noted [9]-[13]. This measurement interpretation is generally known as quantum trajectory theory [14]. By adding filter cavities as part of the apparatus, even the quantum jumps in the dressed state model can be interpreted as approximations to measurement-induced jumps [15].

The question arises whether an explanation of these jumps can be found to result from a Colombeau solution [16]-[18] $(\Psi_\varepsilon(x, t; \hbar))_\varepsilon$ of the Schrödinger equation alone without additional postulates. We found exact quasi-classical asymptotic of the quantum averages with position variable with localized initial data.

$$(i, t, x_0; \hbar, \varepsilon) = \left(\int x_i |\Psi_\varepsilon(x, t, x_0; \hbar)|^2 dx \right)_\varepsilon, \quad \varepsilon \in (0, 1], \quad x, x_0 \in \mathbb{R}^d, \quad i = 1, \dots, d \tag{1.1}$$

i.e. we found the limiting Colombeau quantum averages (limiting Colombeau quantum trajectories) such that [18]:

$$(i, t, x_0; \varepsilon) = (\varinjlim_{\hbar \rightarrow 0} i, t, x_0; \hbar, \varepsilon) = \left(\varinjlim_{\hbar \rightarrow 0} \int x_i |\Psi_\varepsilon(x, t, x_0; \hbar)|^2 dx \right)_\varepsilon, \tag{1.2}$$

$$\varepsilon \in (0, 1], \quad x \in \mathbb{R}^d, \quad i = 1, \dots, d, \quad t \in [0, T]$$

and limiting quantum trajectories $q(t, x_0) = \{q_1(t, x_0), \dots, q_d(t, x_0)\} \in \mathbb{R}^d, \quad t \in [0, T]$ such that

$$q_i(t, x_0) = \langle i, t, x_0 \rangle = \lim_{\varepsilon \rightarrow 0} \varinjlim_{\hbar \rightarrow 0} \langle i, t, x_0; \hbar, \varepsilon \rangle \tag{1.3}$$

$$= \lim_{\varepsilon \rightarrow 0} \varinjlim_{\hbar \rightarrow 0} \int x_i |\Psi_\varepsilon(x, t, x_0; \hbar)|^2 dx, \quad x \in \mathbb{R}^d, \quad i = 1, \dots, d.$$

if limit in LHS of Equation (1.3) exists.

The physical interpretation of these asymptotic given below, shows that the answer is “yes” for the limiting quantum trajectories with localized initial data.

Note that an axiom of quantum measurement is: if the particle is in some state $|\Psi_{t,\varepsilon}\rangle$ that the probability $P(x, \delta x)$ of getting a result $x \in \mathbb{R}^d$ at instant t with an accuracy of $\|\delta x\| \ll 1$ will be given by

$$P_t(x, \delta x, x_0; \hbar, \varepsilon) = \int_{\|x-x'\| \leq \|\delta x\|} \left| \langle x' | \Psi_{t,\varepsilon} \rangle \right|^2 dx' = \int_{\|x-x'\| \leq \|\delta x\|} |\Psi_\varepsilon(x', t, x_0; \hbar)|^2 dx'. \tag{1.4}$$

We rewrite now Equation (1.4) of the form

$$P_t(x, \delta x, x_0; \hbar, \varepsilon) = \frac{1}{(2\pi)^{d/2} \delta x^d} \int_{\mathbb{R}^d} |\Psi_\varepsilon(x', t, x_0; \hbar)|^2 \exp\left[-\frac{\|x-x'\|^2}{\|\delta x\|^2}\right] dx' \tag{1.5}$$

We define well localized limiting quantum trajectories $q(t) = q(t, x_0) = \{q_1(t, x_0), \dots, q_d(t, x_0)\} \in \mathbb{R}^d, \quad q(0, x_0) = x_0, \quad t \in [0, T]$ such that:

$$\left(\varinjlim_{\|\delta q(t)\| \rightarrow 0} \varinjlim_{\hbar \rightarrow 0} P_t(q(t), \delta q(t), x_0; \hbar, \varepsilon) \right)_\varepsilon = 1 \tag{1.6}$$

and well localized limiting quantum trajectories $q(t) = q(t, x_0) = \{q_1(t, x_0), \dots, q_d(t, x_0)\} \in \mathbb{R}^d, \quad q(0, x_0) = x_0, \quad t \in [0, T]$ such that:

$$\lim_{\varepsilon \rightarrow 0} \varinjlim_{\|\delta q(t)\| \rightarrow 0} \varinjlim_{\hbar \rightarrow 0} P_t(q(t), \delta q(t), x_0; \hbar, \varepsilon) = 1 \tag{1.7}$$

if limit in LHS of Equation (1.7) exists.

2. Colombeau Solutions of the Schrödinger Equation and Corresponding Path Integral Representation

Let \mathbf{H} be a complex infinite dimensional separable Hilbert space, with inner product $\langle \cdot, \cdot \rangle$ and norm $\|\cdot\|$.

Let us consider Schrödinger equation:

$$-i\hbar\left(\frac{\partial\Psi(t)}{\partial t}\right) + \hat{H}(t)\Psi(t) = 0, \quad \Psi(0) = \Psi_0(x), \tag{2.1}$$

$$H(t) = -\left(\frac{\hbar^2}{2m}\right)\Delta + V(x,t). \tag{2.2}$$

Here operator $H(t): \mathbb{R} \times \mathbf{H} \rightarrow \mathbf{H}$ is essentially self-adjoint, $\hat{H}(t)$ is the closure of $H(t)$.

Theorem 2.1. [19] [20]. Assume that: (1) $\Psi_0(x) \in L_2(\mathbb{R}^d)$, (2) $V(x,t)$ is continuous and $\sup_{x \in \mathbb{R}^d, t \in [0,T]} |V(x,t)| < +\infty$. Then corresponding solution of the Schrödinger Equations (2.1)-(2.2) exist and can be represented via formulae

$$\Psi(t,x) = \lim_{n \rightarrow \infty} \left(\frac{nm}{4\pi i t \hbar}\right)^{d(n+1)/2} \int_{\mathbb{R}^d} \dots \int_{\mathbb{R}^d} dx_0 dx_1 \dots dx_n \Psi_0(x_0) \exp\left[\frac{i}{\hbar} S(x_0, x_1, \dots, x_n, x_{n+1}; t)\right], \tag{2.3}$$

where we have set $x_{n+1} = x$ and

$$S(x_0, x_1, \dots, x_n, x_{n+1}; t) = \sum_{i=1}^n \left[\frac{m |x_{i+1} - x_i|^2}{4(t/n)^2} - V(x_{i+1}, t_i) \right], \tag{2.4}$$

where $t_i = \frac{it}{n}$, Let $q_n(t)$ be a trajectory; that is, a function from $[0,t]$ to \mathbb{R}^d with $q_n(0) = x_0$ and set $q_n(t_i) = x_i, i = 1, \dots, n+1$. We rewrite Equation (2.3) for a future application symbolically for short of the following form

$$\Psi(t,x) = \lim_{n \rightarrow \infty} \int_{q_n(t)=x} D[q_n(t)] \Psi_0(q_n(0)) \exp\left[\frac{i}{\hbar} S(q_n(t), x; t)\right], \tag{2.5}$$

where we have set 1) $S(q_n(t), x; t) = S(x_0, x_1, \dots, x_n, x_{n+1}; t)$ and 2) $D[q_n(t)]$ that is, a

$$D[q_n(t)] = \left(\frac{nm}{4\pi i t \hbar}\right)^{d(n+1)/2} \prod_{j=0}^n dx_j, \tag{2.6}$$

Trotter and Kato well known classical results give a precise meaning to the Feynman integral when the potential $V(x,t)$ is sufficiently regular [18] [19]. However if potential $V(x,t)$ is a non-regular this is well known problem to represent solution of the Schrödinger Equations (2.1)-(2.2) via formulae (2.3), see [19].

We avoided this difficulty using contemporary Colombeau framework [16]-[18]. Using replacement

$x_i \rightarrow \frac{x_i}{1 + \varepsilon^{2k} |x_i|^{2k}}, \varepsilon \in (0,1], k \geq 1$ we obtain from potential $V(x,t)$ regularized

Potential $V_\varepsilon(x,t), \varepsilon \in (0,1]$ such that $V_{\varepsilon=0}(x,t) = V(x,t)$ and

1) $(V_\varepsilon(x,t))_\varepsilon \in G(\mathbb{R}^d)$,

$$\sup_{x \in \mathbb{R}^d, t \in [0,T]} |V_\varepsilon(x,t)| < +\infty, \quad \varepsilon \in (0,1]. \tag{2.7}$$

Here $G(\mathbb{R}^d)$ is Colombeau algebra of Colombeau generalized functions [16]-[18].

Finally we obtain regularized Schrödinger equation of Colombeau form [21]-[24]:

$$-i\hbar\left(\frac{\partial\Psi_\varepsilon(t)}{\partial t}\right)_\varepsilon + (\hat{H}_\varepsilon(t)\Psi_\varepsilon(t))_\varepsilon = 0, \quad (\Psi_\varepsilon(0))_\varepsilon = \Psi_0(x), \tag{2.8}$$

$$H_\varepsilon(t) = -\left(\frac{\hbar^2}{2m}\right)\Delta + V_\varepsilon(x,t). \tag{2.9}$$

Using the inequality (2.7) Theorem 2.1 asserts again that corresponding solution of the Schrödinger Equations (2.8)-(2.9) exist and can be represented via formulae:

$$(\Psi_\varepsilon(t, x))_\varepsilon = \left(\lim_{n \rightarrow \infty} \left(\frac{nm}{4\pi i t \hbar} \right)^{d(n+1)/2} \int_{\mathbb{R}^d} \cdots \int_{\mathbb{R}^d} dx_0 dx_1 \cdots dx_n \Psi_0(x_0) \exp \left[\frac{i}{\hbar} S_\varepsilon(x_0, x_1, \dots, x_n, x_{n+1}; t) \right] \right)_\varepsilon, \quad (2.10)$$

where we have set $x_{n+1} = x$ and

$$S_\varepsilon(x_0, x_1, \dots, x_n, x_{n+1}; t) = \sum_{i=1}^n \left[\frac{m |x_{i+1} - x_i|^2}{4 (t/n)^2} - V_\varepsilon(x_{i+1}, t_i) \right], \quad (2.11)$$

where we have set $t_i = \frac{it}{n}$.

We rewrite Equation (2.10) for a future application symbolically of the following form

$$(\Psi_\varepsilon(t, x))_\varepsilon = \left(\lim_{n \rightarrow \infty} \int_{q_n(t)=x} D[q_n(t)] \Psi_0(q_n(0)) \exp \left[\frac{i}{\hbar} S_\varepsilon(q_n(t); t) \right] \right)_\varepsilon, \quad (2.12)$$

or of the following form

$$(\Psi_\varepsilon(t, x))_\varepsilon = \left(\lim_{n \rightarrow \infty} \Psi_{\varepsilon, n}(t, x) \right)_\varepsilon = \left(\lim_{n \rightarrow \infty} \int_{q(t)=x} D_n[q(t)] \Psi_0(q_n(0)) \exp \left[\frac{i}{\hbar} S_\varepsilon(\dot{q}(t), q(t); t) \right] \right)_\varepsilon. \quad (2.13)$$

For the limit in RHS of (2.12) and (2.13) we will be used canonical path integral notation

$$(\Psi_\varepsilon(t, x))_\varepsilon = \left(\int_{q(t)=x} D[q(t)] \Psi_0(q(0)) \exp \left[\frac{i}{\hbar} S_\varepsilon(\dot{q}(t), q(t)) \right] \right)_\varepsilon, \quad (2.14)$$

where $S_\varepsilon(\dot{q}(t), q(t)) = \int_0^t \left[\frac{m}{4} \dot{q}^2(s) - V_\varepsilon(q(s), s) \right] ds$.

Substitution $n = 8k + 7$ into RHS of the Equation (2.10) gives

$$(\Psi_\varepsilon(t, x))_\varepsilon = \left(\lim_{k \rightarrow \infty} \left(\frac{(8k+7)m}{4\pi t \hbar} \right)^{d(4k+4)} \int_{\mathbb{R}^d} \cdots \int_{\mathbb{R}^d} dx_0 dx_1 \cdots dx_{8k+7} \Psi_0(x_0) \exp \left[\frac{i}{\hbar} S_\varepsilon(x_0, x_1, \dots, x_{8k+7}, x_{8k+8}; t) \right] \right)_\varepsilon. \quad (2.15)$$

We rewrite Equation (2.15) for a future application symbolically of the following form

$$(\Psi_\varepsilon(t, x))_\varepsilon = \left(\lim_{n \rightarrow \infty} \int_{q_n(t)=x} D^+[q_n(t)] \Psi_0(q_n(0)) \exp \left[\frac{i}{\hbar} S_\varepsilon(q_n(t); t) \right] \right)_\varepsilon, \quad (2.16)$$

or of the following form

$$(\Psi_\varepsilon(t, x))_\varepsilon = \left(\lim_{n \rightarrow \infty} \int_{q(t)=x} D_n^+[q(t)] \Psi_0(q(0)) \exp \left[\frac{i}{\hbar} S_\varepsilon(\dot{q}(t), q(t); t) \right] \right)_\varepsilon. \quad (2.17)$$

For the limit in RHS of (2.16) and (2.17) we will be used following path integral notation

$$(\Psi_\varepsilon(t, x))_\varepsilon = \left(\int_{q(t)=x} D^+[q(t)] \Psi_0(q(0)) \exp \left[\frac{i}{\hbar} S_\varepsilon(\dot{q}(t), q(t)) \right] \right)_\varepsilon. \quad (2.18)$$

Let us consider now regularized oscillatory integral

$$(J_{\varepsilon, n}(t; \hbar))_\varepsilon = \left(\int_{\mathbb{R}^d} \cdots \int_{\mathbb{R}^d} dx_0 dx_1 \cdots dx_n f(x_0, x_1, \dots, x_n) \Psi_0(x_0) \exp \left[\frac{i}{\hbar} S_\varepsilon(x_0, x_1, \dots, x_n; t) \right] \right)_\varepsilon. \quad (2.19)$$

Lemma 2.1. (Localization Principle [25] [26]) Let Ω be a domain in $\mathbb{R}^{d \times n}$ and $f \in C_0^\infty(\Omega)$ be a smooth function of compact support, $S_\varepsilon \in C^\infty(\Omega)$, $\varepsilon \in (0, 1]$ be a real valued smooth function without stationary points in $\text{supp}(f)$, i.e. $\partial_x S_\varepsilon(x) \neq 0$ for $x \in \Omega$. Let L be a differential operator

$$L(f) = -\sum_{i=1}^n \frac{\partial}{\partial x_i} \left(|S'_{\varepsilon, x}(x)|^{-2} \frac{\partial S_\varepsilon}{\partial x_i} f \right). \quad (2.20)$$

Then $(|J_{\varepsilon,n}(t; \hbar)|)_{\varepsilon} \leq \hbar \int_{\Omega} L(f(x)) dx$. $\forall n, m \in \mathbb{N}$, $\forall \hbar \leq 1$ there exist c_m such that

$$(J_{\varepsilon,n}(t; \hbar))_{\varepsilon} \leq c_m \hbar^m \|f\|, \quad \|f\| = \sup_{x \in \Omega} \sum_{|a| \leq m} |D^a f|. \quad (2.21)$$

Lemma 2.2. (Generalized Localization Principle) Let Ω_n be a domain in \mathbb{R}^{dx_n} and $f_n \in C_0^\infty(\Omega_n)$ be a real valued smooth function without stationary points in $\text{supp}(f)$, i.e. $\partial_x S_\varepsilon(x) \neq 0$ for $x \in \Omega_n$ and let $(\varphi_{\varepsilon,n}(t; \hbar))_{\varepsilon}$ be infinite sequence $n \in \mathbb{N}$:

$$(\varphi_{\varepsilon,n}(t; \hbar))_{\varepsilon} = \left(\left(\frac{nm}{4\pi i t \hbar} \right)^{d(n+1)/2} \int_{\Omega_n} dx_0 dx_1 \cdots dx_n f_n(x_0, x_1, \dots, x_n) \Psi_0(x_0) \exp \left[\frac{i}{\hbar} S_\varepsilon(x_0, x_1, \dots, x_n; t) \right] \right)_{\varepsilon}. \quad (2.22)$$

Then there exist infinite sequence $\{\hbar_k\}_{k \in \mathbb{N}}$, $\lim_{k \rightarrow \infty} \hbar_k = 0$ such that

$$\left(\lim_{\substack{n \rightarrow \infty \\ k \rightarrow \infty}} \varphi_{\varepsilon,n}(t; \hbar_k) \right)_{\varepsilon} = 0.$$

Proof. Equality (2.23) immediately follows from (2.21).

Remark 2.1. From Lemma 2.2 follows that stationary phase approximation is not a valid asymptotic approximation in the limit $\hbar \rightarrow 0$ for a path-integral (2.14) and (2.18).

3. Exact Quasi-Classical Asymptotic Beyond Maslov Canonical Operator

Theorem 3.1. Let us consider Cauchy problem (2.8) with initial data $\Psi_0(x)$ is given via formula

$$\Psi_0(x) = \frac{\eta^{d/4}}{(2\pi)^{d/4} \hbar^{d/4}} \exp \left[-\frac{\eta(x-x_0)^2}{2\hbar} \right], \quad (3.1)$$

where $0 < \hbar \ll \eta \ll 1$ and $x^2 = \langle x, x \rangle$.

1) We assume now that: a) $(V_\varepsilon(x, t))_{\varepsilon} \in G(\mathbb{R}^d)$, b) $V_{\varepsilon=0}(x, t) = V(x, t): \mathbb{R}_+ \times \mathbb{R}^d \rightarrow \mathbb{R}$ and c) $\forall t \in \mathbb{R}_+$ function $V(x, t)$ is a polynomial on variable $x = (x_1, \dots, x_d)$, i.e.

$$V(x, t) = \sum_{\|\alpha\| \leq m} g_\alpha(t) x^\alpha, \quad \alpha = (i_1, \dots, i_d), \quad x^\alpha = x_1^{i_1} \times \dots \times x_d^{i_d}, \quad \|\alpha\| = \sum_{r=1}^d i_r. \quad (3.2)$$

2) Let $u(\tau, t, \lambda, x, y) = (u_1(\tau, t, \lambda, x, y), \dots, u_d(\tau, t, \lambda, x, y))$ be the solution of the boundary problem

$$\frac{\partial^2 u^T(\tau, t, \lambda, x, y)}{\partial \tau^2} = \text{Hess}[V(\lambda, \tau)] u^T(\tau, t, \lambda, x, y) + [V'(\lambda, \tau)]^T, \quad (3.3)$$

$$u(0, t, \lambda, x, y) = y, \quad u(t, t, \lambda, x, y) = x. \quad (3.4)$$

Here $\lambda = (\lambda_1, \dots, \lambda_d) \in \mathbb{R}^d$, $u^T(\tau, t, \lambda, x, y) = (u_1(\tau, t, \lambda, x, y), \dots, u_d(\tau, t, \lambda, x, y))^T$,

$$V'(\lambda, \tau) = \left(\left[\frac{\partial V(x, t)}{\partial x_1} \right]_{x=\lambda}, \dots, \left[\frac{\partial V(x, t)}{\partial x_d} \right]_{x=\lambda} \right) \quad \text{and} \quad \text{Hess}[V(\lambda, \tau)] = \left[\frac{\partial^2 V(x, t)}{\partial x_i \partial x_j} \right]_{x=\lambda}. \quad (3.5)$$

3) Let $S(t, \lambda, x, y)$ be the master action given via formula

$$S(t, \lambda, x, y) = \int_0^t L(\dot{u}(\tau, t, \lambda, x, y), u(\tau, t, \lambda, x, y), \tau) d\tau, \quad (3.6)$$

where master Lagrangian $L(\dot{u}, u, \tau)$ are

$$L(\dot{u}, u, \tau) = \frac{m}{2} \dot{u}^2(\tau, t, \lambda, x, y) - \hat{V}(u(\tau, t, \lambda, x, y), \tau), \quad \dot{u} = \left(\frac{\partial u_1}{\partial \tau}, \dots, \frac{\partial u_d}{\partial \tau} \right), \quad \dot{u}^2 = \langle \dot{u}, \dot{u} \rangle, \quad (3.7)$$

$$\hat{V}((\tau, t, \lambda, x, y), \tau) = u(\tau, t, \lambda, x, y) \text{Hess}[V(\lambda, \tau)] u^T(\tau, t, \lambda, x, y) + V'(\lambda, \tau) u^T(\tau, t, \lambda, x, y). \quad (3.8)$$

Let $y_{cr} = y_{cr}(t, \lambda, x) \in \mathbb{R}^d$ be solution of the linear system of the algebraic equations

$$\left[\frac{\partial S(t, \lambda, x, y)}{\partial y_i} \right]_{y=y_{cr}} = 0, \quad i = 1, \dots, d. \tag{3.9}$$

4) Let $\hat{x} = \hat{x}(t, \lambda, x_0) \in \mathbb{R}^d$ be solution of the linear system of the algebraic equations

$$y_{cr}(t, \lambda, \hat{x}) + \lambda - x_0 = 0. \tag{3.10}$$

Assume that: for a given values of the parameters t, λ, x_0 the point $\hat{x} = \hat{x}(t, \lambda, x_0)$ is not a focal point on a corresponding trajectory is given by corresponding solution of the boundary problem (3.3). Then for the limiting quantum average given via formulae (1.1) the inequalities is satisfied:

$$\lim_{\varepsilon \rightarrow 0} \lim_{\hbar \rightarrow 0} |i, t, x_0; \hbar - \lambda_i| \leq 2 \left[\det S_{y_{cr} y_{cr}}(t, \lambda, \hat{x}(t, \lambda, x_0), y_{cr}(t, \lambda, \hat{x}(t, \lambda, x_0))) \right]^{-1} |\hat{x}_i(t, \lambda, x_0)|, \quad i = 1, \dots, d. \tag{3.11}$$

Thus one can to calculate the limiting quantum trajectory corresponding to potential $V(x, t)$ by using *transcendental master equation*

$$\hat{x}_i(t, \lambda, x_0) = 0, \quad i = 1, \dots, d. \tag{3.12}$$

Proof. From inequality (A.15) and Theorem A1, using inequalities (A.53.a) and (A.53.b) we obtain

$$\lim_{\sigma \rightarrow 0} \lim_{\varepsilon \rightarrow 0} |\langle \hat{x}_i, T; \sigma, l, \lambda, \varepsilon \rangle - \lambda_i| \leq \lim_{\hbar \rightarrow 0} [\mathcal{R}_1(T, \lambda) + \mathcal{R}_2(T, \lambda)], \quad i = 1, \dots, d, \tag{3.13}$$

where

$$\mathcal{R}_1(T, \lambda) = \int dx \left\{ \int_{q(T)=x} D^+ [q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} \cos \left[\frac{1}{\hbar} S_1(\dot{q}, q, \lambda, T) \right] \right\}^2, \tag{3.14}$$

$$\mathcal{R}_2(T, \lambda) = \int dx \left\{ \int_{q(T)=x} D^+ [q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} \sin \left[\frac{1}{\hbar} S_1(\dot{q}, q, \lambda, T) \right] \right\}^2. \tag{3.15}$$

We note that

$$\mathcal{R}_1(T, \lambda) = \int dx [\tilde{\mathcal{R}}_1(x, T, \lambda)]^2, \quad \mathcal{R}_2(T, \lambda) = \int dx [\tilde{\mathcal{R}}_2(x, T, \lambda)]^2, \tag{3.16}$$

where

$$\begin{aligned} \tilde{\mathcal{R}}_1(x, T, \lambda) &= \int_{q(T)=x} D^+ [q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} \cos \left[\frac{1}{\hbar} S_1(\dot{q}, q, \lambda, T) \right] \\ &= \int dy \int_{\substack{q(T)=x \\ q(0)=y}} D^+ [q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} \cos \left[\frac{1}{\hbar} S_1(\dot{q}, q, \lambda, T) \right] \\ &= \int dy \tilde{\mathcal{R}}_1(x, y, T, \lambda), \end{aligned} \tag{3.17}$$

$$\tilde{\mathcal{R}}_1(x, y, T, \lambda) = \int_{\substack{q(T)=x \\ q(0)=y}} D^+ [q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} \cos \left[\frac{1}{\hbar} S_1(\dot{q}, q, \lambda, T) \right] \tag{3.18}$$

and

$$\begin{aligned} \tilde{\mathcal{R}}_2(x, T, \lambda) &= \int_{q(T)=x} D^+ [q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} \sin \left[\frac{1}{\hbar} S_1(\dot{q}, q, \lambda, T) \right] \\ &= \int dy \int_{\substack{q(T)=x \\ q(0)=y}} D^+ [q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} \sin \left[\frac{1}{\hbar} S_1(\dot{q}, q, \lambda, T) \right] \\ &= \int dy \tilde{\mathcal{R}}_2(x, y, T, \lambda), \end{aligned} \tag{3.19}$$

$$\tilde{\mathcal{R}}_2(x, y, T, \lambda) = \int_{\substack{q(T)=x \\ q(0)=y}} D^+ [q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} \sin \left[\frac{1}{\hbar} S_1(\dot{q}, q, \lambda, T) \right]. \tag{3.20}$$

From Equation (3.18) one obtain

$$\tilde{\mathcal{R}}_1(x, y, T, \lambda) = \frac{1}{2} [\tilde{\mathcal{R}}_{1,1}(x, y, T, \lambda) + \tilde{\mathcal{R}}_{1,2}(x, y, T, \lambda)], \quad (3.21)$$

where

$$\tilde{\mathcal{R}}_{1,1}(x, y, T, \lambda) = \int_{q(0)=y}^{q(T)=x} D^+ [q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} \exp \left[\frac{i}{\hbar} S_1(\dot{q}, q, \lambda, T) \right], \quad (3.22)$$

$$\tilde{\mathcal{R}}_{1,2}(x, y, T, \lambda) = \int_{q(0)=y}^{q(T)=x} D^+ [q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} \exp \left[-\frac{i}{\hbar} S_1(\dot{q}, q, \lambda, T) \right]. \quad (3.23)$$

Let us calculate now path integral $\tilde{\mathcal{R}}_{1,1}(x, y, T, \lambda)$ and path integral $\tilde{\mathcal{R}}_{1,2}(x, y, T, \lambda)$, using stationary phase approximation. From Equation (A.23) follows directly that action $S_1(\dot{q}, q, \lambda, T)$ coincide with *master action* $S(t, \lambda, x, y)$ is given via formulae (3.6)-(3.8) and therefore from Equation (3.22) and Equation (3.23) one obtain

$$\begin{aligned} \tilde{\mathcal{R}}_{1,1}(x, y, T, \lambda) &= \int_{q(0)=y}^{q(T)=x} D^+ [q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} \exp \left[\frac{i}{\hbar} S_1(\dot{q}, q, \lambda, T) \right] \\ &= [x_i]^{\frac{1}{2}} \Psi(y) \exp \left[\frac{i}{\hbar} S(t, \lambda, x, y) \right], \end{aligned} \quad (3.24)$$

and

$$\begin{aligned} \tilde{\mathcal{R}}_{1,2}(x, y, T, \lambda) &= \int_{q(0)=y}^{q(T)=x} D^+ [q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} \exp \left[\frac{i}{\hbar} S_1(\dot{q}, q, \lambda, T) \right] \\ &= [x_i]^{\frac{1}{2}} \Psi(y) \exp \left[-\frac{i}{\hbar} S(t, \lambda, x, y) \right]. \end{aligned} \quad (3.25)$$

From Equation (3.17) and Equation (3.24) we obtain

$$\tilde{\mathcal{R}}_1(x, T, \lambda) = \int dy \tilde{\mathcal{R}}_1(x, y, T, \lambda). \quad (3.26)$$

Substitution Equation (3.25) into Equation (3.26) gives

$$\tilde{\mathcal{R}}_{1,1}(x, T, \lambda) = [x_i]^{\frac{1}{2}} \int dy \Psi(y) \exp \left[\frac{i}{\hbar} S(t, \lambda, x, y) \right]. \quad (3.26')$$

Similarly one obtain

$$\tilde{\mathcal{R}}_{1,2}(x, T, \lambda) = [x_i]^{\frac{1}{2}} \int dy \Psi(y) \exp \left[-\frac{i}{\hbar} S(t, \lambda, x, y) \right]. \quad (3.27)$$

Let us calculate now integral $\tilde{\mathcal{R}}_{1,1}(x, T, \lambda)$ and integral $\tilde{\mathcal{R}}_{1,2}(x, T, \lambda)$ using stationary phase approximation. Let $y_{cr} = y_{cr}(t, \lambda, x) \in \mathbb{R}^d$ be the stationary point of master action $S(t, \lambda, x, y)$ and therefore Equation (3.9) is satisfied. Having applied stationary phase approximation one obtain

$$\begin{aligned} &\tilde{\mathcal{Q}}_{1,1}(x, y_{cr}(t, \lambda, x), T, \lambda) \\ &= \left[\det S_{y_{cr} y_{cr}}(t, \lambda, x, y_{cr}(t, \lambda, x)) \right]^{-\frac{1}{2}} [x_i]^{\frac{1}{2}} \Psi(y_{cr}(t, \lambda, x)) \exp \left[\frac{i}{\hbar} S(t, \lambda, x, y_{cr}(t, \lambda, x)) \right], \end{aligned} \quad (3.28)$$

$$\begin{aligned} &\tilde{\mathcal{Q}}_{1,2}(x, y_{cr}(t, \lambda, x), T, \lambda) \\ &= \left[\det S_{y_{cr} y_{cr}}(t, \lambda, x, y_{cr}(t, \lambda, x)) \right]^{-\frac{1}{2}} [x_i]^{\frac{1}{2}} \Psi(y_{cr}(t, \lambda, x)) \exp \left[-\frac{i}{\hbar} S(t, \lambda, x, y_{cr}(t, \lambda, x)) \right]. \end{aligned} \quad (3.29)$$

Substitution Equations (3.28)-(3.29) into Equation (3.21) gives

$$\begin{aligned}
 \tilde{\mathcal{R}}_1(x, y_{cr}(t, \lambda, x), T, \lambda) &= \frac{1}{2} [\tilde{\mathcal{R}}_{1,1}(x, y_{cr}(t, \lambda, x), T, \lambda) + \tilde{\mathcal{R}}_{1,2}(x, y_{cr}(t, \lambda, x), T, \lambda)] \\
 &= \left[\det S_{y_{cr} y_{cr}}(t, \lambda, x, y_{cr}(t, \lambda, x)) \right]^{-\frac{1}{2}} [x_i]^{-\frac{1}{2}} \Psi(y_{cr}(t, \lambda, x)) \\
 &\quad \times \left\{ \exp \left[\frac{i}{\hbar} S(t, \lambda, x, y_{cr}(t, \lambda, x)) \right] + \exp \left[-\frac{i}{\hbar} S(t, \lambda, x, y_{cr}(t, \lambda, x)) \right] \right\} \\
 &= \left[\det S_{y_{cr} y_{cr}}(t, \lambda, x, y_{cr}(t, \lambda, x)) \right]^{-\frac{1}{2}} [x_i]^{-\frac{1}{2}} \Psi(y_{cr}(t, \lambda, x)) \cos \left[\frac{1}{\hbar} S(t, \lambda, x, y_{cr}(t, \lambda, x)) \right].
 \end{aligned} \tag{3.30}$$

Substitution Equation (3.30) into Equation (3.16) gives

$$\begin{aligned}
 \mathcal{R}_1(T, \lambda) &= \int dx \int dx \left[\tilde{\mathcal{R}}_1(x, T, \lambda) \right]^2 \\
 &= \int dx \left[\det S_{y_{cr} y_{cr}}(t, \lambda, x, y_{cr}(t, \lambda, x)) \right]^{-1} |x_i| \Psi^2(y_{cr}(t, \lambda, x)) \cos^2 \left[\frac{1}{\hbar} S(t, \lambda, x, y_{cr}(t, \lambda, x)) \right].
 \end{aligned} \tag{3.31}$$

Similarly one obtain

$$\begin{aligned}
 \mathcal{R}_2(T, \lambda) &= \int dx \int dx \left[\tilde{\mathcal{R}}_2(x, T, \lambda) \right]^2 \\
 &= \int dx \left[\det S_{y_{cr} y_{cr}}(t, \lambda, x, y_{cr}(t, \lambda, x)) \right]^{-1} |x_i| \Psi^2(y_{cr}(t, \lambda, x)) \sin^2 \left[\frac{1}{\hbar} S(t, \lambda, x, y_{cr}(t, \lambda, x)) \right].
 \end{aligned} \tag{3.32}$$

Therefore

$$\mathcal{R}(T, \lambda) = \mathcal{R}_1(T, \lambda) + \mathcal{R}_2(T, \lambda) = 2 \int dx \left[\det S_{y_{cr} y_{cr}}(t, \lambda, x, y_{cr}(t, \lambda, x)) \right]^{-1} |x_i| \Psi^2(y_{cr}(t, \lambda, x)). \tag{3.33}$$

Substitution Equation (3.1) into Equation (3.33) gives

$$\mathcal{R}(T, \lambda) = 2 \frac{\eta^{d/2}}{(2\pi)^{d/2} \hbar^{d/2}} \int dx \left[\det S_{y_{cr} y_{cr}}(t, \lambda, x, y_{cr}(t, \lambda, x)) \right]^{-1} |x_i| \exp \left[-\frac{\eta(y_{cr}(t, \lambda, x) - x_0)^2}{\hbar} \right]. \tag{3.34}$$

Let us calculate now integral (3.34) using Laplace's approximation. It is easy to see that corresponding stationary point $\hat{x} = \hat{x}(t, \lambda, x_0) \in \mathbb{R}^d$ is the solution of the linear system of the algebraic Equation (3.10). Therefore finally we obtain

$$\mathcal{R}(T, \lambda) = 2 |\hat{x}_i(t, \lambda, x_0)| \left[\det S_{y_{cr} y_{cr}}(t, \lambda, \hat{x}(t, \lambda, x_0), y_{cr}(t, \lambda, \hat{x}(t, \lambda, x_0))) \right]^{-1} + O(\hbar^d), \quad i = 1, \dots, d. \tag{3.35}$$

Substitution Equation (3.35) into inequality (3.13) gives the inequality (3.11). The inequality (3.11) completed the proof.

4. Quantum Anharmonic Oscillator with a Cubic Potential Supplemented by Additive Sinusoidal Driving

In this subsection we calculate exact quasi-classical asymptotic for quantum anharmonic oscillator with a cubic potential supplemented by additive sinusoidal driving. Using Theorem 3.1 we obtain corresponding limiting quantum trajectories given via Equation (1.3).

Let us consider quantum anharmonic oscillator with a cubic potential

$$V(x) = \frac{m\omega^2}{2} x^2 - ax^3 + bx, \quad x \in \mathbb{R}, \quad a, b > 0. \tag{4.1}$$

Supplemented by an additive sinusoidal driving. Thus

$$V(x, t) = \frac{m\omega^2}{2} x^2 - ax^3 + bx - [A \sin(\Omega t)] x. \tag{4.2}$$

The corresponding master Lagrangian given by (3.7), are

$$L(\dot{u}, u, \tau) = \left(\frac{m}{2}\right)\dot{u}^2 - m\left[\left(\frac{\omega^2}{2}\right) + \left(\frac{3a\lambda}{m}\right)\right]u^2 - (m\omega^2\lambda + 3a\lambda^2 - b - A\sin(\Omega t))u. \tag{4.3}$$

We assume now that: $\frac{\omega^2}{2} + \frac{3a\lambda}{m} \geq 0$ and rewrite (4.3) of the form

$$L(\dot{u}, u, \tau) = (m/2)\dot{u}^2 - (m\varpi^2\lambda/2)u^2 + g(\lambda, t)u. \tag{4.4}$$

where $\varpi(\lambda) = \sqrt{2\left[\frac{\omega^2}{2} + \frac{3a\lambda}{m}\right]}$ and $g(\lambda, t) = -[m\omega^2\lambda + 3a\lambda^2 - b - A\sin(\Omega \cdot t)]$.

The corresponding master action $S(t, \lambda, x, y)$ given by Equation (3.6), are

$$S(t, \lambda, x, y) = \frac{m\varpi}{2\sin\varpi t} \left[(\cos\varpi t)(y^2 + x^2) - 2xy + \frac{2x}{m\varpi} \int_0^t g(\lambda, \tau) \sin(\varpi\tau) d\tau + \frac{2y}{m\varpi} \int_0^t g(\lambda, \tau) \sin(\varpi(t-\tau)) d\tau - \frac{2}{m^2\varpi^2} \int_0^t \int_0^\tau g(\lambda, \tau) g(\lambda, s) \sin\varpi(t-\tau) \sin(\varpi s) ds d\tau \right]. \tag{4.5}$$

The linear system of the algebraic Equation (3.9) are

$$\frac{\partial S(t, \lambda, x, y)}{\partial y} = 2y\cos\varpi t - 2x + \frac{2}{m\varpi} \int_0^t g(\lambda, \tau) \sin(\varpi(t-\tau)) d\tau = 0 \tag{4.6}$$

Therefore

$$y_{cr}(t, \lambda, x) = \frac{x}{\cos\varpi t} - \frac{1}{m\varpi\cos\varpi t} \int_0^t g(\lambda, \tau) \sin(\varpi(t-\tau)) d\tau. \tag{4.7}$$

The linear system of the algebraic Equation (3.10) are

$$\frac{x}{\cos\varpi t} - \frac{1}{m\varpi\cos\varpi t} \int_0^t g(\lambda, \tau) \sin(\varpi(t-\tau)) d\tau + \lambda - x_0 = 0. \tag{4.8}$$

Therefore the solution of the linear system of the algebraic Equation (3.10) are

$$\hat{x}(t, \lambda, x_0) = \frac{1}{m\varpi} \int_0^t g(\lambda, \tau) \sin(\varpi(t-\tau)) d\tau + (\lambda(t) - x_0) \cos\varpi t. \tag{4.9}$$

Transcendental master Equation (3.11) are

$$\frac{1}{m\varpi} \int_0^t g(\lambda(t), \tau) \sin(\varpi(t-\tau)) d\tau + (\lambda(t) - x_0) \cos\varpi t = 0 \tag{4.10}$$

Finally from Equation (4.10) one obtain

$$d(\lambda(t)) \left(\frac{\cos(\varpi t)}{\varpi} - \frac{1}{\varpi} \right) + \frac{A(\varpi\sin(\Omega t) - \Omega\sin(\varpi t))}{\varpi^2 - \Omega^2} - (\lambda(t) - x_0) m\varpi\cos(\varpi t) = 0. \tag{4.11}$$

where $d(\lambda) = m\omega^2\lambda + 3a\lambda^2 - b$.

Numerical Examples

Example 1 (in **Figure 1** and **Figure 2**). $x_0 = 0, m = 1, \Omega = 0, \omega = 9, a = 3, b = 10, A = 0$.

5. Comparison Exact Quasi-Classical Asymptotic with Stationary-Point Approximation

We set now $d = 1$. Let us consider now path integral (2.14) with $S_\varepsilon(\dot{q}(t), q(t); t)$ given via formula

$$S_\varepsilon(\dot{q}(t), q(t), t) = \frac{1}{4} \int_0^t [\dot{q}(\tau) - F_\varepsilon(q(\tau), \tau)]^2 d\tau. \tag{5.1}$$




Figure 1. Limiting quantum trajectory $\lambda(t)$ with a jump.




Figure 2. Limiting quantum trajectory $\lambda(t)$ with a jump.

Note that for corresponding propagator $K_\varepsilon(x, t|y, 0)$ the time discretized path-integral representation $K_{\varepsilon, N}(x, t|y, 0)$ are:

$$K_{\varepsilon, N}(x, t|y, 0) = \int \frac{dx_1 dx_2 \cdots dx_{N-1}}{(4\pi\hbar\Delta t)^{N/2}} \exp\left[\frac{i}{\hbar} S_{\varepsilon, N}(x_0, \dots, x_N)\right], \quad (5.2)$$

where $S_{\varepsilon, N}(x_0, \dots, x_N)$ are:

$$S_{\varepsilon, N}(x_0, \dots, x_N) = \frac{\Delta t}{4} \sum_{n=0}^{N-1} \left[\frac{x_{n+1} - x_n}{\Delta t} - F_\varepsilon(x_n, t_n) \right]^2. \quad (5.3)$$

Here the initial- x_0 and end-points x_N are fixed by the prescribed x_0 and by the additional constraint $x_N = y$.

Let us calculate now integral (5.2) using stationary-point approximation. Denoting an critical points of the discrete-time action (5.3) by $\mathbf{x}_{\varepsilon, k} = (x_{1, \varepsilon, k}, \dots, x_{N-1, \varepsilon, k})$ it follows that $\mathbf{x}_{\varepsilon, k}$ satisfies the critical point conditions are

$$\frac{\partial S_{\varepsilon, N}(x_{0, \varepsilon, k}, \dots, x_{N-1, \varepsilon, k}, x_{N, \varepsilon, k})}{\partial x_{n, \varepsilon, k}} = 0 \quad (5.4)$$

for $n = 1, \dots, N-1$, supplemented by the prescribed boundary conditions for $n = 0, n = N: x_{0, \varepsilon, k} = x_0, x_{N, \varepsilon, k} = x$.

From Equation (5.2) in the limit $\hbar \rightarrow 0$ using formally stationary-point approximation one obtain

$$K_{\varepsilon, N}(x, t|x_0, 0) \cong Z_{\varepsilon, N-1}(\mathbf{x}_{\varepsilon, k}, x_{0, \varepsilon, k}, x_{N, \varepsilon, k}) \exp\left[\frac{i}{\hbar} S_{\varepsilon, N}(\mathbf{x}_{\varepsilon, k}, x_{0, \varepsilon, k}, x_{N, \varepsilon, k})\right] + O(\hbar). \quad (5.5)$$

Here the pre-factor $Z_{\varepsilon, N}(\mathbf{x}_{\varepsilon, k})$ is given via N -dimensional Gaussian integral of the canonical form as

$$\mathbf{Z}_{\varepsilon, N-1}(\mathbf{x}_{\varepsilon, k}, x_{0, \varepsilon, k}, x_{N, \varepsilon, k}) = \int \frac{dy_1 dy_2 \cdots dy_{N-1}}{(4\pi\hbar\Delta t)^{N/2}} \exp \left[\frac{i}{2\hbar} \sum_{n,m=1}^{N-1} y_n \frac{\partial^2 \mathcal{S}_{\varepsilon, N}(\mathbf{x}_{\varepsilon, k}, x_{0, \varepsilon, k}, x_{N, \varepsilon, k})}{\partial x_{n, \varepsilon, k} \partial x_{m, \varepsilon, k}} y_m \right]. \quad (5.6)$$

The Gaussian integral in (5.6) is given via canonical formula

$$\mathbf{Z}_{\varepsilon, N-1}(\mathbf{x}_{\varepsilon, k}, x_{0, \varepsilon, k}, x_{N, \varepsilon, k}) = \left[4\pi\hbar\Delta t \det \left(2\Delta t \frac{\partial^2 \mathcal{S}_{\varepsilon, N}(\mathbf{x}_{\varepsilon, k}, x_{0, \varepsilon, k}, x_{N, \varepsilon, k})}{\partial x_{n, \varepsilon, k} \partial x_{m, \varepsilon, k}} \right) \right]^{-1/2}, \quad n, m = 1, \dots, N-1. \quad (5.7)$$

Here $\det(A_{n,m})$ denote the determinant of an $(N-1) \times (N-1)$ matrix with elements $A_{n,m}$. Let us consider now Cauchy problem (2.8) with initial data $\Psi_0(x)$ is given via formula

$$\Psi_0(x) = \frac{\eta^{1/4}}{(2\pi)^{1/4} \hbar^{1/4}} \exp \left[-\frac{\eta(x-z_0)^2}{2\hbar} \right].$$

Note that for corresponding Colombeau solution $\Psi_\varepsilon(t, x)$ given via path-integral (2.14) the time discretized path-integral representation $\Psi_{\varepsilon, N}(t, x)$ are

$$\begin{aligned} \Psi_{\varepsilon, N}(t, x) &= \int \Psi_0(x_0) K_{\varepsilon, N}(x, t | x_0, 0) dx_0 \\ &= \sum_k \int dx_0 \mathbf{Z}_{\varepsilon, N-1}(\mathbf{x}_{\varepsilon, k}, x_0, x_{N, \varepsilon, k}) \Psi_0(x_0) \exp \left[\frac{i}{\hbar} \mathcal{S}_{\varepsilon, N}(\mathbf{x}_{\varepsilon, k}, x_0, x_{N, \varepsilon, k}) \right] [1 + O(\hbar)]. \end{aligned} \quad (5.8)$$

Let us calculate now integrals in RHS of Equation (5.8) using stationary-point approximation. Corresponding critical point conditions are

$$\frac{\partial \mathcal{S}_{\varepsilon, N}(x_{0, \varepsilon, k}, \dots, x_{N-1, \varepsilon, k}, x_{N, \varepsilon, k})}{\partial x_{0, \varepsilon, k}} = 0. \quad (5.9)$$

From (5.8) we obtain

$$\Psi_{\varepsilon, N}(t, x) = \sum_k \mathbf{Z}_{\varepsilon, N}(\mathbf{x}_{\varepsilon, k}, x_{0, \varepsilon, k}, x) \Psi_0(x_{0, \varepsilon, k}) \exp \left[\frac{i}{\hbar} \mathcal{S}_{\varepsilon, N}(\mathbf{x}_{\varepsilon, k}, x_{0, \varepsilon, k}, x) \right] [1 + O(\hbar)]. \quad (5.10)$$

$$\mathbf{Z}_{\varepsilon, N}(\mathbf{x}_{\varepsilon, k}, x_{0, \varepsilon, k}, x) = \left[\Delta t \det \left(2\Delta t \frac{\partial^2 \mathcal{S}_{\varepsilon, N}(\mathbf{x}_{\varepsilon, k}, x_{0, \varepsilon, k}, x)}{\partial x_{n, \varepsilon, k} \partial x_{m, \varepsilon, k}} \right) \right]^{-1/2}, \quad n, m = 0, 1, \dots, N-1. \quad (5.11)$$

Let as denote $\mathbf{x}_{\varepsilon, 0} = (x_{0, \varepsilon, 0}, x_{1, \varepsilon, 0}, \dots, x_{N-1, \varepsilon, 0}) = (x_{0, \varepsilon, 0}(x), x_{1, \varepsilon, 0}(x), \dots, x_{N-1, \varepsilon, 0}(x))$ the critical point for which the critical point conditions (5.4) are

$$\frac{x_{n+1, \varepsilon, 0} - x_{n, \varepsilon, 0}}{\Delta t} - F_\varepsilon(x_{n, \varepsilon, 0}, t_n) = 0, \quad n = 0, 1, \dots, N-1. \quad (5.12)$$

Therefore the time discretized path-integral representation of the Colombeau quantum averages given by Equation (1.1) are

$$\begin{aligned} &(1, t, z_0; \hbar, \varepsilon)_\varepsilon \\ &= \left(\int x |\Psi_{\varepsilon, N}(x, t; \hbar)|^2 dx \right)_\varepsilon \\ &= \frac{\eta^{1/2}}{(2\pi)^{1/2} \hbar^{1/2}} \left(\int dx x \mathbf{Z}_{\varepsilon, N}^2(\mathbf{x}_{\varepsilon, k}, x_{0, \varepsilon, 0}, x) \exp \left[-\frac{\eta(x_{0, \varepsilon, 0}(x) - z_0)^2}{\hbar} \right] \right)_\varepsilon [1 + O(\hbar^2)] \\ &+ \frac{\eta^{1/2}}{(2\pi)^{1/2} \hbar^{1/2}} \left(\sum_{k \geq 1} \int dx x \mathbf{Z}_{\varepsilon, N}^2(\mathbf{x}_{\varepsilon, k}, x_{0, \varepsilon, 0}, x) \exp \left[-\frac{\eta(x_{0, \varepsilon, 0}(x) - z_0)^2}{\hbar} \right] \right)_\varepsilon [1 + O(\hbar^2)] + O(\exp(-c\hbar^{-1})), \end{aligned} \quad (5.13)$$

where $c > 0$, $\varepsilon \in (0, 1]$, $x \in \mathbb{R}$. Let us calculate now integrals in RHS of Equation (5.13) using stationary-point approximation. Corresponding critical point conditions are

$$x_{0,\varepsilon,0} (x_{N,\varepsilon,0}) - z_0 = 0, \tag{5.14}$$

$$x_{0,\varepsilon,k} (x_{N,\varepsilon,k}) - z_0 = 0, \quad k \geq 1. \tag{5.15}$$

Here $x_{N,\varepsilon,0}$ can be calculated using linear recursion (5.12) with initial data $x_{0,\varepsilon,0} = z_0$. From Equations (5.13)-(5.14) one obtain

$$(1, t, z_0; \hbar, \varepsilon)_\varepsilon \cong (x_{N,\varepsilon,0} \mathbf{Z}_{\varepsilon,N}^2 (x_{\varepsilon,k}, x_{0,\varepsilon,0}, x_{N,\varepsilon,0}))_\varepsilon, \tag{5.16}$$

and $\mathbf{Z}_{\varepsilon,N} (x_{\varepsilon,k}, x_{0,\varepsilon,k}, x_{N,\varepsilon,0}) = \left[\Delta t \det \left(2\Delta t \frac{\partial^2 S_{\varepsilon,N} (x_{\varepsilon,k}, x_{0,\varepsilon,k}, x_{N,\varepsilon,0})}{\partial x_{n,\varepsilon,k} \partial x_{m,\varepsilon,k}} \right) \right]^{-1/2}$, $n, m = 0, 1, \dots, N$. (5.17)

As demonstrated in [24] the determinant appearing in (5.11) can be calculated using second order linear recursion:

$$\begin{aligned} \frac{Q_{n+1,\varepsilon,k} - 2Q_{n,\varepsilon,k} - Q_{n-1,\varepsilon,k}}{(\Delta t)^2} &= 2 \frac{Q_{n,\varepsilon,k} F'_x (x_{n,\varepsilon,k}, t_n) - Q_{n-1,\varepsilon,k} F'_x (x_{n-1,\varepsilon,k}, t_{n-1})}{\Delta t} \\ &\quad - Q_{n,\varepsilon,k} \left[\frac{x_{n+1,\varepsilon,k} - x_{n,\varepsilon,k}}{\Delta t} - F_\varepsilon (x_{n,\varepsilon,k}, t_n) \right] F''_{\varepsilon,x^2} (x_{n,\varepsilon,k}, t_n) \\ &\quad + Q_{n,\varepsilon,k} \left[F'_{\varepsilon,x} (x_{n,\varepsilon,k}, t_n) \right]^2 - Q_{n-1,\varepsilon,k} \left[F'_{\varepsilon,x} (x_{n-1,\varepsilon,k}, t_{n-1}) \right]^2, \end{aligned} \tag{5.18}$$

$$F'_{\varepsilon,x} (x, t) = \frac{\partial F_\varepsilon (x, t)}{\partial x}, \quad F''_{\varepsilon,x} (x, t) = \frac{\partial^2 F_\varepsilon (x, t)}{\partial x^2}.$$

with initial data $Q_{1,\varepsilon,k} = \Delta t$, $Q_{2,\varepsilon,k} = Q_{1,\varepsilon,k} + \Delta t + O((\Delta t)^2)$ (5.19)

from which the pre-factor $\mathbf{Z}_{\varepsilon,N} (x_{\varepsilon,k}, x_{0,\varepsilon,k}, x_{N,\varepsilon,0})$ in (5.16) follows as

$$\mathbf{Z}_{\varepsilon,N} (x_{\varepsilon,k}, x_{0,\varepsilon,k}, x_{N,\varepsilon,0}) = \sqrt{Q_{N,\varepsilon,k}}. \tag{5.20}$$

In the limit $\Delta t \rightarrow 0$ from critical point conditions (5.12) and (5.14) one obtain

$$\dot{x}(t) - F_\varepsilon (x(t), t) = 0, \quad x(0) = z_0. \tag{5.21}$$

In the limit $\Delta t \rightarrow 0$ from a second order linear recursion (5.18) one obtain the second order linear differential equation

$$\ddot{Q}_{\varepsilon,0} (t) = 2 \frac{d}{dt} [Q_{\varepsilon,0} (t) F'_{\varepsilon,x} (x(t), t)] \tag{5.22}$$

with initial data

$$Q_{\varepsilon,0} (0) = 0, \quad \dot{Q}_{\varepsilon,0} (0) = 1. \tag{5.23}$$

By integration Equation (5.22) one obtain the first order linear differential equation

$$\dot{Q}_{\varepsilon,0} (t) = 2Q_{\varepsilon,0} (t) F'_{\varepsilon,x} (x(t), t) + 1, \quad Q_{\varepsilon,0} (0) = 0. \tag{5.24}$$

In the limit $\Delta t \rightarrow 0$ from Equation (5.16), Equations (5.20)-(5.21) and Equation (5.24) one obtain

$$E(t) = ((1, t, z_0; \hbar, \varepsilon))_\varepsilon \cong x(t) Q_{\varepsilon,0}^{-2} (t). \tag{5.25}$$

We set now in Equation (5.1)

$$F_\varepsilon (q, \tau) = -aq^3 + bx + \text{Asin}(\Omega\tau). \tag{5.26}$$

Corresponding differential master equation are

$$\dot{q} = -(3a\lambda^2 - b)q - (a\lambda^3 - b\lambda) + A\sin(\Omega\tau), \quad q(0) = z_0 - \lambda. \tag{5.27}$$

From Equation (5.27) one obtain that corresponding transcendental master equation are

$$\begin{aligned} [z_0 - \lambda(t)] - \frac{\Delta[\lambda(t)]}{\Theta[\lambda(t)]} \{ \exp(t\Delta[\lambda(t)]) - 1 \} + A \frac{\exp(t\Delta[\lambda(t)])}{\Omega^2 + \Theta^2[\lambda(t)]} \{ \Theta[\lambda(t)] \sin(\Omega t) - \Omega \cos(\Omega t) \} \\ + A \frac{\Omega}{\Omega^2 + \Theta^2[\lambda(t)]} = 0. \end{aligned} \tag{5.28}$$

Numerical Examples

Comparison of the: 1) classical dynamics calculated by using Equation (5.1) (red curve), 2) limiting quantum trajectory $\lambda(t)$ calculated by using master equation Equation (5.28) (blue curve) and 3) limiting quantum trajectory calculated by using stationary-point approximation given by Equation (5.25) (green curve) (in **Figure 3** and **Figure 4**).




Figure 3. Limiting quantum trajectory $\lambda(t)$ without jumps $a = 0.3, b = 1, A = 2$.




Figure 4. Limiting quantum trajectory $\lambda(t)$ with a jump $a = 1, b = 1, A = 0.3$.

6. Conclusions

We pointed out that there existed limiting quantum trajectories given via Equation (1.3) with a jump. Such jump does not depend on any single measurement of particle position $q(t)$ at instant t and is obtained without any reference to a phenomenological master-equation in Lindblad's form.

An axiom of quantum mechanics is that we cannot predict the result of any single measurement of an observable of a quantum mechanical system in a superposition of eigenstates. However we can predict the result of any single measurement of particle position $q(t)$ at instant t with a probability $P(t, q, \delta q) \cong 1$ if valid the condition: $\lim_{\varepsilon \rightarrow 0} \lim_{\|\delta q(t)\| \rightarrow 0} \lim_{\hbar \rightarrow 0} P_t(q(t), \delta q(t), x_0; \hbar, \varepsilon) = 1$, where P_t is given by Equation (1.5).

Acknowledgements

A reviewer provided important clarification.

References

- [1] Vijay, R., Slichter, D.H. and Siddiqi, I. (2011) Observation of Quantum Jumps in a Superconducting Artificial Atom. *Physical Review Letters*, **106**, Article ID: 110502. <http://dx.doi.org/10.1103/PhysRevLett.106.110502>
- [2] Peil, S. and Gabrielse, G. (1999) Observing the Quantum Limit of an Electron Cyclotron: QND Measurements of Quantum Jumps between Fock States. *Physical Review Letters*, **83**, 1287-1290.
- [3] Sauter, T., Neuhauser, W., Blatt, R. and Toschek, P.E. (1986) Observation of Quantum Jumps. *Physical Review Letters*, **57**, 1696-1698. <http://dx.doi.org/10.1103/PhysRevLett.57.1696>
- [4] Bergquist, J.C., Hulet, R.G., Itano, W.M. and Wineland, D.J. (1986) Observation of Quantum Jumps in a Single Atom. *Physical Review Letters*, **57**, 1699-1702. <http://dx.doi.org/10.1103/PhysRevLett.57.1699>
- [5] Bohr, N. (1913) On the Constitution of Atoms and Molecules. *Philosophical Magazine*, **26**, 1-25. <http://dx.doi.org/10.1080/14786441308634955>
- [6] Dum, R., Zoller, P. and Ritsch, H. (1992) Monte Carlo Simulation of the Atomic Master Equation for Spontaneous Emission. *Physical Review A*, **45**, 4879. <http://dx.doi.org/10.1103/PhysRevA.45.4879>
- [7] Dalibard, J., Castin, Y. and Mølmer, K. (1992) Wave-Function Approach to Dissipative Processes in Quantum Optics. *Physical Review Letters*, **68**, 580.
- [8] Gatarek, D. (1991) Continuous Quantum Jumps and Infinite-Dimensional Stochastic Equations. *Journal of Mathematical Physics*, **32**, 2152-2157.
- [9] Molmer, K., Castin, Y. and Dalibard, J. (1993) Monte Carlo Wave-Function Method in Quantum Optics. *Journal of the Optical Society of America B*, **10**, 524-538. <http://dx.doi.org/10.1364/JOSAB.10.000524>
- [10] Gardiner, C.W., Parkins, A.S. and Zoller, P. (1992) Wave-Function Quantum Stochastic Differential Equations and Quantum-Jump Simulation Methods. *Physical Review A*, **46**, 4363-4381.
- [11] Vogt, N., Jeske, J. and Cole, J.H. (2013) Stochastic Bloch-Redfield Theory: Quantum Jumps in a Solid-Stateenvironment. *Physical Review B*, **88**, Article ID: 174514.
- [12] Reiner, J.E., Wiseman, H.M. and Mabuchi, H. (2003) Quantum Jumps between Dressed States: A Proposed Cavity-QED Test Using Feedback. *Physical Review A*, **67**, Article ID: 042106. <http://dx.doi.org/10.1103/PhysRevA.67.042106>
- [13] Berkeland, D.J., Raymondson, D.A. and Tassin, V.M. (2004) Tests for Non-Randomness in Quantum Jumps. *Physical Review A*, **69**, Article ID: 052103.
- [14] Wiseman, H.M. and Milburn, G.J. (1993) Interpretation of Quantum Jump and Diffusion Processes Illustrated on the Bloch Sphere. *Physical Review A*, **47**, 1652-1666. <http://dx.doi.org/10.1103/PhysRevA.47.1652>
- [15] Wiseman, H.M. and Toombes, G.E. (1999) Quantum Jumps in a Two-Level Atom: Simple Theories versus Quantum Trajectories. *Physical Review A*, **60**, 2474. <http://dx.doi.org/10.1103/PhysRevA.60.2474>
- [16] Stojanovic, M. (2008) Regularization for Heat Kernel in Nonlinear Parabolic Equations. *Taiwanese Journal of Mathematics*, **12**, 63-87. <http://www.tjm.nsysu.edu.tw/>
- [17] Garetto, C. (2008) Fundamental Solutions in the Colombeau Framework: Applications to Solvability and Regularity Theory. *Acta Applicandae Mathematicae*, **102**, 281-318. <http://dx.doi.org/10.1007/s10440-008-9220-8>
- [18] Foukzon, J., Potapov, A.A. and Podosenov, S.A. (2011) Exact Quasiclassical Asymptotics beyond Maslov Canonical

- Operator. *International Journal of Recent advances in Physics*, **3**. <http://arxiv.org/abs/1110.0098>
- [19] Feynman, E.N. (1964) Integrals and the Schrödinger Equation. *Journal of Mathematical Physics*, **5**, 332-343. <http://dx.doi.org/10.1063/1.1704124>
- [20] Maslov, V.P. (1976) Complex Markov Chains and Continual Feynman Integral. Nauka, Moscow.
- [21] Apostol, T.M. (1974) Mathematical Analysis. 2nd Edition, Addison-Wesley, Boston.
- [22] Chen, W.W.L. (2003) Fundamentals of Analysis. Published by Chen, W.W.L. via Internet.
- [23] Gupta, S.L. and Rani, N. (1998) Principles of Real Analysis. Vikas Publishing House, New Delhi.
- [24] Lehmann, J., Reimann, P. and Hänggi, P. (2000) Surmounting Oscillating Barriers: Path-Integral Approach for Weak Noise. *Physical Review E*, **62**, 6282. <http://dx.doi.org/10.1103/PhysRevE.62.6282>
- [25] Fedoryuk, M.V. (1977) The Method of Steepest Descent. Moscow. (In Russian)
- [26] Fedoryuk, M.V. (1989) Asymptotic Methods in Analysis. *Encyclopaedia of Mathematical Sciences*, **13**, 83-191. http://dx.doi.org/10.1007/978-3-642-61310-4_2
- [27] Shun, Z. and McCullagh, P. (1995) Laplace Approximation of High Dimensional Integrals. *Journal of the Royal Statistical Society: Series B (Methodological)*, **57**, 749-760.
- [28] Foukzon, J. (2014) Strong Large Deviations Principles of Non-Freidlin-Wentzell Type-Optimal Control Problem with Imperfect Information—Jumps Phenomena in Financial Markets. *Communications in Applied Sciences*, **2**, 230-363.

Appendix

Let us consider now regularized Feynman-Colombeau propagator $(K_\varepsilon(x, T|y, 0))_\varepsilon$ given by Feynman path integral:

$$\tilde{K}_\varepsilon(x, T|y, 0; \sigma, l) = \int_{q(0)=y}^{q(T)=x} D^+[q(t)] \exp\left[-\frac{1}{\hbar} S_1(q, T; \sigma, l)\right] \exp\left[-\frac{1}{\hbar} S_2(q(T), \lambda)\right] \exp\left[\frac{i}{\hbar} S_\varepsilon(\dot{q}, q, T)\right], \quad (\text{A.1})$$

where $\hbar \in (0, 1]$,

$$S_1(q, T; \sigma, l) = \int_0^T dt \left\{ [q(t) - \lambda]^2; \sigma, l \right\}, \quad (\text{A.2})$$

$$S_2(q(T), \lambda) = [q(T) - \lambda]^2, \quad \lambda = (\lambda_1, \dots, \lambda_d) \in \mathbb{R}^d, \quad (\text{A.3})$$

$$S_\varepsilon(\dot{q}, q, T) = \int_0^T L(\dot{q}(t), q(t), t) dt, \quad L(\dot{q}(t), q(t), t) = \frac{m}{2} \dot{q}^2(t) - V_\varepsilon(q(t), t), \quad (\text{A.4})$$

$$V_{\varepsilon=0}(x, t) = V(x, t), \quad (\text{A.5})$$

$$V(x, t) = g_1(t)x + g_2(t)x^2 + g_3(t)x^3 + \dots + g_\alpha(t)x^\alpha, \quad (\text{A.6})$$

$$\alpha = (i_1, \dots, i_d), \quad x^\alpha = x_1^{i_1} \times \dots \times x_d^{i_d}, \quad \|\alpha\| = \sum_{r=1}^d i_r,$$

$$V_\varepsilon(q(t), t) = V(q_\varepsilon(t), t), \quad q_\varepsilon(t) = (q_{1,\varepsilon}(t), \dots, q_{d,\varepsilon}(t)), \quad (\text{A.7})$$

$$q_{i,\varepsilon}(t) = \frac{q_i(t)}{1 + \varepsilon^{2k} |q(t)|^{2k}}, \quad \varepsilon \in (0, 1], \quad k \geq 1. \quad (\text{A.8})$$

Here: 1) $\sigma \in (0, 1]$, $\hbar \ll \sigma$ and 2) for each path $q(t)$ such that $q(t) = \sum_{i=1}^\infty a_n \sin\left(\frac{n\pi t}{T}\right) + u(t, T, y, x)$, $u(0, T, y, x) = y$, $u(T, T, y, x) = x$, where $u(t, T, y, x)$ is a given function, operator $\{p(t); \sigma, l\}$ are

$$\{q(t); \sigma, l\} = \sum_{n=1}^l \sigma a_n \sin\left(\frac{n\pi t}{T}\right) + \sum_{n=l+1}^\infty a_n \sin\left(\frac{n\pi t}{T}\right). \quad (\text{A.9})$$

3) $D^+[q(t)]$ is a positive Feynman “measure”.

Therefore regularized Colombeau solution of the Schrödinger equation corresponding to regularized propagator (A.1) are

$$\begin{aligned} & (\Psi_\varepsilon(T, x; \sigma, l, \lambda))_\varepsilon \\ &= \left(\int_{-\infty}^\infty dy \Psi(y) \tilde{K}_\varepsilon(x, T|y, 0; \sigma, l) \right)_\varepsilon \\ &= \left(\int_{q(T)=x} D^+[q(t)] \Psi(q(0)) \exp\left[-\frac{1}{\hbar} S(q, T; \sigma, l, \lambda)\right] \exp\left[\frac{i}{\hbar} S_\varepsilon(\dot{q}, q, T)\right] \right)_\varepsilon \\ &= \left(\int dy \int_{q(0)=y}^{q(T)=x} D^+[q(t)] \Psi(q(0)) \exp\left[-\frac{1}{\hbar} S(q, T; \sigma, l, \lambda)\right] \exp\left[\frac{i}{\hbar} S_\varepsilon(\dot{q}, q, T)\right] \right)_\varepsilon. \end{aligned} \quad (\text{A.10})$$

$$\text{Here } S(q, T; \sigma, l, \lambda) = S_1(q, T; \sigma, l) + S_2(q(T), \lambda). \quad (\text{A.11})$$

Let us consider now regularized quantum average

$$(\hat{x}_i, T; \sigma, l, \lambda, \varepsilon)_\varepsilon = \left(\int_{-\infty}^\infty dx x_i |\Psi_\varepsilon(T, x; \sigma, l, \lambda)|^2 \right)_\varepsilon. \quad (\text{A.12})$$

From (A.5) and (A.12) one obtain

$$\begin{aligned}
 & \left(|\hat{x}_i, T; \sigma, l, \lambda, \varepsilon| \right)_\varepsilon \\
 & \leq \left(\int dx \left\{ \int_{q(T)=x} D^+ [q(t)] \Psi(q(0)) [q_i(T)]^{1/2} \exp \left[-\frac{1}{\hbar} S(q, T; \sigma, l, \lambda) \right] \cos \left[\frac{1}{\hbar} S_\varepsilon(\dot{q}, q, T) \right] \right\}^2 \right)_\varepsilon \\
 & \quad + \left(\int dx \left\{ \int_{q(T)=x} D^+ [q(t)] \Psi(q(0)) [q_i(T)]^{1/2} \exp \left[-\frac{1}{\hbar} S(q, T; \sigma, l, \lambda) \right] \sin \left[\frac{1}{\hbar} S_\varepsilon(\dot{q}, q, T) \right] \right\}^2 \right)_\varepsilon.
 \end{aligned} \tag{A.13}$$

From Equations (A.5)-(A.13) one obtain

$$\begin{aligned}
 & |\hat{x}_i, T; \sigma, l, \lambda, \varepsilon - \lambda_i| \\
 & = \left| \hat{x}_i, T; \sigma, l, \lambda, \varepsilon - \lambda_i \int_{-\infty}^{\infty} dx |\Psi_\varepsilon(T, x; \sigma, l, \lambda)|^2 \right| \\
 & = \left| \int_{-\infty}^{\infty} dx [x_i - \lambda_i] |\Psi_\varepsilon(T, x; \sigma, l, \lambda)|^2 \right| \leq \int_{-\infty}^{\infty} dx |x_i - \lambda_i| |\Psi_\varepsilon(T, x; \sigma, l, \lambda)|^2 \\
 & = \int dx \left\{ \int_{q(T)=x} D^+ [q(t)] \Psi(q(0)) [q_i(T) - \lambda_i]^{1/2} \exp \left[-\frac{1}{\hbar} S(q, T; \sigma, l, \lambda) \right] \cos \left[\frac{1}{\hbar} S_\varepsilon(\dot{q}, q, T) \right] \right\}^2 \\
 & \quad + \int dx \left\{ \int_{q(T)=x} D^+ [q(t)] \Psi(q(0)) [q_i(T) - \lambda_i]^{1/2} \exp \left[-\frac{1}{\hbar} S(q, T; \sigma, l, \lambda) \right] \sin \left[\frac{1}{\hbar} S_\varepsilon(\dot{q}, q, T) \right] \right\}^2.
 \end{aligned} \tag{A.14}$$

Using replacement $q_i(t) - \lambda_i := q_i(t)$, $i = 1, \dots, d$ into RHS of the Equation (A.9) one obtain

$$\begin{aligned}
 & |\hat{x}_i, T; \sigma, l, \lambda, \varepsilon - \lambda_i| \\
 & \leq \int dx \left\{ \int_{q(T)=x} D^+ [q(t)] \Psi(q(0)) [q_i(T)]^{1/2} \exp \left[-\frac{1}{\hbar} S(q, T; \sigma, l) \right] \cos \left[\frac{1}{\hbar} S_\varepsilon(\dot{q}, q + \lambda, T) \right] \right\}^2 \\
 & \quad + \int dx \left\{ \int_{q(T)=x} D^+ [q(t)] \Psi(q(0)) [q_i(T)]^{1/2} \exp \left[-\frac{1}{\hbar} S(q, T; \sigma, l) \right] \sin \left[\frac{1}{\hbar} S_\varepsilon(\dot{q}, q + \lambda, T) \right] \right\}^2 \\
 & = \int dx [I_1^2(x, T; \sigma, l, \lambda, \varepsilon)] + \int dx [I_2^2(x, T; \sigma, l, \lambda, \varepsilon)].
 \end{aligned} \tag{A.15}$$

Here

$$\begin{aligned}
 S(q, T; \sigma, l) &= S_1(q, T; \sigma, l) + S_2(q(T)), \\
 S_1(q, T; \sigma, l) &= \int_0^T dt \left[\left[q(t) \right]^2; \sigma, l \right], \quad \lambda \in \mathbb{R}^d. \\
 S_2(q(T)) &= [q(T)]^2,
 \end{aligned} \tag{A.16}$$

And

$$\begin{aligned}
 & I_1(x, T; \sigma, l, \lambda, \varepsilon) \\
 & = \int_{q(T)=x} D^+ [q(t)] \Psi(q(0)) [q_i(T)]^{1/2} \exp \left[-\frac{1}{\hbar} S(q, T; \sigma, l) \right] \cos \left[\frac{1}{\hbar} S_\varepsilon(\dot{q}, q + \lambda, T) \right]
 \end{aligned} \tag{A.17}$$

$$\begin{aligned}
 & I_2(x, T; \sigma, l, \lambda, \varepsilon) \\
 & = \int_{q(T)=x} D^+ [q(t)] \Psi(q(0)) [q_i(T)]^{1/2} \exp \left[-\frac{1}{\hbar} S(q, T; \sigma, l) \right] \sin \left[\frac{1}{\hbar} S_\varepsilon(\dot{q}, q + \lambda, T) \right]
 \end{aligned} \tag{A.18}$$

Let us rewrite a function $V_\varepsilon(q(t) + \lambda, t)$ in the following equivalent form:

$$V_\varepsilon(q(t) + \lambda, t) = V_{\varepsilon,0}(q(t), t, \lambda) + V_{\varepsilon,1}(q(t), t, \lambda), \tag{A.19}$$

$$V_{\varepsilon,0}(q(t),t,\lambda) = a_{\varepsilon,1}(q(t),t,\lambda)q(t) + a_{\varepsilon,2}(q(t),t,\lambda)q^2(t), \quad (\text{A.20})$$

$$V_{\varepsilon,1}(q(t),t,\lambda) = a_{\varepsilon,3}(q(t),t,\lambda)q^3(t) + \dots + a_{\varepsilon,\alpha}(q(t),t,\lambda)q^\alpha(t), \quad (\text{A.21})$$

where $a_{\varepsilon=0,1}(q(t),t,\lambda) = c_1(t,\lambda)$, $a_{\varepsilon=0,2}(q(t),t,\lambda) = c_2(t,\lambda)$, \dots , $a_{\varepsilon=0,\alpha}(q(t),t,\lambda) = c_\alpha(t,\lambda)$.

Let us evaluate now path integral $I_1(T;\sigma,l,\lambda)$ given via Equation (A.17). Substitution Equation (A.19) into RHS of the Equation (A.17) gives

$$\begin{aligned} I_1(x,T;\sigma,l,\lambda,\varepsilon) &= I_1^{(1)}(x,T;\sigma,l,\lambda,\varepsilon) + I_1^{(2)}(x,T;\sigma,l,\lambda,\varepsilon) \\ &= \int_{q(T)=x} D^+[q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} \exp\left[-\frac{1}{\hbar} S(q,T;\sigma,l)\right] \cos\left[\frac{1}{\hbar} S_{\varepsilon,1}(\dot{q},q+\lambda,T)\right] \cos\left[\frac{1}{\hbar} S_{\varepsilon,2}(q+\lambda,T)\right] \\ &\quad + \int_{q(T)=x} D^+[q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} \exp\left[-\frac{1}{\hbar} S(q,T;\sigma,l)\right] \sin\left[\frac{1}{\hbar} S_{\varepsilon,1}(\dot{q},q,\lambda,T)\right] \sin\left[-\frac{1}{\hbar} S_{\varepsilon,2}(q,\lambda,T)\right], \end{aligned} \quad (\text{A.22.a})$$

$$\begin{aligned} I_2(x,T;\sigma,l,\lambda,\varepsilon) &= I_2^{(1)}(x,T;\sigma,l,\lambda,\varepsilon) + I_2^{(2)}(x,T;\sigma,l,\lambda,\varepsilon) \\ &= \int_{q(T)=x} D^+[q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} \exp\left[-\frac{1}{\hbar} S(q,T;\sigma,l)\right] \cos\left[\frac{1}{\hbar} S_{\varepsilon,1}(\dot{q},q+\lambda,T)\right] \sin\left[\frac{1}{\hbar} S_{\varepsilon,2}(q+\lambda,T)\right] \\ &\quad + \int_{q(T)=x} D^+[q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} \exp\left[-\frac{1}{\hbar} S(q,T;\sigma,l)\right] \sin\left[\frac{1}{\hbar} S_{\varepsilon,1}(\dot{q},q,\lambda,T)\right] \cos\left[-\frac{1}{\hbar} S_{\varepsilon,2}(q,\lambda,T)\right] \end{aligned} \quad (\text{A.22.b})$$

where

$$S_{\varepsilon,1}(\dot{q},q,\lambda,T) = \int_0^T L_\varepsilon(\dot{q}(t),q(t),t,\lambda) dt, \quad L_\varepsilon(\dot{q}(t),q(t),t,\lambda) = \frac{m}{2} \dot{q}^2(t) - V_{\varepsilon,0}(q(t),t,\lambda), \quad (\text{A.23})$$

$$S_{\varepsilon,2}(q,\lambda,T) = \int_0^T V_{\varepsilon,1}(q(t),t,\lambda) dt, \quad (\text{A.24})$$

$$\begin{aligned} I_1^{(1)}(x,T;\sigma,l,\lambda,\varepsilon) &= \int_{q(T)=x} D^+[q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} \exp\left[-\frac{1}{\hbar} S(q,T;\sigma,l)\right] \cos\left[\frac{1}{\hbar} S_{\varepsilon,1}(\dot{q},q,\lambda,T)\right] \cos\left[\frac{1}{\hbar} S_{\varepsilon,2}(q,\lambda,T)\right] \end{aligned} \quad (\text{A.25.a})$$

$$\begin{aligned} I_2^{(1)}(x,T;\sigma,l,\lambda,\varepsilon) &= \int_{q(T)=x} D^+[q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} \exp\left[-\frac{1}{\hbar} S(q,T;\sigma,l)\right] \cos\left[\frac{1}{\hbar} S_{\varepsilon,1}(\dot{q},q,\lambda,T)\right] \sin\left[\frac{1}{\hbar} S_{\varepsilon,2}(q,\lambda,T)\right], \end{aligned} \quad (\text{A.25.b})$$

$$\begin{aligned} I_1^{(2)}(x,T;\sigma,l,\lambda,\varepsilon) &= \int_{q(T)=x} D^+[q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} \exp\left[-\frac{1}{\hbar} S(q,T;\sigma,l)\right] \sin\left[\frac{1}{\hbar} S_{\varepsilon,1}(\dot{q},q,\lambda,T)\right] \sin\left[-\frac{1}{\hbar} S_{\varepsilon,2}(q,\lambda,T)\right]. \end{aligned} \quad (\text{A.26.a})$$

$$\begin{aligned} I_2^{(2)}(x,T;\sigma,l,\lambda,\varepsilon) &= \int_{q(T)=x} D^+[q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} \exp\left[-\frac{1}{\hbar} S(q,T;\sigma,l)\right] \sin\left[\frac{1}{\hbar} S_{\varepsilon,1}(\dot{q},q,\lambda,T)\right] \cos\left[-\frac{1}{\hbar} S_{\varepsilon,2}(q,\lambda,T)\right]. \end{aligned} \quad (\text{A.26.b})$$

Let us evaluate now n -dimensional path integral $I_{1,n}^{(1)}(x, T; \sigma, l, \lambda, \varepsilon)$:

$$\begin{aligned}
 & I_{1,n}^{(1)}(x, T; \sigma, l, \lambda, \varepsilon) \\
 &= \int_{q(T)=x} D_n^+ [q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} \exp \left[-\frac{1}{\hbar} S(q, T; \sigma, l) \right] \cos \left[\frac{1}{\hbar} S_{\varepsilon,1}(\dot{q}, q, \lambda, T) \right] \cos \left[\frac{1}{\hbar} S_{\varepsilon,2}(q, \lambda, T) \right] \\
 &= \int_{q(T)=x} D_n^+ [q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} \exp \left[-\frac{1}{\hbar} S(q, T; \sigma, l) \right] \left\{ \cos \left[\frac{1}{\hbar} S_{\varepsilon,1}(\dot{q}, q, \lambda, T) \right] + 1 \right\} \cos \left[\frac{1}{\hbar} S_{\varepsilon,2}(q, \lambda, T) \right] \\
 &\quad - \int_{q(T)=x} D_n^+ [q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} \exp \left[-\frac{1}{\hbar} S(q, T; \sigma, l) \right] \cos \left[\frac{1}{\hbar} S_{\varepsilon,2}(q, \lambda, T) \right].
 \end{aligned} \tag{A.27}$$

From Equation (A.27) one obtain the inequality

$$\begin{aligned}
 & \left| I_{1,n}^{(1)}(x, T; \sigma, l, \lambda, \varepsilon) \right| \\
 &\leq \left| \int_{q(T)=x} D_n^+ [q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} \exp \left[-\frac{1}{\hbar} S(q, T; \sigma, l) \right] \left\{ \cos \left[\frac{1}{\hbar} S_{\varepsilon,1}(\dot{q}, q, \lambda, T) \right] + 1 \right\} \right| \\
 &\quad - \int_{q(T)=x} D_n^+ [q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} \exp \left[-\frac{1}{\hbar} S(q, T; \sigma, l) \right] \cos \left[\frac{1}{\hbar} S_{\varepsilon,2}(q, \lambda, T) \right] \\
 &= \left| \int_{q(T)=x} D_n^+ [q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} \exp \left[-\frac{1}{\hbar} S(q, T; \sigma, l) \right] \cos \left[\frac{1}{\hbar} S_{\varepsilon,1}(\dot{q}, q, \lambda, T) \right] \right| \\
 &\quad + \int_{q(T)=x} D_n^+ [q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} \exp \left[-\frac{1}{\hbar} S(q, T; \sigma, l) \right] \\
 &\quad - \int_{q(T)=x} D_n^+ [q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} \exp \left[-\frac{1}{\hbar} S(q, T; \sigma, l) \right] \cos \left[\frac{1}{\hbar} S_{\varepsilon,2}(q, \lambda, T) \right].
 \end{aligned} \tag{A.28}$$

From In Equation (A.28) one obtain the inequality

$$\begin{aligned}
 & \left| I_{1,n}^{(1)}(x, T; \sigma, l, \lambda, \varepsilon) \right| \\
 &\leq \left| \int_{q(T)=x} D_n^+ [q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} \exp \left[-\frac{1}{\hbar} S(q, T; \sigma, l) \right] \cos \left[\frac{1}{\hbar} S_{\varepsilon,1}(\dot{q}, q, \lambda, T) \right] \right| \\
 &\quad - \sum_{i=1}^{\infty} \frac{(-1)^i \hbar^{-2i}}{(2i)!} \int_{q(T)=x} D_n^+ [q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} [S_{\varepsilon,2}(q, \lambda, T)]^{2i} \exp \left[-\frac{1}{\hbar} S(q, T; \sigma, l) \right] \\
 &= \left| \mathbf{J}_{1,n}^{(1)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar) \right| - \sum_{i=1}^{\infty} \frac{(-1)^i \hbar^{-2i}}{(2i)!} \mathcal{R}_{\varepsilon}^{(i)}(x, T; \sigma, l, n),
 \end{aligned} \tag{A.29}$$

where

$$\begin{aligned}
 & \mathbf{J}_{1,n}^{(1)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar) \\
 &= \int_{q(T)=x} D_n^+ [q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} \exp \left[-\frac{1}{\hbar} S(q, T; \sigma, l) \right] \cos \left[\frac{1}{\hbar} S_{\varepsilon,1}(\dot{q}, q, \lambda, T) \right],
 \end{aligned} \tag{A.30}$$

$$\begin{aligned}
 & \square_{\varepsilon}^{(i)}(x, T; \sigma, l, n) \\
 &= \int_{q(T)=x} D_n^+ [q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} [S_{\varepsilon,2}(q, \lambda, T)]^{2i} \exp \left[-\frac{1}{\hbar} S(q, T; \sigma, l) \right].
 \end{aligned} \tag{A.31}$$

Using replacement $q_i(t) := \hbar^{\frac{1}{2}} q_i(t)$, $t \in [0, T]$, $i = 1, \dots, d$ into RHS of the Equation (A.31) one obtain

$$\begin{aligned}
 \mathcal{R}_\varepsilon^{(i)}(x, T; \sigma, l, n) &= \hbar^{l/4} \int_{q(T)=\frac{x}{\sqrt{\hbar}}} \check{D}_n^+ [q(t)] \Psi \left(\hbar^{\frac{1}{2}} q(0) \right) \left[|q_i(T)| \right]^{\frac{1}{2}} \left[\mathcal{S}_{\varepsilon, 2}(\hbar^{1/2} q, \lambda, T) \right]^{2i} \exp \left[-\frac{1}{\hbar} S(\hbar^{1/2} q, T; \sigma, l) \right] \\
 &= \hbar^{l/4} \hbar^{i/2} \int_{\substack{q(T)=\frac{x}{\sqrt{\hbar}} \\ q(0)=\frac{y}{\sqrt{\hbar}}}} dy \int \check{D}_n^+ [q(t)] \check{\Psi}(q(0)) \left[|q_i(T)| \right]^{\frac{1}{2}} \left[\hat{\mathcal{S}}_{\varepsilon, 2}(q, \lambda, T, \hbar) \right]^{2i} \exp[-S(q, T; \sigma, l)] \\
 &= \hbar^{l/4} \hbar^{i/2} \hat{\mathcal{R}}_\varepsilon^{(i)}(x, T; \sigma, l, n),
 \end{aligned} \tag{A.32}$$

$$\begin{aligned}
 \check{D}_n^+ [q(t)] &= D_n^+ \left[\hbar^{\frac{1}{2}} q(t) \right], \quad t \in [0, T], \quad \check{\Psi}(q(0)) = \frac{\eta^{d/4}}{(2\pi)^{d/4} \hbar^{d/4}} \exp \left[\frac{\eta q^2(0)}{2} \right], \text{ see Equation (3.1) and} \\
 \hat{\mathcal{S}}_{\varepsilon, 2}(q, \lambda, T, \hbar) &= \int_0^T \hat{V}_{\varepsilon, 1}(q(t), t, \lambda, \hbar) dt,
 \end{aligned} \tag{A.33}$$

$$\hat{V}_{\varepsilon, 1}(q(t), t, \lambda, \hbar) = a_{\varepsilon, 3}(q(t), t, \lambda) q^3(t) + \dots + \hbar^{\frac{\alpha-3}{2}} a_{\varepsilon, \alpha}(q(t), t, \lambda) q^\alpha(t). \tag{A.34}$$

$$\hat{\mathcal{R}}_\varepsilon^{(i)}(x, T; \sigma, l, n) = \int dy \int_{\substack{q(T)=\frac{x}{\sqrt{\hbar}} \\ q(0)=\frac{y}{\sqrt{\hbar}}}} \check{D}_n^+ [q(t)] \check{\Psi}(q(0)) \left[|q_i(T)| \right]^{\frac{1}{2}} \left[\hat{\mathcal{S}}_{\varepsilon, 2}(q, \lambda, T, \hbar) \right]^{2i} \exp[-S(q, T; \sigma, l)]. \tag{A.35}$$

From (A.29)-(A.35) one obtain

$$\begin{aligned}
 \left| I_{1, n}^{(1)}(x, T; \sigma, l, \lambda, \varepsilon) \right| &\leq \left| \mathbf{J}_{1, n}^{(1)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar) \right| - \hbar^{\frac{1}{4}} \sum_{i=1}^{\infty} \frac{(-1)^i \hbar^i}{(2i)!} \hat{\mathcal{R}}_\varepsilon^{(i)}(x, T; \sigma, l, n) \\
 &\leq \left| \mathbf{J}_{1, n}^{(1)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar) \right| - \hbar^{\frac{1}{4}} \Xi_\varepsilon(x, T; \sigma, l, n),
 \end{aligned} \tag{A.36}$$

$$\Xi_\varepsilon(x, T; \sigma, l, \hbar, n) = \sum_{i=1}^{\infty} \frac{(-1)^i \hbar^i}{(2i)!} \hat{\mathcal{R}}_\varepsilon^{(i)}(x, T; \sigma, l, n). \tag{A.37}$$

Proposition A.1. [27] [28] Let $\{s_{n, m}\}_{n, m=1}^{n, m=\infty}$ be a double sequence $s: \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{C}$. Let $\lim_{n, m \rightarrow \infty} s_{n, m} = a$.

Then the iterated limit: $\lim_{n \rightarrow \infty} (\lim_{m \rightarrow \infty} s_{n, m})$ exist and equal to a if and only if $\lim_{m \rightarrow \infty} s_{n, m}$ exists for each $n \in \mathbb{N}$.

Proposition A.2. Let $I_1^{(1)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar) = I_1^{(1)}(x, T; \sigma, l, \lambda, \varepsilon)$, where $I_1^{(1)}(x, T; \sigma, l, \lambda, \varepsilon)$ is given via Equation (A.25) and let $I_1^{(2)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar) = I_1^{(2)}(x, T; \sigma, l, \lambda, \varepsilon)$, where $I_1^{(2)}(x, T; \sigma, l, \lambda, \varepsilon)$ is given via Equation (A.26). Then

$$\begin{aligned}
 &\lim_{\sigma \rightarrow 0} \lim_{\hbar \rightarrow 0} \int dx \left[I_1^{(1)}(x, T; \sigma, l, \lambda, \varepsilon) \right]^2 \\
 1) &\leq \lim_{\hbar \rightarrow 0} \int dx \left\{ \int_{q(T)=x} D^+ [q(t)] \Psi(q(0)) \left[|q_i(T)| \right]^{\frac{1}{2}} \cos \left[\frac{1}{\hbar} S_1(\dot{q}, q, \lambda, T) \right] \right\}^2, \\
 2) &\lim_{\sigma \rightarrow 0} \lim_{\hbar \rightarrow 0} \int dx \left[I_1^{(2)}(x, T; \sigma, l, \lambda, \varepsilon) \right]^2 = 0, \\
 3) &\lim_{\sigma \rightarrow 0} \lim_{\hbar \rightarrow 0} \int dx \left[I_1^{(2)}(x, T; \sigma, l, \lambda, \varepsilon) I_1^{(2)}(x, T; \sigma, l, \lambda, \varepsilon) \right] = 0, \\
 &\lim_{\sigma \rightarrow 0} \lim_{\hbar \rightarrow 0} \int dx \left[I_2^{(2)}(x, T; \sigma, l, \lambda, \varepsilon) \right]^2 \\
 4) &\leq \lim_{\hbar \rightarrow 0} \int dx \left\{ \int_{q(T)=x} D^+ [q(t)] \Psi(q(0)) \left[|q_i(T)| \right]^{\frac{1}{2}} \sin \left[\frac{1}{\hbar} S_1(\dot{q}, q, \lambda, T) \right] \right\}^2,
 \end{aligned}$$

- 5) $\lim_{\sigma \rightarrow 0} \lim_{\hbar \rightarrow 0} \int dx \left[I_2^{(1)}(x, T; \sigma, l, \lambda, \varepsilon) \right]^2 = 0,$
 6) $\lim_{\sigma \rightarrow 0} \lim_{\hbar \rightarrow 0} \int dx \left[I_2^{(2)}(x, T; \sigma, l, \lambda, \varepsilon) I_2^{(1)}(x, T; \sigma, l, \lambda, \varepsilon) \right] = 0.$

Here

$$S_1(\dot{q}, q, \lambda, T) = S_{\varepsilon=0,1}(\dot{q}, q, \lambda, T) = \int_0^T L_{\varepsilon=0}(\dot{q}(t), q(t), t, \lambda) dt,$$

$$L_{\varepsilon=0}(\dot{q}(t), q(t), t, \lambda) = \frac{m}{2} \dot{q}^2(t) - V_{\varepsilon=0,0}(q(t), t, \lambda).$$

Proof (I) Let us to choose an sequence $\{\hbar_m\}_{m=1}^\infty$ such that

1) $\lim_{m \rightarrow \infty} \hbar_m = 0$ and

2) $\lim_{m, n \rightarrow \infty} \int dx \left\{ \Xi_\varepsilon^{(m)}(x, T; \sigma, l, \hbar_m, n) \right\}^2 = \lim_{m, n \rightarrow \infty} \int dx \left\{ \sum_{i=1}^m \frac{(-1)^i \hbar_m^i}{(2i)!} \hat{\mathcal{R}}_\varepsilon^{(i)}(x, T; \sigma, l, n) \right\}^2 = 0.$

We note that from (ii) follows that: perturbative expansion

$$\begin{aligned} & \int dx \left\{ \Xi_\varepsilon(x, T; \sigma, l, \hbar_m, n) \right\}^2 \\ &= \hbar_m^{1/4} \int dx \left\{ \sum_{i=1}^\infty \frac{(-1)^i \hbar_m^i}{(2i)!} \hat{\Xi}_\varepsilon^{(i)}(x, T; \sigma, l, n) \right\}^2 \end{aligned}$$

vanishes in the limit $m, n \rightarrow \infty$. From (A.36) and Schwarz's inequality using Proposition A.1, one obtain

$$\begin{aligned} & \lim_{m, n \rightarrow \infty} \int dx \left[I_{1,n}^{(1)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar_m) \right]^2 \\ & \leq \lim_{m, n \rightarrow \infty} \int dx \left\{ \left| \mathbf{J}_{1,n}^{(1)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar_m) \right| - \hbar_m^{1/4} \Xi_\varepsilon(x, T; \sigma, l, \hbar, n) \right\}^2 \\ & \leq \lim_{m, n \rightarrow \infty} \int dx \left\{ \mathbf{J}_{1,n}^{(1)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar_m) \right\}^2 \\ & \quad + \lim_{m, n \rightarrow \infty} \left\{ 2\hbar_m^{1/4} \sqrt{\int dx \left\{ \mathbf{J}_{1,n}^{(1)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar_m) \right\}^2 \int dx \left\{ \Xi_\varepsilon(x, T; \sigma, l, \hbar_m, n) \right\}^2} + \int dx \left\{ \Xi_\varepsilon(x, T; \sigma, l, \hbar_m, n) \right\}^2 \right\} \\ & = \lim_{m, n \rightarrow \infty} \int dx \left\{ \mathbf{J}_{1,n}^{(1)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar_m) \right\}^2 \\ & = \lim_{\hbar \rightarrow 0} \lim_{n \rightarrow \infty} \int dx \left\{ \mathbf{J}_{1,n}^{(1)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar) \right\}^2 \\ & = \lim_{\hbar \rightarrow 0} \int dx \left\{ \int_{q(T)=x} D^+ [q(t)] \Psi(q(0)) \left[|q_i(T)| \right]^{\frac{1}{2}} \cos \left[\frac{1}{\hbar} S_1(\dot{q}, q, \lambda, T) \right] \right\}^2. \end{aligned} \tag{A.38}$$

Let us to choose now an subsequence $\{\hbar_{m_k}\}_{m_k=1}^\infty$ such that the limit: $\lim_{k, n \rightarrow \infty} \int dx \left[I_{1,n}^{(1)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar_{m_k}) \right]^2$

exist and $\lim_{k, n \rightarrow \infty} \int dx \left[I_{1,n}^{(1)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar_{m_k}) \right]^2 = \lim_{k, n \rightarrow \infty} \int dx \left[I_{1,n}^{(1)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar_{m_k}) \right]^2,$ (A.39)

From (A.39) and Proposition A.1 one obtain

$$\begin{aligned} & \lim_{k, n \rightarrow \infty} \int dx \left[I_{1,n}^{(1)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar_{m_k}) \right]^2 \\ & = \lim_{k \rightarrow \infty} \left\{ \lim_{n \rightarrow \infty} \int dx \left[I_{1,n}^{(1)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar_{m_k}) \right]^2 \right\}. \end{aligned} \tag{A.40}$$

From (A.39), (A.40) and (A.38) one obtain

$$\begin{aligned}
 & \lim_{\sigma \rightarrow 0} \lim_{\varepsilon \rightarrow 0} \lim_{\hbar \rightarrow 0} \int dx \left[I_1^{(1)}(x, T; \sigma, l, \lambda, \varepsilon) \right]^2 \\
 & \leq \lim_{\sigma \rightarrow 0} \lim_{\varepsilon \rightarrow 0} \lim_{k \rightarrow \infty} \left\{ \lim_{n \rightarrow \infty} \int dx \left[I_{1,n}^{(1)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar_{m_k}) \right]^2 \right\} \\
 & = \lim_{\sigma \rightarrow 0} \lim_{\varepsilon \rightarrow 0} \lim_{k \rightarrow \infty} \int dx \left[I_1^{(1)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar_{m_k}) \right]^2 \\
 & \leq \lim_{\hbar \rightarrow 0} \int dx \left\{ \int_{q(T)=x} D^+ [q(t)] \Psi(q(0)) \left[|q_i(T)| \right]^{\frac{1}{2}} \cos \left[\frac{1}{\hbar} S_1(\dot{q}, q, \lambda, T) \right] \right\}^2.
 \end{aligned} \tag{A.41}$$

The inequality (A.41) completed the proof of the statement (1).

(II) Let us estimate now n -dimensional path integral

$$\begin{aligned}
 & I_{1,n}^{(2)}(x, T; \sigma, l, \lambda, \varepsilon) \\
 & = \int_{q(T)=x} D_n^+ [q(t)] \Psi(q(0)) \left[|q_i(T)| \right]^{1/2} \exp \left[-\frac{1}{\hbar} S(q, T; \sigma, l) \right] \sin \left[\frac{1}{\hbar} S_{\varepsilon,1}(\dot{q}, q, \lambda, T) \right] \sin \left[-\frac{1}{\hbar} S_{\varepsilon,2}(q, \lambda, T) \right],
 \end{aligned} \tag{A.42}$$

From Equation (A.42) one obtain the inequality

$$\begin{aligned}
 \left| I_{1,n}^{(2)}(x, T; \sigma, l, \lambda, \varepsilon) \right| & \leq \int_{q(T)=x} D_n^+ [q(t)] \Psi(q(0)) \left[|q_i(T)| \right]^{1/2} \exp \left[-\frac{1}{\hbar} S(q, T; \sigma, l) \right] \left| \sin \left[-\frac{1}{\hbar} S_{\varepsilon,2}(q, \lambda, T) \right] \right| \\
 & \leq \sum_{i=0}^{\infty} \frac{\hbar^{-(2i+1)}}{(2i+1)!} \int_{q(T)=x} D_n^+ [q(t)] \Psi(q(0)) \left[|q_i(T)| \right]^{\frac{1}{2}} \left[|S_{\varepsilon,2}(q, \lambda, T)| \right]^{(2i+1)} \exp \left[-\frac{1}{\hbar} S(q, T; \sigma, l) \right] \\
 & = \sum_{i=0}^{\infty} \frac{\hbar^{-(2i+1)}}{(2i+1)!} \wp_{\varepsilon}^{(i)}(x, T; \sigma, l, n),
 \end{aligned} \tag{A.43}$$

where

$$\wp_{\varepsilon}^{(i)}(x, T; \sigma, l, n) = \int_{q(T)=x} D_n^+ [q(t)] \Psi(q(0)) \left[|q_i(T)| \right]^{\frac{1}{2}} \left[|S_{\varepsilon,2}(q, \lambda, T)| \right]^{(2i+1)} \exp \left[-\frac{1}{\hbar} S(q, T; \sigma, l) \right]. \tag{A.44}$$

Using replacement $q_i(t) := \hbar^{\frac{1}{2}} \tilde{q}_i(t)$, $t \in [0, T]$, $i = 1, \dots, d$ into RHS of the Equation (A.44) one obtain

$$\begin{aligned}
 \wp_{\varepsilon}^{(i)}(x, T; \sigma, l, n) & = \hbar^{1/4} \int_{q(T)=\frac{x}{\sqrt{\hbar}}} \tilde{D}_n^+ [q(t)] \Psi \left(\hbar^{\frac{1}{2}} q(0) \right) \left[|q_i(T)| \right]^{\frac{1}{2}} \left[|S_{\varepsilon,2}(q, \lambda, T)| \right]^{(2i+1)} \exp \left[-\frac{1}{\hbar} S(\hbar^{1/2} q, T; \sigma, l) \right] \\
 & = \hbar^{1/4} \hbar^{i/2} \int_{\substack{q(T)=\frac{x}{\sqrt{\hbar}} \\ q(0)=\frac{y}{\sqrt{\hbar}}}} dy \int \tilde{D}_n^+ [q(t)] \tilde{\Psi}(q(0)) \left[|q_i(T)| \right]^{\frac{1}{2}} \left[|\hat{S}_{\varepsilon,2}(q, \lambda, T)| \right]^{(2i+1)} \exp[-S(q, T; \sigma, l)] \\
 & = \hbar^{1/4} \hbar^{i/2} \hat{\wp}_{\varepsilon}^{(i)}(x, T; \sigma, l, n),
 \end{aligned} \tag{A.45}$$

$$\tilde{D}_n^+ [q(t)] = D_n^+ \left[\hbar^{\frac{1}{2}} q(t) \right], \quad t \in [0, T], \quad \tilde{\Psi}(q(0)) = \frac{\eta^{d/4}}{(2\pi)^{d/4} \hbar^{d/4}} \exp \left[\frac{\eta q^2(0)}{2} \right], \text{ see Equation (3.1) and}$$

$$\hat{S}_{\varepsilon,2}(q, \lambda, T, \hbar) = \int_0^T \hat{V}_{\varepsilon,1}(q(t), t, \lambda, \hbar) dt, \tag{A.46}$$

$$\hat{V}_{\varepsilon,1}(q(t), t, \lambda, \hbar) = a_{\varepsilon,3}(q(t), t, \lambda) q^3(t) + \dots + \hbar^{\frac{\alpha-3}{2}} a_{\varepsilon,\alpha}(q(t), t, \lambda) q^{\alpha}(t). \tag{A.47}$$

$$\hat{\rho}_\varepsilon^{(i)}(x, T; \sigma, l, n) = \int_{q(0)=\frac{x}{\sqrt{\hbar}}}^{\int_{q(T)=\frac{x}{\sqrt{\hbar}}} D_n^+} \tilde{\Psi}(q(0)) [q_i(T)]^{\frac{1}{2}} \left[\hat{S}_{\varepsilon, 2}(q, \lambda, T) \right]^{(2i+1)} \exp[-S(q, T; \sigma, l)]. \quad (\text{A.48})$$

From (A.43)-(A.48) one obtain

$$\left| I_{1,n}^{(2)}(x, T; \sigma, l, \lambda, \varepsilon) \right| \leq \hbar^{\frac{1}{4}} \sum_{i=0}^{\infty} \frac{\hbar^{2(i+1)}}{(2i+1)!} \hat{\rho}_\varepsilon^{(i)}(x, T; \sigma, l, n) = \Theta_\varepsilon(x, T; \sigma, l, \hbar, n). \quad (\text{A.49})$$

Let us to choose an sequence $\{\hbar_m\}_{m=1}^{\infty}$ such that

1) $\lim_{m \rightarrow \infty} \hbar_m = 0$ and

$$2) \lim_{m, n \rightarrow \infty} \int dx \left\{ \Theta_\varepsilon^{(m)}(x, T; \sigma, l, \hbar_m, n) \right\}^2 = \lim_{m, n \rightarrow \infty} \int dx \left\{ \sum_{i=0}^m \frac{\hbar_m^{2(i+1)}}{(2i+1)!} \hat{\rho}_\varepsilon^{(i)}(x, T; \sigma, l, n) \right\}^2 = 0.$$

We note that from 2) follows that: perturbative expansion

$$\int dx \left\{ \Theta_\varepsilon(x, T; \sigma, l, \hbar_m, n) \right\}^2 = \hbar_m^{1/4} \int dx \left\{ \sum_{i=0}^{\infty} \frac{\hbar_m^{2(i+1)}}{(2i+1)!} \hat{\rho}_\varepsilon^{(i)}(x, T; \sigma, l, n) \right\}^2,$$

Vanishes in the limit $m, n \rightarrow \infty$. From (A.49) one obtain

$$\underline{\lim}_{m, n \rightarrow \infty} \int dx \left[I_{1,n}^{(2)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar_m) \right]^2 \leq \underline{\lim}_{m, n \rightarrow \infty} \int dx \left\{ \Theta_\varepsilon(x, T; \sigma, l, \hbar_m, n) \right\}^2. \quad (\text{A.50})$$

Let us to choose now an subsequence $\{\hbar_{m_k}\}_{m_k=1}^{\infty}$ such that the limit: $\lim_{k, n \rightarrow \infty} \int dx \left[I_{1,n}^{(2)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar_{m_k}) \right]^2$

$$\text{exist and } \lim_{k, n \rightarrow \infty} \int dx \left[I_{1,n}^{(2)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar_{m_k}) \right]^2 = \underline{\lim}_{m, n \rightarrow \infty} \int dx \left[I_{1,n}^{(2)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar_m) \right]^2 \quad (\text{A.51})$$

From (A.51) and Proposition A.1 one obtain

$$\lim_{k, n \rightarrow \infty} \int dx \left[I_{1,n}^{(2)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar_{m_k}) \right]^2 = \lim_{k \rightarrow \infty} \left\{ \lim_{n \rightarrow \infty} \int dx \left[I_{1,n}^{(2)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar_{m_k}) \right]^2 \right\}. \quad (\text{A.52})$$

From (A.50), (A.51) and (A.52) one obtain

$$\begin{aligned} \underline{\lim}_{\sigma \rightarrow 0} \underline{\lim}_{\hbar \rightarrow 0} \int dx \left[I_1^{(2)}(x, T; \sigma, l, \lambda, \varepsilon) \right]^2 &\leq \underline{\lim}_{\sigma \rightarrow 0} \lim_{k \rightarrow 0} \left\{ \lim_{n \rightarrow 0} \int dx \left[I_{1,n}^{(2)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar_{m_k}) \right]^2 \right\} \\ &= \underline{\lim}_{\sigma \rightarrow 0} \lim_{k \rightarrow 0} \int dx \left[I_1^{(2)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar_{m_k}) \right]^2 = 0, \end{aligned}$$

Proof of the statements (3)-(6) is similarly to the proof of the statements (1)-(2).

Theorem A.1. Let $I_1(x, T; \sigma, l, \lambda, \varepsilon, \hbar) = I_1(x, T; \sigma, l, \lambda, \varepsilon)$, $I_2(x, T; \sigma, l, \lambda, \varepsilon, \hbar) = I_2(x, T; \sigma, l, \lambda, \varepsilon)$, where $I_1(x, T; \sigma, l, \lambda, \varepsilon)$ is given via Equations (A.22a)-(A.22b). Then

$$\underline{\lim}_{\sigma \rightarrow 0} \underline{\lim}_{\hbar \rightarrow 0} \int dx \left[I_1^2(x, T; \sigma, l, \lambda, \varepsilon) \right] \leq \lim_{\hbar \rightarrow 0} \int dx \left\{ \int_{q(T)=x} D^+ [q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} \sin \left[\frac{1}{\hbar} S_1(\dot{q}, q, \lambda, T) \right] \right\}^2 \quad (\text{A.53.a})$$

$$\underline{\lim}_{\sigma \rightarrow 0} \underline{\lim}_{\hbar \rightarrow 0} \int dx \left[I_2^2(x, T; \sigma, l, \lambda, \varepsilon) \right] \leq \lim_{\hbar \rightarrow 0} \int dx \left\{ \int_{q(T)=x} D^+ [q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} \sin \left[\frac{1}{\hbar} S_1(\dot{q}, q, \lambda, T) \right] \right\}^2 \quad (\text{A.53.b})$$

Here

$$S_1(\dot{q}, q, \lambda, T) = S_{\varepsilon=0,1}(\dot{q}, q, \lambda, T) = \int_0^T L_{\varepsilon=0}(\dot{q}(t), q(t), t, \lambda) dt, \quad (\text{A.54})$$

$$L_{\varepsilon=0}(\dot{q}(t), q(t), t, \lambda) = \frac{m}{2} \dot{q}^2(t) - V_{\varepsilon=0,0}(q(t), t, \lambda). \quad (\text{A.55})$$

Proof. We remain that

$$I_1(x, T; \sigma, l, \lambda, \varepsilon, \hbar) = I_1^{(1)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar) + I_1^{(2)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar). \quad (\text{A.56})$$

From Equation (A.56) we obtain

$$\begin{aligned} \int dx [I_1^2(x, T; \sigma, l, \lambda, \varepsilon, \hbar)] &\leq \int dx [I_1^{(1)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar)]^2 + \int dx [I_1^{(2)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar)]^2 \\ &\quad + 2 \int dx [I_1^{(1)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar) I_1^{(2)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar)] \\ &= \int dx [I_1^{(1)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar)]^2 + \int dx [I_1^{(2)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar)]^2 \\ &\quad + 2 \sqrt{\int dx [I_1^{(1)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar)]^2 \int dx [I_1^{(2)}(x, T; \sigma, l, \lambda, \varepsilon, \hbar)]^2}. \end{aligned} \quad (\text{A.57})$$

Let us to choose now an sequences $\{\hbar_m\}_{m=1}^\infty$, $\{\varepsilon_k\}_{k=1}^\infty$, $\{\sigma_l\}_{l=1}^\infty$ such that:

$$\begin{aligned} 1) \quad &\lim_{m \rightarrow \infty} \hbar_m = 0, \quad \lim_{k \rightarrow \infty} \varepsilon_k = 0, \quad \lim_{l \rightarrow \infty} \sigma_l = 0, \\ 2) \quad &\lim_{l \rightarrow \infty} \lim_{k \rightarrow \infty} \lim_{m \rightarrow \infty} \int dx [I_1^{(2)}(x, T; \sigma_l, l, \lambda, \varepsilon_k, \hbar_m)]^2 = \lim_{l \rightarrow \infty} \lim_{k \rightarrow \infty} \lim_{m \rightarrow \infty} \int dx [I_1^{(2)}(x, T; \sigma_l, l, \lambda, \varepsilon_k, \hbar_m)]^2 = 0, \end{aligned} \quad (\text{A.58})$$

$$\begin{aligned} &\lim_{l \rightarrow \infty} \lim_{k \rightarrow \infty} \lim_{m \rightarrow \infty} \int dx [I_1^{(1)}(x, T; \sigma_l, l, \lambda, \varepsilon_k, \hbar_m)]^2 \\ 3) \quad &\leq \lim_{\hbar \rightarrow 0} \int dx \left\{ \int_{q(T)=x} D^+ [q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} \cos \left[\frac{1}{\hbar} S_1(\dot{q}, q, \lambda, T) \right] \right\}^2. \end{aligned} \quad (\text{A.59})$$

Therefore from inequality (A.57), Equation (A.58) and inequality (A.59) we obtain

$$\begin{aligned} &\lim_{\sigma \rightarrow 0} \lim_{\hbar \rightarrow 0} \int dx [I_1^2(x, T; \sigma, l, \lambda, \varepsilon, \hbar)] \\ &\leq \lim_{l \rightarrow \infty} \lim_{k \rightarrow \infty} \lim_{m \rightarrow \infty} \int dx [I_1^2(x, T; \sigma_l, l, \lambda, \varepsilon_k, \hbar_m)] \\ &\leq \lim_{l \rightarrow \infty} \lim_{k \rightarrow \infty} \lim_{m \rightarrow \infty} \int dx [I_1^{(1)}(x, T; \sigma_l, l, \lambda, \varepsilon_k, \hbar_m)]^2 + \lim_{l \rightarrow \infty} \lim_{k \rightarrow \infty} \lim_{m \rightarrow \infty} \int dx [I_1^{(2)}(x, T; \sigma_l, l, \lambda, \varepsilon_k, \hbar_m)]^2 \\ &\quad + 2 \lim_{l \rightarrow \infty} \lim_{k \rightarrow \infty} \lim_{m \rightarrow \infty} \sqrt{\int dx [I_1^{(1)}(x, T; \sigma_l, l, \lambda, \varepsilon_k, \hbar_m)]^2 \int dx [I_1^{(2)}(x, T; \sigma_l, l, \lambda, \varepsilon_k, \hbar_m)]^2} \\ &= \lim_{l \rightarrow \infty} \lim_{k \rightarrow \infty} \lim_{m \rightarrow \infty} \int dx [I_1^{(1)}(x, T; \sigma_l, l, \lambda, \varepsilon_k, \hbar_m)]^2 \\ &= \lim_{\hbar \rightarrow 0} \int dx \left\{ \int_{q(T)=x} D^+ [q(t)] \Psi(q(0)) [q_i(T)]^{\frac{1}{2}} \cos \left[\frac{1}{\hbar} S_1(\dot{q}, q, \lambda, T) \right] \right\}^2. \end{aligned} \quad (\text{A.60})$$