

Microbiological Identification of Atmospheric Particles in Hermosillo, Sonora, México

Santos-Romo Alvaro¹, Sau-Acosta Nancy Janeth¹, Certucha-Barragán María Teresa², Almendáriz-Tapia Francisco Javier², Monge-Amaya Onofre², Hernández-López Jorge¹, Zepeda-Ibarra Joel³

¹Nortwest Biological Investigation Centre, Hermosillo, Mexico

²Department of Chemical Engineering & Metallurgy within the Engineering Division of the University of Sonora, Hermosillo, Mexico

³Urban Development and Ecology, General Coordination of Infrastructure, Municipal Institute of Ecology, Hermosillo, Mexico

Email: jhlopez04@cibnor.mx

Received 18 February 2014; revised 16 March 2014; accepted 12 April 2014

Copyright © 2014 by authors and Scientific Research Publishing Inc.

This work is licensed under the Creative Commons Attribution International License (CC BY).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

Abstract

The air quality in many cities has been deteriorating and affecting population's health. Airborne particles can block the defense mechanisms of respiratory system and may lead biological elements such as pollen, bacteria, fungi and viruses into the lungs. This paper presents the study of biologically contaminated air in the city of Hermosillo, analyzing filters containing particulate matter with a diameter equal or less than 10 micrometers (PM₁₀) and Total Suspended Particles (TSP). Three sampling periods were analyzed during 2011. Samples taken from the filters were placed in tubes containing nutritive broth BHI (Brain Heart Infusion) and incubated at 37 °C for 24 hours. The identification of the isolated microorganisms was done by a biochemical test recommended in Bergey's Manual. Results showed a high bacteria concentration in the presence of microorganisms such as *Klebsiella pneumoniae sub pneumoniae*, *Citrobacter*, *Enterococcus faecalis*, *Proteus penneri*, *Butiauxella agrestis*, *Salmonella paratyphi A* and *Enterobacter sacazakii*. When bacterial concentration was analyzed separately, the lowest number of bacteria was observed during February for the south location and in August for the north location. The highest degree of bacterial air pollution was found in the central area of the city.

Keywords

Air Pollution, Airborne Particles, Bacteria

1. Introduction

In most cities, the increase of air quality deterioration has reached a level that affects the population's health. One of the most serious pollutants are atmospheric particles, such as dust, pollen, carbon and metals like lead and cadmium, asbestos, salts, sulfuric acid droplets, dioxins, pesticides, etc. [1], that remain suspended in the atmosphere.

These particles are capable of blocking the respiratory system air defense mechanisms at the upper airway level and alveoli which may carry biological elements such as pollen, bacteria, fungi and viruses that are conducted to the lungs. Particles larger than 10 microns are retained primarily in the upper respiratory tract; particles which are less than 10 microns remain in the breathable fraction and penetrate into the alveolar space of the lungs, Mexican Official Norm (NOM) [2]. High concentration of airborne particles can cause or worsen cardiovascular and lung diseases, heart attacks and arrhythmias, they also affect the central nervous and reproductive systems and may cause cancer [3] [4].

The atmosphere lacks a native microbiota, which acts as a dispersion medium for many types of microorganisms (spores, bacteria, viruses and fungi), from others environments. Microorganisms dispersed into the air have a high biological and economic importance; they also produce illness in plants, animals and human beings since they can quickly be transported in bioaerosol forms causing alterations in foods and organic materials, and contribute to the deterioration and corrosion of monuments and metal buildings.

Several viral, bacterial and fungal infections of human beings and animals are airborne transmitted and often produce epidemic outbreaks. These can be classified into respiratory diseases (pneumonia, whooping cough, tuberculosis, legionellosis, cold, flu), systemic diseases (meningitis, measles, varicella, fungal infections) and allergies [5]. Acute respiratory infections are not produced by inert particles, but by a microorganism that is associated to them, therefore the importance of biological identification of PM₁₀ and TSP.

In the city of Hermosillo, Mexico, the number of cases of Acute Respiratory Infections (ARI's) between 2006 and 2010, were provided by the Sonora State Children's Hospital. The number of ARI's between 2006 and 2010 affected mainly children ranging ages from 1 to 4 years. The causal agents of these infections were, among others, *Klebsiella pneumoniae sub pneumoniae* and *Pseudomonas*.

Microbiological analysis of atmospheric particles including infectious, allergens and fungal agents among others, were determined by capturing them in solid, liquid or agar media for subsequent qualitative or quantitative analysis using biochemical, microscopic, immunochemical, microbiological and molecular techniques, that offer various degrees of sensitivity and specificity [6]. Many of the aerosols sampling devices use particle separation methods from airflow through capture in preselected medium. More common sampling methods used to separate and capture bioaerosols include impaction, filtration and impingement. Impaction method employs a surface of a broad spectrum solid medium (agar); the filtration uses a membrane filter and the impingement an isotonic liquid medium (water-based) [7]. These techniques are used for low volume samples and work for small periods (minutes).

The main goal of this investigation was to identify the microflora present in the air of Hermosillo, México, using high volume samplers, which suck air from the environment and retain particles in glass and quartz microfiber filters. Then the particles were analyzed using biochemical techniques for microorganism identification. As we know, there are no previous studies on bacterial analysis of air quality using this type of filters with samples of PM₁₀ or TSP.

2. Methodology

Three sampling periods were analyzed during 2011; February 8 to 26, April 27 to May 15 and August 25 to September 12 at four different sampling stations. Samples were obtained every six days at each sampling station following the NOM protocol. For the PM₁₀ a Wedding and Associated Brand high volume sampler was employed using quartz filters (Whatman), whereas for TSP sampling, a high volume sampler TSP model TE-5000 with fiber glass filter was used. The size of the filter was 20.3 x 25.4 cm. For PM₁₀, the sample had a continuous flow of 1.02 to 1.24 m³/min, and 1.1 to 1.7 m³/min for TSP, for 24 consecutive hours. The samplers used are property of Municipal Institute of Ecology and employees through the program assessment of air quality.

2.1. Sample Locations

High volume samples were strategically located in four sampling stations in the city, being each one of these:

South station: COBACH High School; coordinates 29°02'40.15"N and 110°57'30.46"W. This site is surrounded by residential and commercial offices, high traffic areas and unpaved roads.

Central station: Direction of Inspection and Vigilance Building of the City of Hermosillo; coordinates 29°04'44.01"N and 110°56'31.91"W. Surrounded by residential and commercial offices and high traffic areas.

North station: Sonora State University (UES); 29°07'17.70"N and 110°57'40.12"W. Surrounded by residential and medium traffic areas.

Northwest station: CBTIS 206 High School; located at 29°07'07.25"N and 111°00'23.86"W. Surrounded by residential areas and unpaved roads. **Figure 1** shows the four sampling sites.

2.2. Stabilization of Filters

The filters were stabilized and weighed before and after sampling under the same conditions of temperature and humidity. Filters were kept for 24 hours in a desiccator with humidity indicator. An analytical electronic balance was used, Sartorius Brand model A200S-D1B installed on anti-vibration table. The filters were handled using powder-free gloves.

2.3. Calculation of Particles Concentration PM₁₀ and TSP in the Air

The concentration of particulate matter in air at standard conditions (P = 760mmHg and T = 298 K) is determined by the mass of collected particles and the volume of air suctioned by the equipment during the sampling time (24 hours) following the protocol established in the Code Federal Regulation from United State Environmental Protection Agency [8].

2.3.1. Concentration of PM₁₀

The equation used for calculating PM₁₀ concentration (1):

$$PM_{10} = \frac{(P2 - P1) \times 10^6}{V} \quad (1)$$

where P1= Initial weight; P2 = Final weight; V = Volume of air sampled.

Air sampled volume is calculated by Equation (2):

$$V = Q_{std} \times 60 \times \text{sampling hours} \quad (2)$$

where: Qstd = Standard flow

Air sampled volume

Standard flow was determined by Equation (3):

$$Q_{std} = \frac{Q_a (\text{Average pressure during sampling} \times 298K)}{760 \text{ mmHg} \times \text{Average temperature during sampling}} \quad (3)$$

where: Qa = Current volumetric flow

We obtained the current volumetric flow by using the Look Up Table (manual) where we calculated ratio pressure (Po/Pa) using formula (4):

$$\frac{P_o}{P_a} = 1 - \left(\frac{P_f}{P_a} \right) \quad (4)$$

where

Pa = Standard pressure (760 mmHg).

Pf = Average pressure drop.

where: Pf was determined by Equation (5):

$$P_f = \frac{\text{Inches of H}_2\text{O}}{13.6 \times 25.4} \quad (5)$$

The average equipment pressure drop was determined using data readings taken before and at the end of sampling with water manometer.




Figure 1. Sampling locations.

2.3.2. Concentration of TSP

The equation used for calculating the concentration of TSP (6):

$$TSP = \frac{(P2 - P1) \times 10^6}{V} \quad (6)$$

where P1= Initial weight; P2 = Final weight; V = Volume.

The volume is calculated with equation (7):

$$V = Q_{pm} \cdot t \quad (7)$$

where: Q_{pm} = Standard flow; t = Time.

For calculating Q_{pm} Equation (8) was used:

$$Q_{pm} = \frac{1}{m} \left[\left(I \sqrt{\frac{P_3 T_{pm}}{P_{pm} T_3}} \right) - b \right] \quad (8)$$

where:

M = Slope of the sampler calibration curve.

I = Average flow indicated in the flow recorder (letter Dickson).

P3 = Average flow indicated in the flow recorder (letter Dickson).

T3 = Average temperature during the sampling period (in Kelvin degrees).

Pp_{tn} = Standard pressure (101 kPa or 760 mmHg).

Tp_{tn} = Standard temperature (in Kelvin degrees, 298).

B = Intersection point of the sampler calibration curve line.

2.4. Sampling for Bacteriological Analysis

Samples for analysis were obtained from the filters PM₁₀ and TSP in which small portions were taken with an area of 28.27 mm². The number of samples per filter was calculated taking into consideration that the filter has a sampling area of 41,617.50 mm² and a volume of 1627.20 m³ of air sampled in 24 hours. The unit sample area considered was 28.27 mm² because it is the standard measure. Canon and Roe's formula was used, considering that expected prevalence of bacteria on the filter was 50% and sensitivity of the techniques employed 95%. According to these operations, the number of sampling units was four.

2.5. Bacteriological Analysis

Samples taken from the filters were placed in tubes containing nutritive broth BHI (Brain Heart Infusion) and incubated at 37°C for 24 hours. The following enriched culture mediums were used for sample incubation: Blood and chocolate agar. Colonies were selected according to morphological differentiation and Gram staining was used to differentiate cell morphology. To gram positive cocci, a catalase test was applied to differentiate Micrococaceae and Streptococaceae family. The gram positive cocci catalase negative was cultured in selective medium KF. Gram negative bacilli were cultured in MacConkey agar and the harvested microorganisms were tested with oxidase. The identification of the isolated microorganisms was done by a biochemical test recommended in Bergey's Manual [9] for each particular case.

Colony forming units (CFU) were counted from the total microorganisms in the filters. The concentration was calculated for a sample area of 41,617.50 mm² and a volume of air sampled of 1627.20 m³ in 24 hours and in accordance with the procedures established on NOM [10].

2.6. Statistical Analysis

A SigmaPlot 11 software was used to determine statistical analysis using one way ANOVA.

3. Results

3.1. Respiratory Infections in Hermosillo's Population

Acute respiratory infections are not produced by inert particles, but by a microorganism that is associated to

them. In the city of Hermosillo, ARI's affects mainly people in ages between 1 and 4 years (as shown in **Figure 2**). The most common infections are: *Klebsiella pneumoniae* sub *pneumoniae* and *Pseudomonas*. However, there is no evidence that these bacteria are airborne.

3.2. Concentration of PM₁₀ and TSP

Comparing the PM₁₀ concentrations values found in each of the sampling dates with the value of the maximum permitted concentration established by the Mexican Official Norm [11] which is 120 µg/m³, only the South Station exceeded the permitted concentration during the second sampling period, with a value of 139.71 µg/m³. TSP concentration values obtained in all sampling dates were below the Mexican Official Norms. In **Figure 3**, particles concentration and maximum permissible by NOM is observed.

3.3. Identified Microorganisms

The results of the microbiological analysis, using traditional techniques, revealed the presence of microorganisms associated with respiratory diseases, such as *Klebsiella pneumoniae* sub *pneumoniae* and *Citrobacter sp10* mainly. Besides *Enterococcus faecalis*, other bacteria present in samples were the *Pseudomonas* genus and the *Enterobacteriaceae* family.

In the south and northwest sampling locations, *E. faecalis* was found in 50% of the samples, 100% in the center location and 33% in the north one. In the center and north locations, *Klebsiellapneumoniae* sub *pneumoniae* was found on February 8. *Citrobacter sp.10* was found once in the south and northwest locations, each in different dates. *Butiauxella agrestis* and *Proteus penneri* were found only in the south location. On May 15, *Salmonella paratyphi A* and *Enterobacter sacazakii* were found in the northwest location. **Table 1** shows the microorganisms found and **Figure 4** shows the results of CFU at each station and period analyzed.

3.4. Antibiotic Sensitivity Analysis

Since *E. faecalis* was the microorganism with higher number of appearances, an antibiotic sensitivity analysis was performed with the support of a local laboratory. (LaboratoriosAcuña, Hermosillo, Mexico). Results can be observed in **Table 2**.




Figure 2. Number of ARI's in the city of Hermosillo, México, during the period 2006-2010, Source: Children's Hospital of the State of Sonora (HIES).





Figure 3. PM_{10} concentration in different sampling sites a) First sample, b) Second sample, c) Third sample and TSP d) First sample, e) Second sample, f) Third sample. There is not a statistically significant between sampling dates North ($P = 0.421$), South ($P = 0.201$), Northwest ($P = 0.344$) and Center ($P = 0.154$).

4. Discussion


Air pollution has been widely studied in different countries to identify potential risks of respiratory diseases in human beings. However, the main focus of these studies is about air chemical contamination regarding the presence of carbon monoxide (CO), sulfur dioxide (SO_2), nitrogen oxides (NO_x and NO_y), reduced nitrogen (ammonia, NH_3 ; ammonium, NH_4), ozone (O_3) [12]-[17], heavy metals [18] [19] and particulate matter PM_{10} [12] [20] [21].

Some authors have been analyzing the possible transmission of pathogens through air, but these studies have been targeted to areas where high content of potential pathogens are present, such as water treatment plants and garbage dumps [22]. Few investigations have been done to look for biological contaminants in the air; the most common is the search for particles that cause allergic diseases such as pollen [23] [24].



(a)

■ 08 February
 ■ 15 August
 ■ 18 September



(b)

Figure 4. CFU of mesophylls bacteria obtained at each sampling location at three different periods (a). Sampled area distribution (b). There is not a statistically significant between sampling dates ($P = 0.274$).

Table 1. Microorganisms identified using traditional biochemical tests.

Zones	08-feb	26-feb	27-abr	15-may	25-ago	18-sep
South	<i>Butiaxellaagrestis</i>	<i>E. faecalis</i> <i>Proteus penneri</i>	<i>E. faecalis</i>	<i>Enterobacterias</i>		<i>E. faecalis</i>
		<i>Citrobacter sp. 10</i>				
Center	<i>E. faecalis</i> <i>Klebsiella pneumoniae sub pneumoniae</i>	<i>E. faecalis</i> <i>Pseudomonas</i>	<i>E. faecalis</i> <i>Cedecea sp. 3</i>	<i>E. faecalis</i> <i>Enterobacterias</i>	<i>E. faecalis</i>	<i>E. faecalis</i>
North	<i>E. faecalis</i> <i>Klebsiella pneumoniae sub pneumoniae</i>	<i>Pseudomonas</i>		<i>E. faecalis</i>		
Northwest	<i>Pseudomonas sp.</i>		<i>E. faecalis</i>	<i>E. faecalis</i> <i>Citrobacter sp.10</i> <i>Salmonella paratyphi A</i> <i>Enterobacter sakazakii</i>	<i>E. faecalis</i>	

Table 2. Antibiotic sensitivity test of 6 isolated of *E. faecalis*.

ANTIBIOTIC	1	2	3	4	5	6
Ampicillin	R	S	R	MR	S	S
Azithromycin	MR	MR	MR	R	S	MR
Ciprofloxacin	MR	MR	MR	R	S	MR
Clarithromycin	S	S	R	S	MR	S
Erythromycin	R	S	R	R	R	MR
Nitrofurantoin	R	S	S	S	MR	S
Fosfomicin	R	R	R	R	S	MR
Gentamicin	R	R	R	S	R	S
Llevofloxacin	S	S	MR	MR	S	MR
Norfloxacin	MR	S	S	MR	S	S
Ofloxacin	MR	MR	S	R	MR	MR
Penicillin G	R	S	S	R	S	R
Rifampicin	MR	MR	R	MR	MR	MR
Cotrimoxazole	S	MR	MR	MR	MR	MR
Tetracycline	MR	S	S	S	S	S
Vancomycin	S	S	S	S	S	S

S = Sensitive; MR = Moderately Resistant; R = Resistant.

The concentration of PM₁₀ was analyzed in four different locations in the city of Hermosillo, México. Concentration levels above the Mexican Standard Norms were only observed during May, in the south location. The results indicate that Hermosillo, at least in 2011, did not present mayor problems of pollution for PM₁₀ in the sampled areas.

This investigation also covered the analysis of filters containing particles PM₁₀ and TSP to determine the load and variety of bacteria. The results demonstrated a high bacterial concentration in the city's air. The greatest concentration was observed in February and the lowest in September.

When bacterial concentration was analyzed separately, the lowest number of bacteria was observed during February for the south location and in August for the north location. The highest degree of bacterial air pollution was found in the central area of the city.

Cases of acute respiratory infections (ARI's) in the city occur mainly in younger age groups, as shown in **Figure 2**. The pathogens identified as generators of these ARI's were, *Klebsiella pneumoniae* sub *pneumoniae* and *Pseudomonas*, among others. It was possible to isolate the bacteria found in the air samples; therefore the conclusion is that air is one of the main routes of transmission of pathogens of ARI's.

In the analysis of isolated bacteria, genres that cause health problems were identified. The most vulnerable population groups are children and the elderly. Some of these bacteria such as *Klebsiella pneumoniae* sub *pneumoniae* and *Pseudomonas* sp. are opportunistic pathogens of nosocomial infections (intra hospital), while *Salmonella paratyphi* A, *E. faecalis* and generally Enterobacteriaceae cause intestinal infections when contaminated foods are consumed. Contamination means of these foods can be direct (hands and contaminated water) or, as demonstrated in this work, air.

Isolated colonies of *E. faecalis* were resistant to almost all antibiotics and in most cases; patients receiving treatment for this bacteria will generally not be effective. According to the presented results, vancomycin is the only antibiotic recommended to treat affected patients by this type of bacteria. Due to the toxicity level of this antibiotic, it is recommended to use it as a last option treatment.

This is the first report of systematic measurements looking for bacterial air contamination. The results shown in this paper suggests that respiratory tract pathogenic bacteria can be transported through the air. Specific pathogens research such as *Mycobacterium tuberculosis*, *Streptococcus pneumoniae*, *Haemophyllus influenzae* and *coccidioides* sp. is still necessary.

References

- [1] Sepúlveda, J.D. (2004) Morphological Profile and Elemental Chemical of Respirable Particulates on Punctual Emission

- Sources and Environmental Sampling (Perfil Morfológico y Químico Elemental de Partículas Respirables en Fuentes de Emisión Puntual y Muestras Ambientales). Informe Final INE-CENICA, México.
- [2] SSA (2005) Official Mexican Standard NOM-025-SSA1-1993, Environmental Health. Criteria for Assessing the Allowable Limit for the Concentration of Particulate Matter, Permissible Limit for the Concentration of Total Suspended Particles TSP, Particles Less than 10 Microns PM10 and Particles Smaller than 2.5 Micrometers PM2.5 of Ambient Air Quality. Criteria for assessing the Air Quality (SSA 2005) Norma Oficial Mexicana NOM-025-SSA1-1993, Salud Ambiental. Criterios para Evaluar el valor Límite Permissible para la Concentración de Material Particulado. Valor Límite Permissible para la Concentración de Partículas Suspendingas Totales PST, Partículas Menores de 10 Micrómetros PM10 y Partículas Menores de 2.5 Micrómetros PM2.5 de la Calidad del Aire Ambiente Criterios para Evaluar la calidad del Aire). Diario Oficial de la Federación, México.
 - [3] Cruz, M.E., Gómez, A., Quintero, M. and Varela, J. (2013) Assessment of Air Quality Regarding Total Suspended Particles (TSP) and Heavy Metals (Pb, Cd, Ni, Cu, Cr) in the city of Hermosillo, Sonora, Mexico, during an Annual Period (Evaluación de la Calidad del Aire Respecto de Partículas Suspendingas Totales (PST) y Metales Pesados (Pb, Cd, Ni, Cu, Cr) en la Ciudad de Hermosillo, Sonora, México, Durante un Periodo Anual). *Revista Internacional de Contaminación Ambiental*, **29**, 269-283.
<http://200.77.231.100/work/normas/noms/2005/025ssa12005.pdf>
 - [4] Kappos, A.D., *et al.* (2004) Health Effects of Particles in Ambient Air. *International Journal of Hygiene and Environmental Health*, **207**, 399-407.
 - [5] de la Rosa, M.C., Mosso, M.A. and Ullán, C. (2002) The Air: Habitat and Transmission of Microorganisms via (El Aire: Hábitat y Medio de Transmisión de Microorganismos). *Observatorio Medioambiental*, **5**, 375-402.
 - [6] Reoun, H., Mainelisa, G. and White, L. (2006) Development and Calibration of Real-time PCR for Quantification of Airborne Microorganisms in Air Samples. *Atmospheric Environment*, **40**, 7924-7939.
<http://dx.doi.org/10.1016/j.atmosenv.2006.07.020>
 - [7] Jensen, P.A., Cih, P.E. and Schafer, M.P. (1998) Sampling and Characterization of Bioaerosols. NIOSH Manual of Analytical Methods NIOSH/DPSE, Chapter 2.
 - [8] USEPA (1992) 40 CFR Ch. I Pt. 50, App. B—Reference Method for the Determination of Suspended Particulate Matter in the Atmosphere (High-Volume Method). United States Environmental Protection Agency—Code of Federal Regulations, Washington DC.
 - [9] Breed, R.S., Murray, E.G.D. and Smith, N.R. (1957) *Bergey's Manual Determinative Bacteriology*. 7th Edition, The Williams & Wilkins Company, Baltimore.
 - [10] SSA (Ministry of Health United States of Mexico) (1994) Mexican Official Standard NOM-113-SSA1-1994. Goods and Services. Counting Method for Total Coliform Microorganisms in plate (SSA, Secretaría de Salud Estados Unidos Mexicanos (Norma Oficial Mexicana NOM-113-SSA1-1994. Bienes y Servicios. Método para la Cuenta de Microorganismos Coliformes Totales en Placa). Diario Oficial de la Federación, México.
 - [11] SSA (Ministry of Health United States of Mexico) (1994) Mexican Official Standard NOM-024-SSA1-1993. Environmental Health. Criteria for Assessing Ambient Air Quality with Respect to Total Suspended Particles (TSP). Allowable Value for the Concentration of Total Suspended Particles (TSP) in Ambient Air as a Protection to the Population's Health (SSA, Secretaría de Salud Estados Unidos Mexicanos, Norma Oficial Mexicana NOM-024-SSA1-1993. Salud Ambiental. Criterio para evaluar la calidad del aire ambiente con respecto a partículas suspendingas totales (PST). Valor permissible para la concentración de partículas suspendingas totales (PST) en el aire ambiente como medida de protección a la salud de la población). Diario Oficial de la Federación, México.
 - [12] Aceves, M.A., *et al.* (2012) Analysis of Key Features of Non-Linear Behavior Using Recurrence Plots, Case Study: Urban Pollution in Mexico City. *Journal of Environmental Protection*, **3**, 1147-1160.
<http://dx.doi.org/10.4236/jep.2012.329133>
 - [13] Capilla, C. (2012) Application of Statistical Methods to Assess Carbon Monoxide Pollution Variations within an Urban Area. *International Journal of Geosciences*, **3**, 885-890. <http://dx.doi.org/10.4236/ijg.2012.325090>
 - [14] Naddafi, K., Sowlat, M.H. and Safari, M.H. (2012) Integrated Assessment of Air Pollution in Tehran, Over the Period from September 2008 to September 2009. *Iranian Journal of Public Health*, **41**, 77-86.
 - [15] Kabashi, S., *et al.* (2012) Dynamic Modeling of Air Pollution and Acid Rain from Energy System and Transport in Kosovo. *Open Journal of Air Pollution*, **1**, 82-96. <http://dx.doi.org/10.4236/ojap.2012.13011>
 - [16] Blanchard, C.L., Hidy, G.M., Tanenbaum, S., Edgerton, E.S. and Hartsell, B.E. (2013) The Southeastern Aerosol Research and Characterization (SEARCH) Study: Spatial Variations and Chemical Climatology, 1999-2010. *Journal of the Air & Waste Management Association*, **63**, 260-275. <http://dx.doi.org/10.1080/10962247.2012.749816>
 - [17] Restrepo, C.E., Simonoff, J.S., Thurston, G.D. and Zimmerman, R. (2012) Asthma Hospital Admissions and Ambient Air Pollutant Concentrations in New York City. *Journal of Environmental Protection*, **3**, 1102-1116.
<http://dx.doi.org/10.4236/jep.2012.329129>

- [18] Spangler, J.G. (2012) Diabetes Mortality and Environmental Heavy Metals in North Carolina Counties: An Ecological Study. *Journal of Diabetes Mellitus*, **2**, 369-372. <http://dx.doi.org/10.4236/jdm.2012.24057>
- [19] Monick, M.M., *et al.* (2013) Effects of Eyjafjallajökull Volcanic Ash on Innate Immune System Responses and Bacterial Growth *in Vitro*. *Environmental Health Perspectives*, **121**, 691-698. <http://dx.doi.org/10.1289/ehp.1206004>
- [20] Newth, D. and Gunasekera, D. (2012) An Integrated Agent-Based Framework for Assessing Air Pollution Impacts. *Journal of Environmental Protection*, **3**, 1135-1146. <http://dx.doi.org/10.4236/jep.2012.329132>
- [21] Vivanco, M.G., *et al.* (2012) Impact of the Transboundary Transport of Air Pollutants on Air Quality in Spain. *Journal of Environmental Protection*, **3**, 1167-1175. <http://dx.doi.org/10.4236/jep.2012.329135>
- [22] Kalwasińska, A., Burkowska, A. and Swiontek, M. (In Press) Exposure of Workers of Municipal Landfill Site to Bacterial and Fungal Aerosol. *Clean—Soil, Air and Water*. <http://onlinelibrary.wiley.com/doi/10.1002/clen.201300385/pdf>
- [23] Ghosh, D., *et al.* (2012) Associations between Pollen Counts, Pollutants and Asthma-Related Hospital Admissions in a High-Density Indian Metropolis. *Journal of Asthma*, **49**, 792-799. <http://dx.doi.org/10.3109/02770903.2012.716473>
- [24] Duro, A., Piccione, V. and Zampino, D. (2013) Air Quality Biomonitoring through Pollen Viability of Fabaceae. *Environmental Monitoring and Assessment*, **185**, 3803-3817. <http://dx.doi.org/10.1007/s10661-012-2829-0>