

Examination of a Multiple Universe Version of the Partition Function of the Universe, Based upon Penrose's Cyclic Conformal Cosmology. Leading to Uniform Values of \hbar (Planck's Constant) and Invariant Physical Laws in Each Universe of the "Multiverse"

Andrew Walcott Beckwith

Physics Department, College of Physics, Chongqing University Huxi Campus, Chongqing, China
Email: Rwill9955b@gmail.com, abeckwith@uh.edu

How to cite this paper: Beckwith, A.W. (2016) Examination of a Multiple Universe Version of the Partition Function of the Universe, Based upon Penrose's Cyclic Conformal Cosmology. Leading to Uniform Values of \hbar (Planck's Constant) and Invariant Physical Laws in Each Universe of the "Multiverse". *Journal of High Energy Physics, Gravitation and Cosmology*, 2, 571-580.
<http://dx.doi.org/10.4236/jhepgc.2016.24049>

Received: June 28, 2016

Accepted: September 9, 2016

Published: September 12, 2016

Copyright © 2016 by author and Scientific Research Publishing Inc. This work is licensed under the Creative Commons Attribution International License (CC BY 4.0).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

Abstract

We look at, starting with Shankar's treatment of the partition function, inserting in the data of the modified Heisenberg uncertainty principle as to give a role to the inflaton in the formation of a partition of the universe. The end result will be, even with the existence of a multiverse, *i.e.* simultaneous universes, uniform physical laws throughout the multiple universes.

Keywords

Cyclic Conformal Cosmology (Penrose), Modified HUP

1. What Is Unique about Our Application of Modification of the Penrose Cyclic Cosmology Theory?

We review the modification of the Penrose cyclic conformal cosmology paradigm given in [1] and include in a partition function given in [2], as a way to include in the modified Heisenberg Uncertainty principle, [3] as a way to ascertain the role of the inflaton, as we write it up in using Padmanabhan's reference [4].

Modification of the HUP and included in our representation of the inflaton, in the partition function will then lead to, after we are including in the results from [2] a way

to discuss how to get a uniform value for Planck’s constant, \hbar , in, per cycle creation of new universes.

On the contrary, the supposition is given by Susskind and others, [5] as to up to 10^{100} universes, with only say 10^6 of them surviving due to sufficiently “robust” cosmological values, for stable physical law. The end result is that what we would have instead is a “multiverse” which is dynamic and stable over time. And so we review our present modification of the Penrose cyclic conformal cosmology model to take into account multiple universes.

2. Extending Penrose’s Suggestion of Cyclic Universes, Black Hole Evaporation, and the Embedding Structure Our Universe Is Contained within. This Multiverse Embeds BHs and May Resolve What Appears to Be an Impossible Dichotomy

That there are no fewer than N universes undergoing Penrose “infinite expansion” [1] contained in a mega universe structure. Furthermore, each of the N universes has black hole evaporation, with the Hawking radiation from decaying black holes. If each of the N universes is defined by a partition function, called $\{\Xi_i\}_{i=N}^{i=1}$, then there exist an information ensemble of mixed minimum information correlated as about $10^7 - 10^8$ bits of information per partition function in the set $\{\Xi_i\}_{i=N}^{i=1} \Big|_{\text{before}}$, so minimum information is conserved between a set of partition functions per universe. We are when following this using the notation of [1] while noting that there is a subsequent alteration of the notation used for partition functions.

$$\{\Xi_i\}_{i=N}^{i=1} \Big|_{\text{before}} \equiv \{\Xi_i\}_{i=N}^{i=1} \Big|_{\text{after}} \tag{1}$$

However, there is non-uniqueness of information put into each partition function $\{\Xi_i\}_{i=N}^{i=1}$. Furthermore Hawking radiation from the black holes is collated via a strange attractor collection in the mega universe structure to form a new big bang for each of the N universes represented by $\{\Xi_i\}_{i=N}^{i=1}$. Verification of this mega structure compression and expansion of information with a non-uniqueness of information placed in each of the N universes favors ergodic mixing treatments of initial values for each of N universes expanding from a singularity beginning. The n_f value, will be using (Ng, 2008) $S_{\text{entropy}} \sim n_f$ [6]. How to tie in this energy expression, as in Equation (1) will be to look at the formation of a nontrivial gravitational measure as a new big bang for each of the N universes as by $n(E_i)$. The density of states at a given energy E_i for a partition function (Poplawski, 2011) [7].

$$\{\Xi_i\}_{i=1}^{i=N} \propto \left\{ \int_0^\infty dE_i \cdot n(E_i) \cdot e^{-E_i} \right\}_{i=1}^{i=N} \tag{2}$$

Each of E_i identified with Equation (2) above, are with the iteration for N universes [1]. Then the following holds, namely, from [1].

Claim 1:

$$\frac{1}{N} \cdot \sum_{j=1}^N \Xi_j \Big|_{j\text{-before-nucleation-regime}} \xrightarrow{\text{vacuum-nucleation-transfer}} \Xi_i \Big|_{i\text{-fixed-after-nucleation-regime}} \tag{3}$$

For N number of universes, with each $\Xi_j|_{j\text{-before-nucleation-regime}}$ for $j = 1$ to N being the partition function of each universe just before the blend into the RHS of Equation (3) above for our present universe. Also, each of the independent universes given by $\Xi_j|_{j\text{-before-nucleation-regime}}$ are constructed by the absorption of one to ten million black holes taking in energy. *i.e.* [1]. Furthermore, the main point is similar to what was done in terms of general ergodic mixing.

Claim 2:

$$\Xi_j|_{j\text{-before-nucleation-regime}} \approx \sum_{k=1}^{Max} \tilde{\Xi}_k|_{\text{black-holes-}j\text{th-universe}} \tag{4}$$

What is done in **Claim 1 and Claim 2** is to come up with a protocol as to how a multi dimensional representation of black hole physics enables continual mixing of spacetime largely as a way to avoid the Anthropic principle, as to a preferred set of initial conditions. How can a graviton with a wavelength 10^{-4} the size of the universe interact with a Kere black hole, spatially. Embedding the BH in a multiverse setting may be the only way out.

Claim 1 is particularly important. The idea here is to use what is known as CCC cosmology, which can be thought of as the following.

First. Have a big bang (initial expansion) for the universe. After redshift $z = 10$, a billion years ago, SMBH formation starts. Matter-energy is vacuumed up by the SMBHs, which at a much later date than today (present era) gather up all the matter-energy of the universe and recycles it in a cyclic conformal translation, as follows, namely

$$\begin{aligned} E &= 8\pi \cdot T + \Lambda \cdot g \\ E &= \text{source for gravitational field} \\ T &= \text{mass energy density} \\ g &= \text{gravitational metric} \\ \Lambda &= \text{vacuum energy, rescaled as follows} \end{aligned} \tag{5}$$

$$\Lambda = c_1 \cdot [Temp]^\beta \tag{6}$$

c_1 is, here a constant. Then we have that for consistency in our presentation that the main methodology in the Penrose proposal has been shown in Equation (6) where we are evaluating a change in the metric g_{ab} by a conformal mapping $\hat{\Omega}$ to

$$\hat{g}_{ab} = \hat{\Omega}^2 g_{ab} \tag{7}$$

Penrose’s suggestion has been to utilize the following [1]

$$\hat{\Omega} \xrightarrow{ccc} \hat{\Omega}^{-1} \tag{8}$$

The infall into cosmic black holes has been the main mechanism which the author asserts would be useful for the recycling apparent in Equation (8) above with the caveat that \hbar is kept constant from cycle to cycle as represented by

$$\hbar_{\text{old-cosmology-cycle}} = \hbar_{\text{present-cosmology-cycle}} \tag{9}$$

Equation (9) is to be generalized, as given by a weighing averaging as given by Equa-

tion (3). where the averaging is collated over perhaps thousands of universes, call that number N , with an ergotic mixing of all these universes, with the ergodic mixing represented by Equation (3) to generalize Equation (9) from cycle to cycle.

3. Now for the Mixing Being Put in, and Birkhoff's Ergodic Mixing Theorem

We will, afterwards, do the particulars of the partition function. But before that, we will do the "mixing" of inputs into the Partition function of the Universe, *i.e.* an elaboration on Equation (3) above. To do this, first, look at the following, from [8].

Birkhoff's Ergodic mixing theorem:

Let $T : (X, \mu) \rightarrow (X, \mu)$ be a measure preserving transformation of a probability space $\varphi \in L^1(X, \mu)$.

Then for μ almost every $x \in X$ the following time average exists

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=0}^{n-1} \varphi(T^j(x)) =: \varphi_T(x). \tag{10}$$

In the end, we need to have a way to present how the bona fides of Equation (9) can be established, and the averaging of both Equation (10) and Equation (4) above need to be put to a consistent general treatment for an invariant \hbar for cycle to cycle, of cosmological creation.

To do this, we also refer to the generalized treatment of, from [9]

$$\begin{aligned} \text{Energy (average)} &= kT^2 \left(\frac{\partial \ln Z(T)}{\partial T} \right) \sim \hbar \omega \\ \Leftrightarrow \hbar(\text{averaged-value}) &= \frac{k_B T^2}{\omega} \left(\frac{\partial \ln Z(T)}{\partial T} \right) \\ &= \frac{k_B T^2}{\omega} \left(\frac{\partial \ln Z(T)}{\partial T} \right) = \frac{k_B \lambda T^2}{2\pi c \omega} \left(\frac{\partial \ln Z(T)}{\partial T} \right) \\ \Leftrightarrow \hbar(\text{averaged-value}) &= \frac{k_B \lambda T^2}{2\pi c} \left(\frac{\partial \ln Z(T)}{\partial T} \right) \end{aligned} \tag{11}$$

Having said, that, the remaining constraint is to come up with a suitably averaged value of the Partition function in the above work. Our averaging eventually will have to be reconciled with the Birkhoff Ergodic Mixing theorem.

4. How to Average out the Planck's Constant, Using Partition Function Given in Equation (11)

We begin with what is given in Shankar's treatment of the partition function of [10] as given by

$$Z(T) = Z(\beta) = \int dr \cdot \sqrt{\frac{1}{2\pi}} \cdot \frac{\exp[-\beta V(r)]}{\Delta r}. \tag{12}$$

Using, for Pre Planckian space-time the approximation of [10]

$$\begin{aligned}
 Z(\beta) &\approx \int dr \cdot \sqrt{\frac{1}{2\pi}} \cdot \frac{\exp[-\beta V(r)]}{\Delta r} \sim \Delta r \sqrt{\frac{1}{2\pi}} \cdot \frac{\exp[-\beta V(r_{\text{midpoint}})]}{\Delta r} \\
 &\Leftrightarrow Z(\beta) \approx \sqrt{\frac{1}{2\pi}} \cdot \exp[-\beta V(r_{\text{midpoint}})].
 \end{aligned}
 \tag{13}$$

Approximate using Beckwith's treatment of the HUP, in Pre Planckian space-time [3]

$$\begin{aligned}
 Z(\beta) &\approx \int dr \cdot \sqrt{\frac{1}{2\pi}} \cdot \frac{\exp[-\beta V(r)]}{\Delta r} \sim \Delta r \sqrt{\frac{1}{2\pi}} \cdot \frac{\exp[-\beta V(r_{\text{midpoint}})]}{\Delta r} \\
 &\Leftrightarrow Z(\beta) \approx \sqrt{\frac{1}{2\pi}} \cdot \exp[-\beta V(r_{\text{midpoint}})] \\
 V(r_{\text{midpoint}}) &\sim \Delta E \sim \frac{\hbar}{\delta t \cdot \delta g_{tt}} \sim \frac{\hbar}{\delta t \cdot (a_{\text{min}})^2 \phi} \\
 &\Leftrightarrow Z(\beta) \approx \sqrt{\frac{1}{2\pi}} \cdot \exp\left[-\beta \cdot \frac{\hbar}{\delta t \cdot (a_{\text{min}})^2 \phi}\right].
 \end{aligned}
 \tag{14}$$

Put in now the value of the inflaton given by Padmanbhan, [4] as for

$$\begin{aligned}
 a &\approx a_{\text{min}} t^\gamma \\
 \Leftrightarrow \phi &\approx \sqrt{\frac{\gamma}{4\pi G}} \cdot \ln\left\{\sqrt{\frac{8\pi G V_0}{\gamma \cdot (3\gamma - 1)}} \cdot t\right\} \\
 \Leftrightarrow V &\approx V_0 \cdot \exp\left\{-\sqrt{\frac{16\pi G}{\gamma}} \cdot \phi(t)\right\}
 \end{aligned}
 \tag{15}$$

Put in the value for the inflaton as given in Equation (15) into the partition function of Equation (14)

$$\begin{aligned}
 Z(\beta) &\approx \sqrt{\frac{1}{2\pi}} \cdot \exp\left[-\beta \cdot \frac{\hbar}{\delta t \cdot (a_{\text{min}})^2 \phi}\right] = \sqrt{\frac{1}{2\pi}} \cdot \exp\left[-\beta \cdot \frac{\hbar}{\delta t \cdot (a_{\text{min}})^2 \sqrt{\frac{\gamma}{4\pi G}} \cdot \ln\left\{\sqrt{\frac{8\pi G V_0}{\gamma \cdot (3\gamma - 1)}} \cdot t\right\}}\right] \\
 &= \sqrt{\frac{1}{2\pi}} \cdot \exp\left[\beta \cdot \frac{\hbar}{\delta t \cdot (a_{\text{min}})^2 \sqrt{\frac{\gamma}{4\pi G}} \cdot \ln\left\{1/\left(\sqrt{\frac{8\pi G V_0}{\gamma \cdot (3\gamma - 1)}} \cdot t\right)\right\}}\right] \\
 &\& \ln\left\{1/\left(\sqrt{\frac{8\pi G V_0}{\gamma \cdot (3\gamma - 1)}} \cdot t\right)\right\} = \left[\frac{\left\{1/\left(\sqrt{\frac{8\pi G V_0}{\gamma \cdot (3\gamma - 1)}} \cdot t\right)\right\}^{-1}}{\left\{1/\left(\sqrt{\frac{8\pi G V_0}{\gamma \cdot (3\gamma - 1)}} \cdot t\right)\right\}}\right] + \frac{1}{2} \cdot \left[\frac{\left\{1/\left(\sqrt{\frac{8\pi G V_0}{\gamma \cdot (3\gamma - 1)}} \cdot t\right)\right\}^{-1}}{\left\{1/\left(\sqrt{\frac{8\pi G V_0}{\gamma \cdot (3\gamma - 1)}} \cdot t\right)\right\}}\right]^2 + \dots
 \end{aligned}
 \tag{16}$$

Then using Shankar, [9], we will take the result so Equation (16) and then from there, use them in Equation (15) and also Equation (14). The end result is a massive cancella-

tion of the terms, as to obtain Equation (17) below

$$\begin{aligned}
 \frac{\partial \ln Z(T)}{\partial T} &= \frac{1}{Z(T)} \cdot \frac{\partial Z(T)}{\partial T} \\
 &= \frac{-\hbar}{\delta t \cdot (a_{\min})^2 \sqrt{\frac{\gamma}{4\pi G}} \cdot \ln \left\{ 1 / \left(\sqrt{\frac{8\pi G V_0}{\gamma \cdot (3\gamma - 1)}} \cdot t \right) \right\}} \cdot \frac{1}{k_B T^2} \\
 &= \frac{\hbar}{\delta t \cdot (a_{\min})^2 \sqrt{\frac{\gamma}{4\pi G}} \cdot \ln \left\{ \left(\sqrt{\frac{8\pi G V_0}{\gamma \cdot (3\gamma - 1)}} \cdot t \right) \right\}} \cdot \frac{1}{k_B T^2} \tag{17} \\
 &\sim \frac{\hbar}{(\delta t)^2 \cdot (a_{\min})^2 \sqrt{\frac{8\pi G V_0}{\gamma \cdot (3\gamma - 1)}} \sqrt{\frac{\gamma}{4\pi G}}} \cdot \frac{1}{k_B T^2}
 \end{aligned}$$

Then by use of Equation (11) we obtain

$$\begin{aligned}
 \hbar(\text{averaged-value}) &= \frac{k_B \lambda T^2}{2\pi c} \left(\frac{\partial \ln Z(T)}{\partial T} \right) \\
 &\sim \frac{\lambda}{2\pi c} \frac{\hbar}{(\delta t)^2 \cdot (a_{\min})^2 \sqrt{\frac{8\pi G V_0}{\gamma \cdot (3\gamma - 1)}} \sqrt{\frac{\gamma}{4\pi G}}} \tag{18} \\
 &\Leftrightarrow \frac{\lambda}{2\pi c} \frac{1}{(\delta t)^2 \cdot (a_{\min})^2 \sqrt{\frac{8\pi G V_0}{\gamma \cdot (3\gamma - 1)}} \sqrt{\frac{\gamma}{4\pi G}}} \sim 1.
 \end{aligned}$$

This is the baseline of the constraint which will make Planck’s constant, a constant per universe creation cycle. As given by Equation (9). *i.e.* Equation (9) is confirmed by Equation (18). We will next then go to how this ties into Equation (10) above, via use of averaging is affecting the choice of the inputs into Equation (18) above. Doing this will allow investigation as to how to falsify the Birkhoff Ergodic mixing theorem as mentioned next.

5. Applying the Birkhoff Ergodic Averaging Equation (10) to the Inputs into Equation (18)

To do this, we specifically look at the wavelength, namely, applying [8] to a wavelength λ . One over the wavelength is proportional to frequency, so if we have the wave length, as represented by the following situation, with invariance set in stone. Here we are assuming that the formulation is, as follows: With N the number of recycled “universes”

$$\begin{aligned}
 \varphi(T^j(x)) &= \lambda \\
 \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{j=0}^{N-1} \varphi(T^j(x)) &= \lambda \tag{19}
 \end{aligned}$$

i.e. the averaging by the Burkhoff theorem implies that there is a critical invariance. And this invariance should be linked, then, to the diameter of a nonsingular bounce

point. A nonsingular bounce, *i.e.*, beginning of an expansion of a new universe is the main point of [11]. Furthermore, we have that if we look at applying the insights of [12] [13] we obtain

$$\begin{aligned}
 V_0 &\sim m = \text{mass}(\text{transferred-space-time}) \\
 &= m_{\text{graviton}} \cdot \text{Entropy} = m_{\text{graviton}} n(\text{entropy}) \\
 &\& n(\text{entropy}) \sim 10^{10} - 10^{20} \\
 &\& m_{\text{graviton}} \sim 10^{-62} \text{ grams} \\
 &\& (a_{\text{min}})^2 \sim 10^{-110} \\
 &\& \delta t \sim 10^{-\beta} t_{\text{Planck}}; 0 < \beta < 10
 \end{aligned}
 \tag{20}$$

We will try to show, in a later date that these are invariant per cycle, but the upshot is that if there is a natural fit, as to Equation (19) and if λ is fixed as an invariant per cycle, given by 19, then the invariance of \hbar per cycle is then maintained.

6. Conclusion: Implications of the Invariance of \hbar : Uniform Physics Laws per Universe, and Not the 10^{1000} Created Universes with Only Say 10^{10} Surviving through a Cosmic Cycle

In a word this demolishes the program of the cosmic landscape of string theory [5], and gives credence to the possibility of an invariant multiverse, which would not be collapsing.

If this is confirmed, experimentally, it will do much to reduce what has been at times a post modern fragmentation of basic physics inquiry and to have physics, with a uniform set of laws, regardless of whether there were many worlds, or just one, in terms of one universe, or many universes, and as well as allow investigation of the information theory approach of [14] to event horizons and early universe cosmology. How [14] could influence a choice of partition functions is given in this paper's Appendix.

Acknowledgements

This work is supported in part by National Nature Science Foundation of China grant No. 11375279.

References

- [1] Penrose, R. (2010) *Cycles of Time: An Extraordinary New View of the Universe*. The Bodley Head, the Random House Group, Ltd., London.
- [2] Beckwith, A. (2014) Analyzing Black Hole Super-Radiance Emission of Particles/Energy from a Black Hole as a Gedanken Experiment to Get Bounds on the Mass of a Graviton. *Advances in High Energy Physics*, **2014**, Article ID: 230713. <http://www.hindawi.com/journals/ahep/2014/230713/> <http://dx.doi.org/10.1155/2014/230713>
- [3] Beckwith, A. (2016) Gedanken Experiment for Refining the Unruh Metric Tensor Uncertainty Principle via Schwarzschild Geometry and Planckian Space-Time with Initial Non-

- zero Entropy and Applying the Riemannian-Penrose Inequality and Initial Kinetic Energy for a Lower Bound to Graviton Mass (Massive Gravity). *Journal of High Energy Physics, Gravitation and Cosmology*, **2**, 106-124. <http://dx.doi.org/10.4236/jhepgc.2016.21012>
- [4] Padmanabhan, T. (2005) Understanding Our Universe, Current Status and Open Issues. In: Ashatekar, A., Ed., *100 Years of Relativity, Space-Time Structure. Einstein and Beyond*, World Scientific Publishing Co. Pte. Ltd., Singapore, 175-204. <http://arxiv.org/abs/gr-qc/0503107>
http://dx.doi.org/10.1142/9789812700988_0007
- [5] Susskind, L. (2006) *The Cosmic Landscape: String Theory and the Illusion of Intelligent Design*. Little, Brown, and Company, Time Warner Book Group, New York.
- [6] Ng, Y.J. (2007) Holographic Foam, Dark Energy and Infinite Statistics. *Physics Letters B*, **657**, 10-14. <http://dx.doi.org/10.1016/j.physletb.2007.09.052>
- [7] Poplawski, N. (2011) Cosmological Constant from QCD Vacuum and Torsion. *Annalen der Physik*, **523**, 291-295. <http://dx.doi.org/10.1002/andp.201000162>
- [8] Katok, A. and Hasselblatt, B. (1999) *Introduction to the Modern Theory of Dynamical Systems*. Cambridge University Press, Cambridge.
- [9] Shankar, R. (2014) *Fundamentals of Physics, Mechanics, Relativity and Thermodynamics*. Yale University Press, Yale.
- [10] Shankar, R. (1994) *Principles of Quantum Mechanics*. 2nd Edition, Springer Verlag, Heidelberg. <http://dx.doi.org/10.1007/978-1-4757-0576-8>
- [11] Beckwith, A. (2016) Gedanken Experiment for Delineating the Regime for the Start of Quantum Effects, and Their End, Using Turok's Perfect Bounce Criteria and Radii of a Bounce Maintaining Quantum Effects, as Delineated by Haggard and Rovelli. *Journal of High Energy Physics, Gravitation and Cosmology*, **2**, 287-292. <http://dx.doi.org/10.4236/jhepgc.2016.23024>
- [12] Corda, C. (2012) Gravity's Primordial Breath. *Electronic Journal of Theoretical Physics*, **9**, 1-10. <http://www.ejtp.com/articles/ejtpv9i26p1.pdf>
- [13] Gao, C. (2012) A Model of Nonsingular Universe. *Entropy*, **14**, 1296-1305. <http://dx.doi.org/10.3390/e14071296>
- [14] Lee, J.W. (2010) Quantum Mechanics Emerges from Information Theory Applied to Causal Horizons. *Foundations of Physics*, **41**, 744-753. <http://dx.doi.org/10.1007/s10701-010-9514-3>

Appendix: Highlights of J.-W. Lee's Paper [14]

The following formulation is to highlight how entropy generation blends in with quantum mechanics, and how the breakdown of some of the assumptions used in Lee's paper coincide with the growth of degrees of freedom. What is crucial to Lee's formulation, is Rindler geometry, not the curved space formulation of initial universe conditions. First of all, [14] (Lee, 2010),

“Considering all these recent developments, it is plausible that quantum mechanics and gravity has information as a common ingredient, and information is the key to explain the strange connection between two. If gravity and Newton mechanics can be derived by considering information at Rindler horizons, it is natural to think quantum mechanics might have a similar origin. In this paper, along this line, it is suggested that quantum field theory (QFT) and quantum mechanics can be obtained from information theory applied to causal (Rindler) horizons, and that quantum randomness arises from information blocking by the horizons.”

To start this we look at the Rindler partition function, as by [14] (Lee, 2010)

$$Z_R = \sum_{i=1}^n \exp[-\beta H(x_i)] = \text{Trace} \cdot (\exp[-\beta H]). \quad (\text{A.1})$$

As stated by Lee [14], we expect Z_R to be equal to the quantum mechanical partition function of a particle with mass m in Minkowski space time. Furthermore, there exists the datum that: Lee made an equivalence between Equation (A1) and [14] (Lee, 2010)

$$Z_Q = N_i \int \delta x \cdot \exp\left[\frac{-i}{\hbar} \cdot I(x_i)\right] \quad (\text{A2})$$

where $I(x_i)$ is the action “integral” for each path x_i , leading to a wave function for each path? x_i

$$\psi \sim \exp\left[\frac{-i}{\hbar} \cdot I(x_i)\right] \quad (\text{A3})$$

If we do a rescale $\hbar = 1$, then the above wave equation can lead to a Schrodinger equation.

The example given by (Lee, 2010) is that there is a Hamiltonian for which

$$H(\phi) = \int d^3x \cdot \left\{ \frac{1}{2} \cdot \left(\frac{\partial \phi}{\partial r} \right)^2 + \frac{1}{2} \cdot (\nabla \phi)^2 + V(\phi) \right\}. \quad (\text{A4})$$

Here, V is a potential, and ϕ can have arbitrary values before measurement, and to a degree, Z represent uncertainty in measurement. In Rindler co-ordinates, $H \rightarrow H_R$, in co-ordinates (η, r, x_2, x_3) with proper time variance $ard\eta$ then

$$H_R(\phi) = \int dr dx_{\perp} ar \cdot \left\{ \frac{1}{2} \cdot \left(\frac{\partial \phi}{\partial r} \right)^2 + \frac{1}{2} \cdot \left(\frac{\partial \phi}{ar \partial \eta} \right)^2 + \frac{1}{2} \cdot (\nabla_{\perp} \phi)^2 + V(\phi) \right\}. \quad (\text{A5})$$

Here, the \perp is a plane orthogonal to the (η, r) plane. If so then

$$Z = \text{tr exp}[-\beta H] \mapsto Z_R = \text{tr exp}[-\beta H_R]. \quad (\text{A6})$$

Now, for the above situation, the following are equivalent.

- 1) Z_R Thermal partition function is from information loss about field beyond the Rindler Horizon.
- 2) QFT formation is equivalent to purely information based statistical treatment suggested in this paper.
- 3) QM emerges from information theory emerging from Rindler co-ordinate.

Lee also forms a Euclidian version for the following partition function, if $I_E(x_i)$ is the Euclidian action for the scalar field in the initial frame. *i.e.*

$$Z_Q^E = N_1 \int \phi x \cdot \exp\left[\frac{-i}{\hbar} \cdot I_E(x_i)\right] \quad (\text{A7})$$

There exist analytic continuation of $\tilde{t} \mapsto it$ leading to $Z_Q^E \mapsto Z_Q =$ Usual zero temperature QM partition function of Z_Q for ϕ fields.

Important Claim: The following are equivalent.

- 1) Z_R and Z_Q are obtained by analytic continuation from Z_Q^E .
- 2) Z_R and Z_Q are equivalent.



Submit or recommend next manuscript to SCIRP and we will provide best service for you:

Accepting pre-submission inquiries through Email, Facebook, LinkedIn, Twitter, etc.

A wide selection of journals (inclusive of 9 subjects, more than 200 journals)

Providing 24-hour high-quality service

User-friendly online submission system

Fair and swift peer-review system

Efficient typesetting and proofreading procedure

Display of the result of downloads and visits, as well as the number of cited articles

Maximum dissemination of your research work

Submit your manuscript at: <http://papersubmission.scirp.org/>