



# Weak Insertion of a Continuous Function between Two Comparable $\alpha$ -Continuous (C-Continuous) Functions\*

**Majid Mirmiran**

Department of Mathematics, University of Isfahan, Isfahan, Iran

Email: mirmir@sci.ui.ac.ir

Received 18 February 2016; accepted 4 March 2016; published 9 March 2016

Copyright © 2016 by author and OALib.

This work is licensed under the Creative Commons Attribution International License (CC BY).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

---

## Abstract

**A sufficient condition in terms of lower cut sets is given for the insertion of a continuous function between two comparable real-valued functions.**

## Keywords

**Weak Insertion, Strong Binary Relation, C-Open Set, Semi-Preopen Set,  $\alpha$ -Open Set, Lower Cut Set**

**Subject Areas: Topology**

---

## 1. Introduction

The concept of a  $C$ -open set in a topological space was introduced by E. Hatir, T. Noiri and S. Yksel in 1996 [1]. The authors define a set  $S$  to be a  $C$ -open set if  $S = U \cap A$ , where  $U$  is open and  $A$  is semi-preclosed. A set  $S$  is a  $C$ -closed set if its complement is  $C$ -open set or equivalently if  $S = U \cup A$ , where  $U$  is closed and  $A$  is semi-preopen. The authors show that a subset of a topological space is open if and only if it is an  $\alpha$ -open set and a  $C$ -open set. This enable them to provide the following decomposition of continuity: a function is continuous if and only if it is  $\alpha$ -continuous and  $C$ -continuous.

Recall that a subset  $A$  of a topological space  $(X, \tau)$  is called  $\alpha$ -open if  $A$  is the difference of an open and a nowhere dense subset of  $X$ . A set  $A$  is called  $\alpha$ -closed if its complement is  $\alpha$ -open or equivalently if  $A$  is union of a closed and a nowhere dense set. Sets which are dense in some regular closed subspace are called semi-preopen or  $\beta$ -open. A set is semi-preclosed or  $\beta$ -closed if its complement is semi-preopen or  $\beta$ -open.

The concept of a set  $A$  was  $\beta$ -open if and only if  $A \subseteq Cl(Int(Cl(A)))$  was introduced by J. Dontchev in 1998 [2].

---

\*This work was supported by University of Isfahan and Centre of Excellence for Mathematics (University of Isfahan).

Recall that a real-valued function  $f$  defined on a topological space  $X$  was called  $A$ -continuous if the preimage of every open subset of  $\mathbb{R}$  belongs to  $A$ , where  $A$  was a collection of subset of  $X$  and this the concept was introduced by M. Przemski in 1993 [3]. Most of the definitions of function used throughout this paper are consequences of the definition of  $A$ -continuity. However, for unknown concepts, the reader might refer to papers introduced by J. Dontchev in 1995 [4], M. Ganster and I. Reilly in 1990 [5].

Hence, a real-valued function  $f$  defined on a topological space  $X$  is called  $C$ -continuous (resp.  $\alpha$ -continuous) if the preimage of every open subset of  $\mathbb{R}$  is  $C$ -open (resp.  $\alpha$ -open) subset of  $X$ .

Results of Katětov in 1951 [6] and in 1953 [7] concerning binary relations and the concept of an indefinite lower cut set for a real-valued function, which was due to Brooks in 1971 [8], were used in order to give necessary and sufficient conditions for the strong insertion of a continuous function between two comparable real-valued functions.

If  $g$  and  $f$  are real-valued functions defined on a space  $X$ , we write  $g \leq f$  in case  $g(x) \leq f(x)$  for all  $x$  in  $X$ .

The following definitions were modifications of conditions considered in paper introduced by E. Lane in 1976 [9].

A property  $P$  defined relative to a real-valued function on a topological space is a  $c$ -property provided that any constant function has property  $P$  and provided that the sum of a function with property  $P$  and any continuous function also has property  $P$ . If  $P_1$  and  $P_2$  are  $c$ -property, the following terminology is used: A space  $X$  has the *weak  $c$ -insertion property* for  $(P_1, P_2)$  if and only if for any functions  $g$  and  $f$  on  $X$  such that  $g \leq f$ ,  $g$  has property  $P_1$  and  $f$  has property  $P_2$ , then there exists a continuous function  $h$  such that  $g \leq h \leq f$ .

In this paper, it is given a sufficient condition for the weak  $c$ -insertion property. Also several insertion theorems are obtained as corollaries of this result.

## 2. The Main Result

Before giving a sufficient condition for insertability of a continuous function, the necessary definitions and terminology are stated.

Let  $(X, \tau)$  be a topological space, the family of all  $\alpha$ -open,  $\alpha$ -closed,  $C$ -open and  $C$ -closed will be denoted by  $\alpha O(X, \tau)$ ,  $\alpha C(X, \tau)$ ,  $CO(X, \tau)$  and  $CC(X, \tau)$ , respectively.

**Definition 2.1.** Let  $A$  be a subset of a topological space  $(X, \tau)$ . Respectively, we define the  $\alpha$ -closure,  $\alpha$ -interior,  $C$ -closure and  $C$ -interior of a set  $A$ , denoted by  $\alpha Cl(A)$ ,  $\alpha Int(A)$ ,  $C Cl(A)$  and  $C Int(A)$  as follows:

$$\begin{aligned}\alpha Cl(A) &= \bigcap \{F : F \supseteq A, F \in \alpha C(X, \tau)\} \\ \alpha Int(A) &= \bigcup \{O : O \subseteq A, O \in \alpha O(X, \tau)\} \\ C Cl(A) &= \bigcap \{F : F \supseteq A, F \in CC(X, \tau)\} \\ &\text{and} \\ C Int(A) &= \bigcup \{O : O \subseteq A, O \in CO(X, \tau)\}.\end{aligned}$$

Respectively, we have  $\alpha Cl(A)$ ,  $C Cl(A)$  are  $\alpha$ -closed, semi-preclosed and  $\alpha Int(A)$ ,  $C Int(A)$  are  $\alpha$ -open, semi-preopen.

The following first two definitions are modifications of conditions considered in [6] [7].

**Definition 2.2.** If  $\rho$  is a binary relation in a set  $S$  then  $\bar{\rho}$  is defined as follows:  $x \bar{\rho} y$  if and only if  $y \rho v$  implies  $x \rho v$  and  $u \rho x$  implies  $u \rho y$  for any  $u$  and  $v$  in  $S$ .

**Definition 2.3.** A binary relation  $\rho$  in the power set  $P(X)$  of a topological space  $X$  is called a *strong binary relation* in  $P(X)$  in case  $\rho$  satisfies each of the following conditions:

- 1) If  $A_i \rho B_j$  for any  $i \in \{1, \dots, m\}$  and for any  $j \in \{1, \dots, n\}$ , then there exists a set  $C$  in  $P(X)$  such that  $A_i \rho C$  and  $C \rho B_j$  for any  $i \in \{1, \dots, m\}$  and any  $j \in \{1, \dots, n\}$ .
- 2) If  $A \subseteq B$ , then  $A \bar{\rho} B$ .
- 3) If  $A \rho B$ , then  $Cl(A) \subseteq B$  and  $A \subseteq Int(B)$ .

The concept of a lower indefinite cut set for a real-valued function was defined [8] as follows:

**Definition 2.4.** If  $f$  is a real-valued function defined on a space  $X$  and if

$\{x \in X : f(x) < \ell\} \subseteq A(f, \ell) \subseteq \{x \in X : f(x) \leq \ell\}$  for a real number  $\ell$ , then  $A(f, \ell)$  is called a *lower indefinite cut set* in the domain of  $f$  at the level  $\ell$ .

We now give the following main result:

**Theorem 2.1.** Let  $g$  and  $f$  be real-valued functions on a topological space  $X$  with  $g \leq f$ . If there exists a strong binary relation  $\rho$  on the power set of  $X$  and if there exist lower indefinite cut sets  $A(f, t)$  and  $A(g, t)$  in the domain of  $f$  and  $g$  at the level  $t$  for each rational number  $t$  such that if  $t_1 < t_2$  then  $A(f, t_1) \rho A(g, t_2)$ , then there exists a continuous function  $h$  defined on  $X$  such that  $g \leq h \leq f$ .

**Proof.** Let  $g$  and  $f$  be real-valued functions defined on  $X$  such that  $g \leq f$ . By hypothesis there exists a strong binary relation  $\rho$  on the power set of  $X$  and there exist lower indefinite cut sets  $A(f, t)$  and  $A(g, t)$  in the domain of  $f$  and  $g$  at the level  $t$  for each rational number  $t$  such that if  $t_1 < t_2$  then  $A(f, t_1) \rho A(g, t_2)$ .

Define functions  $F$  and  $G$  mapping the rational numbers  $\mathbb{Q}$  into the power set of  $X$  by  $F(t) = A(f, t)$  and  $G(t) = A(g, t)$ . If  $t_1$  and  $t_2$  are any elements of  $\mathbb{Q}$  with  $t_1 < t_2$ , then  $F(t_1) \bar{\rho} F(t_2)$ ,  $G(t_1) \bar{\rho} G(t_2)$ , and  $F(t_1) \rho G(t_2)$ . By Lemmas 1 and 2 of [7] it follows that there exists a function  $H$  mapping  $\mathbb{Q}$  into the power set of  $X$  such that if  $t_1$  and  $t_2$  are any rational numbers with  $t_1 < t_2$ , then  $F(t_1) \rho H(t_2)$ ,  $H(t_1) \rho H(t_2)$  and  $H(t_1) \rho G(t_2)$ .

For any  $x$  in  $X$ , let  $h(x) = \inf \{t \in \mathbb{Q} : x \in H(t)\}$ .

We first verify that  $g \leq h \leq f$ : If  $x$  is in  $H(t)$  then  $x$  is in  $G(t')$  for any  $t' > t$ ; since  $x$  is in  $G(t') = A(g, t')$  implies that  $g(x) \leq t'$ , it follows that  $g(x) \leq t$ . Hence  $g \leq h$ . If  $x$  is not in  $H(t)$ , then  $x$  is not in  $F(t')$  for any  $t' < t$ ; since  $x$  is not in  $F(t') = A(f, t')$  implies that  $f(x) > t'$ , it follows that  $f(x) \geq t$ . Hence  $h \leq f$ .

Also, for any rational numbers  $t_1$  and  $t_2$  with  $t_1 < t_2$ , we have  $h^{-1}(t_1, t_2) = \text{Int}(H(t_2)) \setminus \text{Cl}(H(t_1))$ . Hence  $h^{-1}(t_1, t_2)$  is an open subset of  $X$ , i.e.,  $h$  is a continuous function on  $X$ . ■

The above proof used the technique of proof of Theorem 1 of [6].

### 3. Applications

The abbreviations  $\alpha c$  and  $Cc$  are used for  $\alpha$ -continuous and  $C$ -continuous, respectively.

**Corollary 3.1.** If for each pair of disjoint  $\alpha$ -closed (resp.  $C$ -closed) sets  $F_1, F_2$  of  $X$ , there exist open sets  $G_1$  and  $G_2$  of  $X$  such that  $F_1 \subseteq G_1$ ,  $F_2 \subseteq G_2$  and  $G_1 \cap G_2 = \emptyset$  then  $X$  has the weak  $c$ -insertion property for  $(\alpha c, \alpha c)$  (resp.  $(Cc, Cc)$ ).

**Proof.** Let  $g$  and  $f$  be real-valued functions defined on the  $X$ , such that  $f$  and  $g$  are  $\alpha c$  (resp.  $Cc$ ), and  $g \leq f$ . If a binary relation  $\rho$  is defined by  $A \rho B$  in case  $\alpha \text{Cl}(A) \subseteq \alpha \text{Int}(B)$  (resp.  $C \text{Cl}(A) \subseteq C \text{Int}(B)$ ), then by hypothesis  $\rho$  is a strong binary relation in the power set of  $X$ . If  $t_1$  and  $t_2$  are any elements of  $\mathbb{Q}$  with  $t_1 < t_2$ , then

$$A(f, t_1) \subseteq \{x \in X : f(x) \leq t_1\} \subseteq \{x \in X : g(x) < t_2\} \subseteq A(g, t_2);$$

since  $\{x \in X : f(x) \leq t_1\}$  is an  $\alpha$ -closed (resp.  $C$ -closed) set and since  $\{x \in X : g(x) < t_2\}$  is an  $\alpha$ -open (resp.  $C$ -open) set, it follows that  $\alpha \text{Cl}(A(f, t_1)) \subseteq \alpha \text{Int}(A(g, t_2))$  (resp.  $C \text{Cl}(A(f, t_1)) \subseteq C \text{Int}(A(g, t_2))$ ). Hence  $t_1 < t_2$  implies that  $A(f, t_1) \rho A(g, t_2)$ . The proof follows from Theorem 2.1. ■

**Corollary 3.2.** If for each pair of disjoint  $\alpha$ -closed (resp.  $C$ -closed) sets  $F_1, F_2$ , there exist open sets  $G_1$  and  $G_2$  such that  $F_1 \subseteq G_1$ ,  $F_2 \subseteq G_2$  and  $G_1 \cap G_2 = \emptyset$  then every  $\alpha$ -continuous (resp.  $C$ -continuous) function is continuous.

**Proof.** Let  $f$  be a real-valued  $\alpha$ -continuous (resp.  $C$ -continuous) function defined on the  $X$ . Set  $g = f$ , then by Corollary 3.1, there exists a continuous function  $h$  such that  $g = h = f$ . ■

**Corollary 3.3.** If for each pair of disjoint subsets  $F_1, F_2$  of  $X$ , such that  $F_1$  is  $\alpha$ -closed and  $F_2$  is  $C$ -closed, there exist open subsets  $G_1$  and  $G_2$  of  $X$  such that  $F_1 \subseteq G_1$ ,  $F_2 \subseteq G_2$  and  $G_1 \cap G_2 = \emptyset$  then  $X$  have the weak  $c$ -insertion property for  $(\alpha c, Cc)$  and  $(Cc, \alpha c)$ .

**Proof.** Let  $g$  and  $f$  be real-valued functions defined on the  $X$ , such that  $g$  is  $\alpha c$  (resp.  $Cc$ ) and  $f$  is  $Cc$  (resp.  $\alpha c$ ), with  $g \leq f$ . If a binary relation  $\rho$  is defined by  $A \rho B$  in case  $C \text{Cl}(A) \subseteq \alpha \text{Int}(B)$  (resp.  $\alpha \text{Cl}(A) \subseteq C \text{Int}(B)$ ), then by hypothesis  $\rho$  is a strong binary relation in the power set of  $X$ . If  $t_1$  and  $t_2$  are any elements of  $\mathbb{Q}$  with  $t_1 < t_2$ , then

$$A(f, t_1) \subseteq \{x \in X : f(x) \leq t_1\} \subseteq \{x \in X : g(x) < t_2\} \subseteq A(g, t_2);$$

since  $\{x \in X : f(x) \leq t_1\}$  is a  $C$ -closed (resp.  $\alpha$ -closed) set and since  $\{x \in X : g(x) < t_2\}$  is an  $\alpha$ -open (resp.

$C$ -open) set, it follows that  $CCl(A(f, t_1)) \subseteq \alpha Int(A(g, t_2))$  (resp.  $\alpha Cl(A(f, t_1)) \subseteq CInt(A(g, t_2))$ ). Hence  $t_1 < t_2$  implies that  $A(f, t_1) \rho A(g, t_2)$ . The proof follows from Theorem 2.1. ■

## Acknowledgements

This research was partially supported by Centre of Excellence for Mathematics(University of Isfahan).

## References

- [1] Hatir, E., Noiri, T. and Yksel, S. (1996) A Decomposition of Continuity. *Acta Mathematica Hungarica*, **70**, 145-150. <http://dx.doi.org/10.1007/BF00113919>
- [2] Dontchev, J. (1998) Between  $\alpha$ - and  $\beta$ -Sets. *Mathematica Balkanica*, **12**, 295-302.
- [3] Przemski, M. (1993) A Decomposition of Continuity and  $\alpha$ -Continuity. *Acta Mathematica Hungarica*, **61**, 93-98. <http://dx.doi.org/10.1007/BF01872101>
- [4] Dontchev, J. (1995) The Characterization of Some Peculiar Topological Space via  $\alpha$ - and  $\beta$ -Sets. *Acta Mathematica Hungarica*, **69**, 67-71. <http://dx.doi.org/10.1007/BF01874608>
- [5] Ganster, M. and Reilly, I. (1990) A Decomposition of Continuity. *Acta Mathematica Hungarica*, **56**, 299-301. <http://dx.doi.org/10.1007/BF01903846>
- [6] Katětov, M. (1951) On Real-Valued Functions in Topological Spaces. *Fundamenta Mathematicae*, **38**, 85-91.
- [7] Katětov, M. (1953) Correction to, "On Real-Valued Functions in Topological Spaces". *Fundamenta Mathematicae*, **40**, 203-205.
- [8] Brooks, F. (1971) Indefinite Cut Sets for Real Functions. *The American Mathematical Monthly*, **78**, 1007-1010. <http://dx.doi.org/10.2307/2317815>
- [9] Lane, E. (1976) Insertion of a Continuous Function. *Pacific Journal of Mathematics*, **66**, 181-190. <http://dx.doi.org/10.2140/pjm.1976.66.181>