

Geographic Differences in Allometric Patterns of Males of the Japanese Firefly *Luciola parvula*

Yutaka Iguchi 

Laboratory of Biology, Okaya, Japan

Email: bio-igu@f8.dion.ne.jp

How to cite this paper: Iguchi, Y. (2024) Geographic Differences in Allometric Patterns of Males of the Japanese Firefly *Luciola parvula*. *Advances in Entomology*, 12, 18-23.

<https://doi.org/10.4236/ae.2024.121002>

Received: October 24, 2023

Accepted: January 2, 2024

Published: January 5, 2024

Copyright © 2024 by author(s) and

Scientific Research Publishing Inc.

This work is licensed under the Creative

Commons Attribution International

License (CC BY 4.0).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

Abstract

It has been well known for many years that males of *Luciola parvula* (Coleoptera, Lampyridae) have two morphs, namely small and large morphs in Japan. This article performed a statistical reanalysis of their morphological measurements based on previously published data and compared them. As a result, the two morphs were found to have different allometric patterns in the Hakone and Nagano areas, Japan. In Hakone, the relationships between their pronotum width and length were expressed by a common allometric line and a shift in location along the line. However, in Nagano, the relationships between their pronotum width and body length were expressed by two parallel allometric lines (a common slope and different intercepts). The allometric pattern of males of this species may vary among different parts of the body or among different areas. Therefore, this article recommends future studies to measure the same parts of the body as previous studies to clarify geographic variation in allometric relationships.

Keywords

Body Size, Male Dimorphism, Allometric Pattern, Hakone, Nagano

1. Introduction

Luciola parvula (Coleoptera, Lampyridae) is a terrestrial firefly species widely distributed in Japan and one of the most popular insects, which is called Himebotaru, or Princess-firefly [1]. Therefore, many studies have been performed on various aspects such as life history [2], morphology [3] [4] [5], behavior [6] [7] [8], molecular phylogeny [9] [10] [11] [12], and biochemistry [13].

One of the interesting topics on this species is the presence of two male morphs, namely large and small male morphs [4] [5]. Male dimorphism in fireflies has been scarcely reported except for *Abseondita cerata* (previously named

Luciola cerata) in Taiwan region [14]. Recently the present author reanalyzed previously published morphological data on this species and found the two male morphs to show a discontinuous allometric relationship [15]. However, the reanalysis data sources were restricted to specimens collected in Nagano Prefecture.

As mentioned above, this species is widely distributed in Japan and has been studied on various aspects for more than forty years. Some reports say that the small male morph is relatively abundant below 800 m altitude in western Japan [4] [7], whereas others say that both morphs are found above 1000 m altitude [16] [17] [18] [19]. It is not yet clear on geographical variation in the two male morphs. One of the reasons is the lack of sufficient statistical analysis of morphological measurements of this species [15]. Therefore, it is important to reanalyze previously published data on specimens collected in various areas and examine the presence of the two morphs before proceeding to further studies on this species.

In order to contribute to the topic of the two male morphs on this species, the present author decided to reanalyze the data in a pioneer study on this topic by the late Dr Nobuyoshi Ohba [3]. He collected male adults around Mt. Hakone, central Japan and then classified them into the large morph above 700 m altitude and the small morph below this altitude. He showed a scatter plot of pronotum width and length, but did not conduct any further statistical analysis.

As far as the present author knows, there have been no detailed statistical studies on morphological data collected around Mt. Hakone so far. Nevertheless, based on Ohba's classification [3], genetic differences were evaluated between the two male morphs [9]. Therefore, the detailed reanalysis of Ohba's data [3] will help researchers plan future studies not only on morphological characteristics but also on molecular genetic characteristics.

2. Materials and Methods

The data analyzed here were obtained from Figure 2.3 of the literature [3] (p. 24) shown as a scatter plot of pronotum width and length in *L. parvula* around Mt. Hakone, Kanagawa Prefecture, Japan. The data consisted of 29 specimens of males of this species collected from 100 to 1200 m altitude, but the raw measurement data were not shown there. Therefore, in the present article, those scatter plot data were first digitized using the GSYS 2.4 software developed by Nuclear Reaction Data Centre, Hokkaido University, Japan [20].

As mentioned above, the 29 specimens were classified into the large morph above 700 m altitude and the small morph below this altitude [3]. Therefore, in the present article as well as in the author's previous article [15], allometric equations were applied to the two morphs using standardized major axis regression with the smatr package in the R software [21]. The data were \log_{10} -transformed into the log-log relationship between pronotum width (x) and pronotum length (y). Consequently, the \log_{10} -transformed allometric equation was expressed as:

$$\log_{10} y = \log_{10} a + b \log_{10} x \quad (1)$$

where a and b are constants. The constant b is the slope of the line, also known as the allometric coefficient.

3. Results

As shown in **Figure 1**, the two allometric lines representing the two morphs did not differ significantly in either slope ($\chi^2 = 0.878$, $df = 1$, $p = 0.349$) or elevation (y -intercept) ($\chi^2 = 0.00135$, $df = 1$, $p = 0.971$), but showed a significant shift along the common slope ($\chi^2 = 76.94$, $df = 1$, $p < 0.001$). The common slope ($b = 1.251$) was not significantly different from 1 ($\chi^2 = 2.342$, $df = 2$, $p = 0.310$), which indicated an isometric relationship between pronotum width and length.

4. Discussion

The allometric analysis of the literature [3] showed that the two male morphs of *L. parvula* around Mt. Hakone shared a common allometric line and shifted in location along the line. This allometric pattern was different from that in Nagano Prefecture found by the author's previous study [15], where the data of the literatures [16] [17] [18] [19] were reanalyzed (**Figure 2**).




Figure 1. Allometric relationship between pronotum width and length in the two male morphs of *L. parvula* fireflies around Mt. Hakone. The data were obtained from Figure 2.3 of the literature [3] (p. 24), where the 29 specimens were classified into the large morph above 700 m altitude and the small morph below this altitude. The standardized major axis regression lines were separately fitted to each group.




Figure 2. Different allometric relationships in the two male morphs of *L. parvula* fireflies in the Nagano and Hakone areas. The relationship in Nagano was analyzed in [15] and that in Hakone was analyzed in this article. The parameter b represents the allometric coefficient.

This difference may be because [3] and [16] [17] [18] [19] measured different parts of the body, namely pronotum width and pronotum length [3], and pronotum width and body length [16] [17] [18] [19]. Allometric relationships can vary geographically because of larval nutrition [22] [23] [24] [25], but they can also vary between different body parts for the same species [22]. Therefore, future studies on the male dimorphism of *L. parvula* will be recommended to compare geographical variations in allometric relationships between the same parts of the body.

Regarding genetic differences in the two male morphs of this species, [9] used allozyme loci and reported that they have genetic distance at the subspecies level. On the other hand, [11] used COI gene sequences and considered that this firefly may be pseudocryptic species with genetic divergence but not with appreciable morphological differences, except for body size. However, [15] revealed that the two male morphs of this species show different allometric relationships as well as different body sizes. Neither [9] nor [11] considered allometric relationships in this species. However, the results shown in the present article suggest that allometric analysis is very important in genetic studies to clarify the dimorphism of this species.

As mentioned above, there are two factors expected to cause the two different allometric patterns in the Nagano and Hakone areas. One is larval nutrition, and the other is gene expression. However, there is no evidence that these factors cause the two different allometric patterns in this species. Therefore, further ecological and genetic investigations are required to find the cause of the two different allometric patterns in this species.

5. Conclusion

Male size dimorphism is rarely observed in firefly species. However, the present study revealed that the large and small male morphs in *L. parvula* exhibit two different allometric patterns in two different areas, Nagano and Hakone, Japan. Further ecological and genetic studies will clarify the cause of the two different allometric patterns.

Acknowledgements

The author declares that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest. The author confirms sole responsibility for the following: study conception and design, data collection, analysis and interpretation of results, and manuscript preparation.

Conflicts of Interest

The author declares no conflicts of interest regarding the publication of this paper.

References

- [1] Takada, K. (2011) Popularity of Different Lampyrid Species in Japanese Culture as Measured by Google Search Volume. *Insects*, **2**, 336-342.
<https://doi.org/10.3390/insects2030336>
- [2] Ohba, N. (1975) Life History of *Hotaria parvula*. *Annual Report of the Yokosuka City Museum*, **21**, 5-8.
- [3] Ohba, N. (1986) Communication of Fireflies. Tokai University Press, Tokyo.
- [4] Ohba, N. (2016) Two Ecological Types of *Luciola parvula*. *Proceedings of the Japan Association for Fireflies Research (Zenkoku Hotaru Kenkyukai-shi)*, **49**, 25-27.
- [5] Mitsuishi, T. (2010) Distribution and Two Kinds (Large Type and Small Type) of *Luciola parvula* Living in Nagano Prefecture. *The Nature and Insects*, **598**, 11-14.
- [6] Ohba, N., Goto, Y., Aiuchi, M. and Watanabe, M. (1987) Flight of Male Firefly, *Hotaria parvula* Searching for Potential Mates. *Science Report of the Yokosuka City Museum*, **35**, 15-22.
- [7] Ohba, N. (2000) Flash Patterns and Communication of the Two Ecotypes in *Hotaria parvula*. *Science Report of the Yokosuka City Museum*, **47**, 1-22.
- [8] Takatsu, H., Minami, M., Tainaka, K.I. and Yoshimura, J. (2012) Spontaneous Flash Communication of Females in an Asian Firefly. *Journal of Ethology*, **30**, 355-360.
<https://doi.org/10.1007/s10164-012-0332-2>
- [9] Suzuki, H., Sato, Y., Fujiyama, S. and Ohba, N. (1993) Genetic Differentiation between Ecological Two Types of the Japanese Firefly, *Hotaria parvula*: An Electrophoretic Analysis of Allozymes. *Zoological Science*, **10**, 697-703.
- [10] Hiyori, Y., Mori, A., Baba, H. and Kusaoke, H. (2006) Morphology and Geographical Distribution of *Luciola parvula* Inferred from Genetic Analysis. *Proceedings of the Japan Association for Fireflies Research (Zenkoku Hotaru Kenkyukai-shi)*, **39**, 56-59.
- [11] Han, T., Kim, S.H., Yoon, H.J., Park, I.G. and Park, H. (2020) Evolutionary History of Species of the Firefly Subgenus *Hotaria* (Coleoptera, Lampyridae, Luciolinae, *Luciola*) Inferred from DNA Barcoding Data. *Contributions to Zoology*, **89**, 127-145.
<https://doi.org/10.1163/18759866-20191420>
- [12] Kusaoke, H., Onagawa, H., Miyahara, M., Yonezawa, M. and Mitamura, Y. (2022) Distribution of and Genetic Characteristics of the Small and Large Adults of *Luciola parvula*. *Proceedings of the Japan Association for Fireflies Research (Zenkoku Hotaru Kenkyukai-shi)*, **55**, 23-29.
- [13] Bessho-Uehara, M. and Oba, Y. (2017) Identification and Characterization of the Luc2-Type Luciferase in the Japanese Firefly, *Luciola parvula*, Involved in a Dim Luminescence in Immobile Stages. *Luminescence*, **32**, 924-931.
<https://doi.org/10.1002/bio.3273>
- [14] Wu, C.H., Jeng, M.L., South, A., Ho, J.Z. and Yang, P.S. (2010) Evidence for Two Male Morphs of *Luciola cerata* Olivier (Coleoptera: Lampyridae) Exhibiting Distinct Mating Behavior, with Implications for Sexual Selection. *The Coleopterists Bulletin*, **64**, 235-242. <https://doi.org/10.1649/0010-065X-64.3.235.9>
- [15] Iguchi, Y. (2023) Allometric Approach to the Two Male Morphs in the Japanese Firefly *Luciola parvula*. *Frontiers in Insect Science*, **3**, Article ID: 1230363.
<https://doi.org/10.3389/finsc.2023.1230363>
- [16] Mitsuishi, T. (2003) Distribution of the Small and Large Adults of *Luciola parvula* in Nagano Prefecture. *Proceedings of the Japan Association for Fireflies Research (Zenkoku Hotaru Kenkyukai-shi)*, **36**, 32-36.

- [17] Mitsuishi, T. (2004) Unique Distribution of the Small and Large Adults of *Luciola parvula* in Nagano Prefecture: I. *Proceedings of the Japan Association for Fireflies Research (Zenkoku Hotaru Kenkyukai-shi)*, **37**, 28-33.
- [18] Mitsuishi, T. (2005) Unique Distribution of the Small and Large Adults of *Luciola parvula* in Nagano Prefecture: II. *Proceedings of the Japan Association for Fireflies Research (Zenkoku Hotaru Kenkyukai-shi)*, **38**, 44-50.
- [19] Mitsuishi, T. (2007) New Distribution Records and Notes on the Unique Habitat of *Luciola parvula*: III. *Proceedings of the Japan Association for Fireflies Research (Zenkoku Hotaru Kenkyukai-shi)*, **40**, 35-39.
- [20] Suzuki, R. (2012) GSYS2.4 Manual. Second Edition- α . Hokkaido University Hospital, Sapporo. <http://www.jcprg.org/gsys/2.4/gsys24-e.pdf>
- [21] R Core Team (2023) R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna. <https://www.r-project.org/>
- [22] Shingleton, A.W., Frankino, W.A., Flatt, T., Nijhout, H.F. and Emlen, D.J. (2007) Size and Shape: The Developmental Regulation of Static Allometry in Insects. *Bio-Essays*, **29**, 536-548. <https://doi.org/10.1002/bies.20584>
- [23] Stern, D.L. and Emlen, D.J. (1999) The Developmental Basis for Allometry in Insects. *Development*, **126**, 1091-1101. <https://doi.org/10.1242/dev.126.6.1091>
- [24] Hooper, A. (2018) The Evolution of Condition-Dependent Life History: From Morphology to Lifespan. Doctoral Dissertation, The University of New South Wales, Sydney, 1-184.
- [25] Thompson, D.B. (2019) Diet-Induced Plasticity of Linear Static Allometry Is Not So Simple for Grasshoppers: Genotype-Environment Interaction in Ontogeny Is Masked by Convergent Growth. *Integrative and Comparative Biology*, **59**, 1382-1398. <https://doi.org/10.1093/icb/icz137>