

# Using Lorentz Violation for Early Universe GW Generation Due to Black Hole Destruction in the Early Universe as by Freeze

Andrew Beckwith

Physics Department, Chongqing University, Chongqing, China

Email: [rwill9955b@gmail.com](mailto:rwill9955b@gmail.com)

**How to cite this paper:** Beckwith, A. (2021) Using Lorentz Violation for Early Universe GW Generation Due to Black Hole Destruction in the Early Universe as by Freeze. *Journal of High Energy Physics, Gravitation and Cosmology*, 7, 993-1004.  
<https://doi.org/10.4236/jhepgc.2021.73058>

**Received:** April 30, 2021

**Accepted:** July 3, 2021

**Published:** July 6, 2021

Copyright © 2021 by author(s) and Scientific Research Publishing Inc. This work is licensed under the Creative Commons Attribution International License (CC BY 4.0).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

---

## Abstract

We are using information from a paper deriving a Lorentz-violating energy-momentum relation entailing an exact momentum cut of as stated by G. Salesi. Salesi in his work allegedly defines Pre Planckian physics, whereas we restrict our given application to GW generation and DE formation in the first  $10^{-39}$  s to  $10^{-33}$  s or so seconds in the early universe. This procedure is enacted due to an earlier work whereas referees exhibited puzzlement as to the physical mechanism for release of Gravitons in the very early universe. The calculation is meant to be complementary to work done in the Book “Dark Energy” by M. Li, X-D. Li, and Y. Wang, and also a calculation for Black hole destruction as outlined by Karen Freeze, *et al.* The GW generation will be when there is sufficient early universe density so as to break apart Relic Black holes but we claim that this destruction is directly linked to a Lorentz violating energy-momentum G. Salesi derived, which we adopt, with a mass  $m$  added in the G. Salesi energy momentum results proportional to a tiny graviton mass, times the number of gravitons in the first  $10^{-43}$  seconds.

## Keywords

Minimum Scale Factor, Cosmological Constant, DE

---

## 1. Introduction

What we are doing here is to utilize having the results of Salesi [1] as to a given actual Lorentz-violating energy-momentum relationship which we utilize to elucidate graviton contributions to an early universe derivation of DE, and the cosmological constant. The idea for this is based upon a referee and academic editor who felt puzzled as to the work done earlier [2] which postulated the existence of a breakup of primordial black holes as by itself contributing to DE,

and this publication is intended to fill in the actual conceptual gaps which lead to [2] having such a rocky reception. [2] was initiated specifically because work done in [3] as to a multiverse, was not well received, for reasons the author was told as in the eight equation of [3] one reviewer made the utterly outrageous statement that the author was modeling the Universe as a harmonic oscillator which is a canard since that equation was comparing the first integral of the Einstein-Hilbert action to an action for an initial spherical well, as by John Klauder as cited by the author in [4] so called enhanced quantization techniques [5].

In a nutshell, the author is trying to defuse mythologies which are in the way of a treatment of DE, which has been put in place by reviewers and a particular academic editor and the latest would allow for understanding the author's use of Miao Li, *et al.* [6].

In particular, the author hopes that judicious use of [1] will allow as to the use of [7] which several academic editors and reviewers took out of context and trivialized.

This will lead us to utilize the statement given by [8] linking the cosmological constant and massive gravity as given by

$$m_g = \frac{\hbar\sqrt{\Lambda}}{c} \quad (1)$$

This release of conditions for massive gravity should be in line with what is in Freeze [7],

$$\rho_{\text{BH breakup density}} = \frac{M_P^4}{32\pi} \cdot \left( \frac{M_P^4}{m^4} \right) \cdot \frac{1}{|1 + 3\omega_\phi|} \quad (2)$$

If the conditions of an early universe, are greater than this value, for Equation (2) [6] then according to Freeze, *et al.* primordial black holes would break apart. We state that this break up of primordial black holes would be enough to create an initial "sea" of gravitons, due to Equation (1) which would then add up to be in effect a value for a sufficient number of early universe gravitons, which would be added up per unit volume, to in fact sum up to an energy density equivalent to Equation (1) so we have massive gravitons and DE. Hence we will be adding up the number of gravitons which may be released due to Equation (2) and [9] which states the number of gravitons which may be emitted due to a black hole as given in it's page 47 is 0.1 percent of emitted energy from a nonrotating black hole. Keep in mind that this is for black holes, as given in [9] with mass:

$$M_{\text{primordial black hole}} \approx 10^{15} \cdot \left( \frac{t}{10^{23} \text{ s}} \right) \text{ grams} \quad (3)$$

For a  $10^{-5}$  gram black hole,  $t$  would have to be about  $10^{-43}$  seconds, and according to inflation expands space by a factor of  $10^{26}$  over a time of the order of  $10^{-43}$  to  $10^{-32}$  seconds. Meaning we had  $10^{-5}$  gram black holes at the start of inflation, and at the time the density of space would be greater than Equation (2) we would have a breakup of black holes if we had space-time density greater than or

equal to about Equation (2) then we have .1% of the mass of the broken BH contributing to gravitons, which after we review it may be relevant to Equation (1) above. One Planck mass is about  $10^{-5}$  grams. And it is worth noting in our development when we go past inflation, that we have Black holes growing to the value of about 1 grams, after  $10^{-40}$  seconds which is for a radii of approximately 1 centimeter, whereas we can and will define Black holes of  $10^{-5}$  grams which would be for less than a centimeter radii just after the start of inflation.

Unfortunately due to an unnamed editor, this picture as attested by Karen Freeze was deemed insufficiently rigorously motivated, and the fact it was put up and deemed allegedly not rigorous, means that we will be forced to have the backup done by [1] as stated. It is important to keep in mind that this was put in to give substance to the modeling given by [6] which is in both [2] and [3],

$$\frac{1}{2} \cdot \sum_i \omega_i \equiv V(\text{volume}) \cdot \int_0^{\tilde{\lambda}} \sqrt{k^2 + m^2} \frac{k^2 dk}{4\pi^2} \approx \frac{\tilde{\lambda}^4}{16\pi^2} \quad (4)$$

$$\xrightarrow{\tilde{\lambda}=M_{\text{Planck}}} \rho_{\text{boson}} \approx 2 \times 10^{71} \text{ GeV}^4 \approx 10^{119} \cdot \left( \rho_{DE} = \frac{\Lambda}{8\pi G} \right)$$

In stating this we have to consider that  $\rho_{DE} = \frac{\Lambda}{8\pi G} \approx \hbar \cdot \frac{(2\pi)^4}{\lambda_{DE}^4}$ , so then that the equation we have to consider is a wavelength  $\lambda_{DE} \approx 10^{30} \ell_{\text{Planck}}$  which is about  $10^{30}$  times a Planck radius of a space-time bubble which we discuss in [2] as a start point for a nonsingular expansion point for Cosmology, at the start of inflation with the space-time bubble of about a Planck length radius we have to consider is a wavelength  $\lambda_{DE} \approx 10^{30} \ell_{\text{Planck}}$  which is about  $10^{30}$  times a Planck length radius of a space-time bubble [3] as a nonsingular expansion point for Cosmology, at start of inflation with the space-time bubble of about a Planck length radius in size. Having said that

$$\lambda_{DE} \approx 10^{30} \ell_{\text{Planck}} \quad (4a)$$

And then we can write up having

$$\rho_{DE} = \frac{\Lambda}{8\pi G} \approx \hbar \cdot \frac{(2\pi)^4}{\lambda_{DE}^4} \quad (4b)$$

This is going to create difficulties which is going to lend us to utilize [1] directly and more so we have a way to refine the argument given in Equation (4), Equation (4a) and Equation (4b). The UNKNOWN academic editor is making this next step compulsory.

## 2. Making Use of [1] and the Use of Lorentz Violation, as in [1] to Define More Precisely the Contribution of Gravitons to both DE, Due to the Breakup of Black Holes

To do this we need to review the Lorentz violating energy-momentum relationship. In short we have that

$$E^2 = p^2 + m^2 - \tilde{\lambda} p^3 \quad (5)$$

where the positive LV parameter  $\tilde{\lambda}$  is usually assumed of the order of Planck mass,  $\lambda \sim 1/M$  (planck mass). This Lorentz violating energy-momentum relationship leads to, according to [1],

$$d\rho = \frac{8\pi p^3 c}{h} \cdot \frac{\sqrt{1 + \frac{m^2}{p^2} - \tilde{\lambda} p}}{\exp\left(\left[\frac{cp}{k_B T_{temp}}\right] \cdot \sqrt{1 + \frac{m^2}{p^2} - \tilde{\lambda} p}\right) - 1} \cdot dp \quad (6)$$

$p \approx 1/\tilde{\lambda}$  is used if we integrate, Equation (6) and if we use the first order Romberg numerical integration scheme as given in [10], page 695, so then for high temperature,

$$\begin{aligned} \rho &\approx \frac{8\pi M_p^3 c}{h} \cdot \frac{\sqrt{\frac{m^2}{M_p^2}}}{\exp\left(\left[\frac{cp}{k_B T_{temp}}\right] \cdot \sqrt{\frac{m^2}{M_p^2}}\right) - 1} \\ &\approx \frac{4M_p^2 c}{\hbar} \cdot \left( \left[\frac{cm}{k_B T_{temp}}\right] - \frac{1}{2} \left(\frac{cm}{k_B T_{temp}}\right)^2 \right) \\ &\xrightarrow{k_B=c=\hbar=M_p=1} 4 \cdot \left( \left[\frac{m}{T_{temp}}\right] - \frac{1}{2} \left(\frac{m}{T_{temp}}\right)^2 \right) \end{aligned} \quad (7)$$

We will then in the next section interpret Equation (7) when we set

$$m \approx m_g \cdot N_g \quad (7a)$$

### 3. Interpreting Equation (7) When Equation (7a) Is Used, So as to Ascertain the Number of Gravitons

We are then looking at [1],

$$\rho \approx \frac{4M_p^2 c}{\hbar} \cdot \left( \left[\frac{cm_g N_g}{k_B T_{temp}}\right] - \frac{1}{2} \left(\frac{cm_g N_g}{k_B T_{temp}}\right)^2 \right) \quad (7b)$$

Using the Planck units renormalized such that  $k_B = c = \hbar = M_p = 1$ , we have that we are looking at resetting Equation (7b) so that the above will be roughly,

$$\rho \approx 4 \cdot \left( \left[\frac{10^{-65} N_g}{(T_{temp}/T_P)}\right] - \frac{1}{2} \left(\frac{10^{-65} N_g}{(T_{temp}/T_P)}\right)^2 \right) \approx 10^{-60} \quad (7c)$$

We then can up to a modeling round off make the following approximation,

$$\left( \frac{10^{-57} N_g}{(T_{temp}/T_P)} \right)^2 - 2 \cdot \left[ \frac{10^{-57} N_g}{(T_{temp}/T_P)} \right] + 4 \times 10^{-60} \approx 0 \quad (7d)$$

‘This value of Equation (7d) will lead to approximately if  $(T_{temp}/T_P) \sim 1$ ,

$$N_g \approx 10^{57} \times (1 \pm (1 - 2 \times 10^{-60})) \approx 2 \times 10^{57} - 2 \times 10^{-3} \quad (8)$$

What is Equation (8) saying? We ascertain Equation (8) especially if Equation (7c) is set to reflect upon the number of Gravitons which may give us Dark Energy. In evaluating Equation (8) we have that  $T_{temp} \leq T_p$  for reasons we will go into in the next section.

#### 4. Interpreting Equation (7) and Equation (8) in Terms of Dark Energy, If Gravitons Produce DE and How This Ties in with the Freeze Suggestion as to the Breakup of Black Holes for Gravitons in First $10^{-27}$ Seconds

Roughly put, it means that there is about 1000 to 10,000 mini black holes, in between  $10^{-43}$  seconds to  $10^{-32}$  seconds which would be destroyed so as to release about  $10^{57}$  gravitons, equivalent to about 1 Planck mass. In terms of space-time this would be commensurate to a density of  $7 \times 10^{-30}$  g/cm<sup>3</sup>, or having a radii of 1000 kilometers for a volume of space for about  $7 \times 10^{-6}$  g/(1000 km)<sup>3</sup> of density for approximately  $10^{57}$  gravitons to have one Planck mass of gravitons released in a volume of space for a radii of about 1000 kilometers after  $10^{-32}$  seconds.

This would be about 1000 to 10,000 destroyed mini black holes in less than 1.057 times  $10^{-10}$  of a light year in radial distance for 1 Planck mass of radiated gravitons, in far less than  $10^{-1}$  seconds in cosmological expansion.

The Universe was once just the radius of the Earth-to-the-Sun, which happened when the Universe was about a trillionth ( $10^{-12}$ ) of a second old, *i.e.* a sphere of 149.6 million km and the region for about  $10^{-43}$  -  $10^{-40}$  seconds was about  $10^3$  kilometers in radial size.

It is important to keep these figures in line and visualize how a sphere of about  $10^3$  kilometers in radial size would be able to have  $10^{57}$  gravitons released in order to have DE formed and possibly then the Cosmological constant. Keep in mind the assumed radii of the universe today is  $4.4 \times 10^{23}$  km.

#### 5. What $10^{57}$ Gravitons in a Radius of 1000 Kilometers Means in Terms of DE and a Cosmological Constant Calculation

We will first of all refer to an early universe treatment of the uncertainty principle is, in the startup of inflationary cosmology [11].

Set

$$a(t) = a_{\text{initial}} t^\gamma \quad (9)$$

$$\rho \approx \frac{\dot{\phi}^2}{2} + V(\phi) \equiv \frac{\gamma}{8\pi G} \cdot t^2 + V_0 \cdot \left\{ \sqrt{\frac{8\pi G V_0}{\gamma(3\gamma-1)}} \cdot t \right\}^{\sqrt{\frac{\gamma}{4\pi G}} - \sqrt{\frac{8\pi G}{\gamma}}} \quad (10)$$

$$V = V_0 \cdot \left\{ \sqrt{\frac{8\pi G V_0}{\gamma(3\gamma-1)}} \cdot t \right\}^{\sqrt{\frac{\gamma}{4\pi G}} - \sqrt{\frac{8\pi G}{\gamma}}} \quad (11)$$

Also we will set

$$\rho \approx \frac{\dot{\phi}^2}{2} + V(\phi) \equiv \frac{\gamma}{8\pi G} \cdot t^2 + V_0 \cdot \left\{ \sqrt{\frac{8\pi G V_0}{\gamma(3\gamma-1)}} \cdot t \right\}^{\sqrt{\frac{\gamma}{4\pi G}} \cdot \sqrt{\frac{8\pi G}{\gamma}}} \quad (12)$$

The value of time  $t$  will be set as  $t \sim (10^{-32} \text{ s}/t(\text{Planck}))$  whereas we can utilize the ideas of having Planck time set  $\sim 5 \times 10^{-44}$  seconds, hence,  $t \sim 10^{12}$ , in Planck Units, whereas  $\hbar = G = \ell_p = m_p = k_B = 1$ , so then we will have, in this situation, Equation, (12) as reset as [11]:

$$\begin{aligned} \rho &\approx \frac{\gamma \cdot (10^{24})}{8\pi} + V_0 \left\{ \frac{8\pi \cdot V_0 \cdot (10^{12})}{\gamma \cdot (3\gamma-1)} \right\}^{\frac{1}{2} \left( \sqrt{\frac{\gamma}{4\pi}} \cdot \sqrt{\frac{\pi}{\gamma}} \right)} \\ &\approx \frac{E_{\text{effective}}}{(1000 \text{ km})^3} \approx \frac{E_{\text{effective}}}{(6.25 \times 10^{40} (\ell_p = 1))^3} \end{aligned} \quad (13)$$

The interesting thing, is that the factor of roughly  $10^{-120}$  shows up in this situation so as to imply that there may be some linkage between setting the effective energy as roughly some proportional power value of Planck Mass. Taking into consideration as to what was done in terms of an earlier document as to Planck mass, consider the afore mentioned DE density,

$$10^2 m_p (\text{gravitons}) \Rightarrow 10^{67} \text{ gravitons per } 10^5 m_p (\text{black hole}) \quad (14)$$

### 6. Making DE Equivalent to a Sea of Initial Gravitons, in Regime $10^{-43}$ to $10^{-32}$ Seconds

Roughly put, one hydrogen atom is about 1.66 times  $10^{-24}$  grams. The weight of a Massive graviton is about  $10^{-65}$  grams [8] [9], hence we are talking about  $10^{-22}$  grams, or about  $10^{44}$  gravitons, with each graviton about  $6 \times 10^{-32}$  eV/c<sup>2</sup> After  $10^{-27}$  seconds, the following in the set of equations given below are Equivalent, and that these together will lead to a cosmological Constant,  $\Lambda$  of the sort which we will be able to refer to later,

$$1 \text{ graviton} \approx 10^{-65} \text{ g} \quad (14a)$$

$$M_{PBH} (10^{-32}) \approx 10^{15} \times \left( \frac{10^{-32} \text{ s}}{10^{-23} \text{ s}} \right) \text{ g} \approx 10^6 \text{ g} \approx 10^{11} M_p \quad (14b)$$

$$M_{PBH} (10^{-38}) \approx 10^{15} \times \left( \frac{10^{-38} \text{ s}}{10^{-23} \text{ s}} \right) \text{ g} \approx 10^0 \text{ g} \approx 10^5 M_p \quad (14c)$$

If so then, using Equation (14c), a  $10^5$  Planck mass sized black hole if it has 1/1000 of its mass converted into gravitons, would have  $10^2$  time Planck mass =  $10^{63}$  gravitons, *i.e.* for  $10^{63}$  gravitons which would occur for  $10^{-38}$  seconds. To match the restrictions as given in Equation (7d) and Equation (8) we need to look at what would allow for  $10^{57}$  gravitons, instead of  $10^{63}$  gravitons. This would entail having

$$M_{PBH} (10^{-43}) \approx 10^{15} \times \left( \frac{10^{-43} \text{ s}}{10^{-23} \text{ s}} \right) \text{ g} \approx 10^{-5} \text{ g} \approx 1M_p \quad (14d)$$

Equation (14d) is in fidelity with having, if 1/1000 of its mass converted to gravitons, a situation where there would be  $10^{57}$  gravitons. At a time of  $10^{-43}$  seconds. We are then examining what happens at the end of inflation. The supposition I have is that one can use 1/1000 of the mass as given in Equation (14d) for Gravitons and to thereby have  $10^{-8}$  g for gravitons, per black hole of mass  $10^{-5}$  g. If one has say  $10^{57}$  gravitons, for a 1000 kilometer regime as say in the first  $10^{-32}$  Seconds we then have for  $10^{-65}$  g per graviton, we are then having for gravitons a value of to be diverted to  $10^{-8}$  g per black hole.

**Table 1**, inputs into forming DE if one is assuming uniform distribution of BH to be broken up by Karen Freeze model as given in the following **Table 1**.

Assuming that gravitons contribute to the Dark Energy value will lead to us using the Karen Freeze model, with gravitons being released in the early universe by the breakup of early universe black holes which have a maximum value of about 1 g, as opposed to the value of the Sun which has about  $10^{33}$  grams, in first  $10^{-32}$  seconds. The divergence from the standard model can be seen in postulating a non singular start to the universe and a simple way to do it, as follows. Assume at the beginning, one has a spherical shell defined by a volume in the regime of radial space defined by  $a - \epsilon \leq r \leq a + \epsilon$  [12],

$$V_{\epsilon} = \frac{8\pi\epsilon}{3} \cdot (3a^2 + \epsilon^2) \tag{15}$$

And then for when one has if one has a heat strength of A, for this radial “shell”  $S_{\epsilon}$ ,

$$\delta_{A,\epsilon}(r) = \begin{cases} A/V_{\epsilon} & \text{if } r \in S_{\epsilon} \\ 0 & \text{otherwise} \end{cases} \tag{16}$$

Then one has the following “integration” in the region of “space-time”,

$$\int_0^{2\pi} \int_0^{\pi} \int_{a-\epsilon}^{a+\epsilon} \delta_{A,\epsilon}(r) \tilde{r}^2 \sin \phi d\tilde{r} d\phi d\theta = A \tag{17}$$

Following the line of reasoning, we will be examining briefly how this bubble-shell start to cosmology could commence, and how to interpret both Equation (16) and Equation (17) and how to make the reset value of when the universe started, and its subsequent inflation in sync with all the formulas in this document: This value of A would be say the beginning set of values of say the point of astrophysics for the breakup of black holes after the end of the inflationary era.

**Table 1.** Summary of inflationary and post inflationary information.

Number of black holes,	Mass of black hole of size 1 Planck mass set aside for gravitons	Mass of black hole for $10^8$ gravitons	Radii of proto universe
$10^8$	$10^{-8}$ grams	$10^{-5}$ g = 1 Planck mass	1000 Kilometers
Volume of Universe is $10^3$ kilometers, cubed	Starting range for Mass of black hole for Gravitons	Assumed to be starting range of BH masses, at about $10^{-43}$ seconds	From less than a meter to 1000 Kilometers for constructing black holes which may be torn asunder by Karen Freeze’s criteria

## 7. What $10^{57}$ Gravitons in a Radius of 1000 Kilometers Means If We Go to the Rosen Early Universe Cosmology

Let us first recall the Shalyt-Margolin and Tregubovich (p.73) [13], Shalyt-Margolin (p.62) [14] [15],

$$\Delta t \geq \frac{\hbar}{\Delta E} + \gamma t_p^2 \frac{\Delta E}{\hbar} \Rightarrow (\Delta E)^2 - \frac{\hbar \Delta t}{\gamma t_p^2} (\Delta E) + \frac{\hbar^2}{\gamma t_p^2} = 0$$

$$\Rightarrow \Delta E = \frac{\hbar \Delta t}{2\gamma t_p^2} \cdot \left( 1 + \sqrt{1 - \frac{4\hbar^2}{\gamma t_p^2 \cdot \left(\frac{\hbar \Delta t}{2\gamma t_p^2}\right)^2}} \right) = \frac{\hbar \Delta t}{2\gamma t_p^2} \cdot \left( 1 \pm \sqrt{1 - \frac{16\hbar^2 \gamma t_p^2}{(\hbar \Delta t)^2}} \right) \quad (18)$$

For sufficiently small  $\gamma$ . The above could be represented by [3] [14],

$$\Delta E \approx \frac{\hbar \Delta t}{2\gamma t_p^2} \cdot \left( 1 \pm \left( 1 - \frac{8\hbar^2 \gamma t_p^2}{(\hbar \Delta t)^2} \right) \right) \quad (19)$$

$$\Rightarrow \Delta E \approx \text{either } \frac{\hbar \Delta t}{2\gamma t_p^2} \cdot \frac{8\hbar^2 \gamma t_p^2}{(\hbar \Delta t)^2}, \text{ or } \frac{\hbar \Delta t}{2\gamma t_p^2} \cdot \left( 2 - \frac{8\hbar^2 \gamma t_p^2}{(\hbar \Delta t)^2} \right)$$

This would lead to a minimal relationship between change in  $E$  and change in time as represented by Equation (19), so that we could to first order, say be looking at something very close to the traditional Heisenberg uncertainty principle results of approximately,

$$\Delta E \approx \frac{\hbar \Delta t}{2\gamma t_p^2} \cdot \frac{8\hbar^2 \gamma t_p^2}{(\hbar \Delta t)^2} \equiv \frac{4\hbar}{\Delta t} \quad (20)$$

Or,

$$\Delta E \Delta t \approx 4\hbar \quad (21)$$

Having brought this up, let us then go to the Rosen [16] version of cosmology, and this needs explanation due to its rescaling of the values of the cosmology time and temperatures involved. The key point of this mini chapter will be to summarize derivation of the space-time temperature [16].

$$T = (\rho_F / \sigma)^{1/4} \cdot \frac{\tilde{a} r^7}{(\tilde{a}^4 + r^4)^2} \quad (22)$$

With  $\tilde{a} = 10^{-3} \text{ cm}$ ,  $(\rho_F / \sigma)^{1/4} = 1.574 \times 10^{32} \text{ K (kelvin)}$

Then according to [2], the initial temperature is

$$T_{\text{initial}} = 2.65 \times 10^{-180} \text{ K (kelvin)} \quad (23)$$

Whereas the temperature where one has the breakup of Primordial black holes starting is at

$$T_{\text{black hole breakup starts}} = 7.41 \times 10^{31} \text{ K (kelvin)} \quad (24)$$

Whereas we start the derivation of Equation (22) in reference [2], and the extreme value of the temperature  $T$  for breaking up black holes, again by [2] leads

to how black holes may contribute to the growth of DE if we have graviton production, so we consider when a body of mass  $m$  and radius  $R$  break apart. As given in [2] by an argument given by Freeze, we have then that if  $R(\text{radius})$  is between 1 meter to say 1000 Kilometers this mass  $m$  breaks up for  $m$  as given by

$$m \approx \frac{8\pi R(\text{radius})^3 \cdot \rho}{3} \quad (25)$$

Here, the density function is given by Equation (12) and Equation (13), for our application and also we obtain for black holes a break up criteria for mass  $m$  Black holes if,

$$m \approx -\left(\frac{4\pi\rho}{3}\right) \cdot \left(1 + 3 \cdot \frac{\rho_G}{p_G}\right) \cdot \frac{8m^3}{M_P^6} \quad (26)$$

So we can have the start of breakup of black holes, if we have gravitons from 1/1000 of the mass of given black holes, and if black holes contribute DE according to when pressure is approximately equal to the negative value of the density which would lead to a Black hole contribution of Equation (27) to DE. As given below.

$$\text{DE from black holes} = 7 \times 10^{-30} \text{ g/cm}^3 = 7 \times 10^{-6} \text{ g/(1000 km)} \quad (27)$$

This rough value of DE, as given in Equation (27) will be directly compared to what we can expect as far as applying in Equation (21) as to comment directly on the  $\Delta t$  time interval for the active generation of DE in the early cosmos.

Keep in mind that J. W. Moffat in [17] postulated in the initial phases of cosmology a situation for which we have no conservation of energy, and in fact this is exactly the situation we could be portraying here, that is if [2] and the description of the Rosen cosmology as in [16] are not wrong.

Quote, from [17]:

The spontaneous symmetry broken phase will induce a violation of conservation of energy and explain the generation of matter in the very early universe.

End of quote:

This would be doable if the initial phases of creation of the Universe follow [16] and if we utilize, initially a near zero temperature start regime in the early universe, as in [16]. The early universe will have an energy input via thermal inputs of a value commensurate with [2].

$$E(\text{thermal energy}) = \frac{d(\text{dim}) \cdot k_B \cdot T_{\text{universe}}}{2} \quad (27)$$

## 8. Conclusion: Using Equation (27) in Conjunction with Reference [16] [17] and Equation [5] for Symmetry Breaking and Its Possible Tie in with DE

According to [17],

Quote:

The spontaneous symmetry breaking mechanism in the vierbein gauge for-

malism has 3 massless degrees of freedom associated with the O(3) rotational invariance, and 3 massive degrees of freedom associated with the broken Lorentz “boosts”. The massive quantum gravity in 3 + 1 dimensions can satisfy unitarity and be renormalizable, in contrast to the  $D = 4$  quantum gravity which will violate unitarity if renormalizable [18].

End of quote:

In short, the idea in the presentation I am doing is that there would be a massless situation until after the turn about point of thermal heating of the Big Bang universe, and this would correspond to symmetry breaking precisely due to the Equation (27) formulation allowing for a massless regime of space-time before the end of inflation, whereas after the end of inflation, we can use the black hole destruction idea given in [7] directly so as to commence.

After we have this value satisfied, set from Equation (11) and Equation (12) setting up the formulation of conditions for the creation of DE according to which we are looking at in the regime of space-time from less than a meter in radii to 1000 km in radii we have the formation of DE with the given frequency, for setting up DE, and then by default the Cosmological constant.

$$V(\phi(t)) \approx \omega_{\text{Graviton generated DE}} \quad (28)$$

Using the ideas of [17] and [18], the Lorentz violation would ensue in this situation as largely a byproduct of the situation outlined in [17] whereas on page 4 of that article we have,

Quote:

At the Planck energy  $E(\text{Planck})$  the local Lorentz vacuum symmetry is spontaneously broken. We postulate the existence of a field,  $\phi$ , and assume that the vacuum expectation value (vev) of the field,  $\langle \phi \rangle$ , will vanish for  $E = E_c < E(\text{Planck})$  or at a temperature  $T < T_c \sim M(\text{Planck})$

End of quote:

We argue that if the build-up of temperature is commensurate with [16] occurs in an early universe and that if we have no DE before the end of inflation and that if DE commences in being created with the use of Equation (28) for the start of DE and the breakup of black holes, via the mechanism brought up by [16]. Mofatt appeals to a generalization of the Wheeler De Witt equation, via complex time whereas our approach takes as its starting point for Lorentz symmetry breaking an influx of “heat” via [16] as well as then applying the nonlinear dispersion relationship given in Equation (5) which is allowed to commence in full generality at the “turning point” of thermality which Rosen and Israelit conceived of.

Furthermore we should investigate if there is a linkage in any sense between Equation (28) and the ideas brought up by Novello [8] which we regard as significant and worthy of review in their own right. And we will be investigating if [19] brings up issues which are relevant to future inquires in this matter so presented. We also state that we are now assuming an interlude of quantum involvement with this problem if and when Equation (21), the deformed relativity

modifications of the HUP are involved, as we claim they will be involved, especially after the turn about regime where temperatures of the Universe cease to be monotonically rising, and the breakup of primordial black holes commences.

The author is aware of prior refereeing, as in opposition to [2] and states that a timely use of Equation (15), Equation (16) and Equation (17) may prove useful in ascertaining a linkage between a pre Planckian Universe state, and the Universe as of today. More of which may be introduced in a timely fashion by the author in future publications plus [20].

## Acknowledgements

Work is supported in part by National Nature Science Foundation of China grant No. 11375279.

## Conflicts of Interest

The author declares that there is no conceivable conflict of interest in this document.

## References

- [1] Salesi, G. (2012) Lorentz-Violating Dynamics in the Pre-Planckian Universe. *Physical Review D*, **85**, Article ID: 063502. <https://doi.org/10.1103/PhysRevD.85.063502>  
<https://arxiv.org/abs/1110.2234>
- [2] Beckwith, A. (2021) A Solution of the Cosmological Constant and DE Using Breakup of Primordial Black Holes, Via a Criteria Brought Up by Dr. Freeze Which Initiates DE as Linked to Inflation. *Preprints*, **2021**, Article ID: 2021040210. <https://doi.org/10.20944/preprints202104.0210.v1>
- [3] Beckwith, A. (2021) A Solution of the Cosmological Constant, Using Multiverse Version of Penrose CCC Cosmology, and Enhanced Quantization Compared. *Preprints*, **2021**, Article ID: 2021020208. <https://doi.org/10.20944/preprints202102.0208.v1>
- [4] Beckwith, A.W. (2021) Using “Enhanced Quantization” to Bound the Cosmological Constant, (for a Bound-On Graviton Mass), by Comparing Two Action Integrals (One Being from General Relativity) at the Start of Inflation. In: Sidharth, B.G., Murillo, J.C., Michelini, M. and Perea, C., Eds., *Fundamental Physics and Physics Education Research*, Springer, Cham, Switzerland, 21-36. [https://doi.org/10.1007/978-3-030-52923-9\\_3](https://doi.org/10.1007/978-3-030-52923-9_3)
- [5] Klauder, J.R. (2015) *Enhanced Quantization: Particles, Fields & Gravity*. World Scientific, Singapore. <https://doi.org/10.1142/9452>
- [6] Li, M., *et al.* (2015) *Dark Energy*. Peking University Press, World Scientific, Singapore.
- [7] Freeze, K., Brown, M. and Kinney, W. (2012) The Phantom Bounce, A New Proposal for an Oscillating Cosmology. In: Mercini-Houghton, L. and Vaas, R., Eds., *The Arrows of Time, A Debate in Cosmology*. Springer, Heidelberg, 149-156. [https://doi.org/10.1007/978-3-642-23259-6\\_7](https://doi.org/10.1007/978-3-642-23259-6_7)
- [8] Novello, M. (2005) The Mass of the Graviton and the Cosmological Constant Puzzle. <https://arxiv.org/abs/astro-ph/0504505>
- [9] Calmet, X., Carr, B. and Winstanley, E. (2014) *Quantum Black Holes*. Springer,

- Heidelberg. <https://doi.org/10.1007/978-3-642-38939-9>
- [10] Willinger, D.Z. (1996) CRC Standard Mathematical Tables, and Formulae. 30th Edition, CRC Press, Boca Raton, USA. <https://doi.org/10.1201/NOE0849324796>
  - [11] Thanu, P. (2006) An Invitation to Astrophysics. World Press Scientific, Singapore. <https://doi.org/10.1142/6010>
  - [12] Hermann, F. (1995) Principles of Analysis [Math 527 (Notes)]. University of Arizona Applied Math in House Publishment, Tucson, USA.
  - [13] Shalyt-Margolin, A.E. (2006) Deformed Density Matrix and 21 Quantum Entropy of the Black Hole. *Entropy*, **8**, 31-43. <https://doi.org/10.3390/e8010031>
  - [14] Shalyt-Margolin, A.E., (2005) The Density Matrix Deformation in Physics of the Early Universe and Some of Its Implications, Quantum Cosmology Research Trends. *Horizons in World Physics*, **246**, 49-91.
  - [15] Shalyt-Margolin, A.E. and Tregubovich, A.Y. (2004) Deformed Density Matrix and Generalized Uncertainty Relation in Thermodynamics. *Modern Physics Letters A*, **19**, 71-81. <https://doi.org/10.1142/S0217732304012812>
  - [16] Rosen, N. and Israelit, M. (1991) A Simple Model of the Universe without Singularities. In: Zichichi, A., de Sabbata, V. and Sánchez, N., Eds., *Gravitation and Modern Cosmology*. Springer, Boston. [https://doi.org/10.1007/978-1-4899-0620-5\\_14](https://doi.org/10.1007/978-1-4899-0620-5_14)
  - [17] Moffat, J.W. (2010) Lorentz Violation of Quantum Gravity. *Classical and Quantum Gravity*, **27**, Article ID: 135016. <https://doi.org/10.1088/0264-9381/27/13/135016>  
<https://arxiv.org/abs/0905.1668>
  - [18] Stelle, K.S. (1977) Renormalization of Higher-Derivative Quantum Gravity. *Physical Review D*, **16**, 953-969. <https://doi.org/10.1103/PhysRevD.16.953>
  - [19] Ng, Y.J. (2021) Holographic Foam Cosmology: From the Late to the Early Universe. *Symmetry*, **13**, 435 p. <https://doi.org/10.3390/sym13030435>
  - [20] Kiefer, C., Polarski, D. and Starobinsky, A.A. (2000) Entropy of Gravitons Produced in the Early Universe. *Physical Review D*, **62**, Article ID: 043518. <https://doi.org/10.1103/PhysRevD.62.043518>  
<https://arxiv.org/abs/gr-qc/9910065>