

# **Compact Non-Linear Gravitomagnetic Fields as Dark Matter**

## Stuart Marongwe<sup>1\*</sup>, Phil Bouchard<sup>2</sup>

<sup>1</sup>Physics Department, University of Botswana, Gaborone, Botswana <sup>2</sup>Fornux Inc., Montreal, Canada Email: \*stuartmarongwe@gmail.com, phil@fornux.com

How to cite this paper: Marongwe, S. and Bouchard, P. (2025) Compact Non-Linear Gravitomagnetic Fields as Dark Matter. Journal of High Energy Physics, Gravitation and Cosmology, 11, 1031-1038. https://doi.org/10.4236/jhepgc.2025.113066

Received: May 23, 2025 Accepted: July 22, 2025 Published: July 25, 2025

Copyright © 2025 by author(s) and Scientific Research Publishing Inc. This work is licensed under the Creative Commons Attribution International License (CC BY 4.0). http://creativecommons.org/licenses/by/4.0/ (cc)

۲ **Open Access** 

### Abstract

We introduce a model of dark matter inspired by non-linear gravito-electromagnetism formulation of Roy Maartens and Bruce Bassett. The model is a fully covariant 1 + 3 electromagnetic analogy of gravity characterized by Weyl gravito-electric and gravito-magnetic spatial tensor fields in which the Bianchi identities are the dynamical equations. The nonlinear rotationally invariant vacuum Bianchi equations describe the properties of space-time within a given locality. We identify the compact gravito-magneto field as a Ricci soliton in Einstein's field equations that describes dark matter. This component provides the extra gravity which is currently attributed to a variety of theorized forms of non-baryonic matter collectively known as dark matter.

## **Keywords**

Dark Matter, Non-Linear Gravito-Electromagnetism, Ricci Soliton, Baryonic Tully-Fisher Relation, Milgrom's Constant

# **1. Introduction**

Currently, one of the most intriguing questions in fundamental physics is, what is the nature and origins of dark matter (DM) and dark energy (DE)? In astrophysics and cosmology, DM is attributed to either some exotic non-baryonic matter or to a modification of the law of gravity at large scales. Particles beyond the standard model of particle physics are actively being searched either through direct detection experiments or through collider experiments. So far, these particles have proved elusive with some models falling out of favor as a series of direct detection experiments have ruled out a gamut of predicted energy scales and cross-sectional areas [1]-[6]. A broad and in-depth review of the variety of proposed ideas on particulate DM can be found in [7]-[11] and the references therein. Within the formalism of modified gravity F(R), F(G) and F(T) gravity theories are actively being explored with the hope that a specific model can be found that reduces to the phenomenological model referred to as MOND [12] [13]. Further reviews of such approaches can be found in [14]-[17]. There are many ways of testing Modified Gravity theories. The most promising methods involve the use of gravitational waves [18] [19]. For instance, the LIGO/VIRGO detection of a neutron star merger event GW170817/GRB170817 provided a significant observational result. This event demonstrated the arrival of gravitational waves (gravitons) and electromagnetic waves (photons) at the Earth detector within 1.7 seconds of each other, challenging modified gravity theories that violated Lorentz Invariance.

In the present work, we attempt to shed some light on the enigmas of DM from the perspective of gravito-electromagnetism (GEM), specifically through the Maartens-Bassett formalism of non-linear GEM [20]-[22]. GEM arises as the name suggests from a rich and detailed correspondence between General Relativity (GR) and electromagnetism [23]-[25]. This strong correspondence is reflected in the Weyl tensor which has a Maxwell like form, the Bel-Robinson tensor with similarities to the electromagnetic energy-momentum tensor and the Bianchi identities' similarities to the dynamical equations of electromagnetism.

The paper is structured as follows: first, in the preliminaries section, we recall the fundamental equations of the Maartens-Basset formalism of non-linear GEM. We then implement this formalism in the linearized form to obtain the modified field equations of GR that include effects attributed to DM.

## 2. Preliminaries

(

In Ref. [23], Ellis provides Maxwell's equations in the 1 + 3 streamlined form as follows:

$$D^a E_a = -2\omega^a H_a + \varrho, \tag{1}$$

$$D^a H_a = 2\omega^a E_a, \tag{2}$$

$$\dot{E}_a - \nabla \times H_a = \frac{2}{3}\theta E_a + \sigma_{ab}E^b - [\omega, E]_a + [\dot{u}, H]_a - j_a,$$
(3)

$$\dot{H}_{a} + \nabla \times E_{a} = \frac{2}{3} \theta H_{a} + \sigma_{ab} H^{b} - [\omega, H]_{a} - [\dot{u}, E]_{a}, \qquad (4)$$

where  $\varrho = -J_a u^a$  is the electric charge density,  $u^a$  the four velocity,  $\theta = D^a u_a$ , the shear  $\sigma_{ab} = D_{\langle a u_b \rangle}$ ,  $\omega_a = -\frac{1}{2} \nabla \times u_a$  is the vorticity and  $\dot{u}_a = \dot{u}_{\langle a \rangle}$  is the four acceleration.

The above operators obey the following covariant identities:

$$D_a \dot{f} = D_a \dot{f} - \frac{1}{3} \theta D_a f + \dot{u}_a \dot{f} - \sigma_a^{\ b} D_b f - \left[\omega, Df\right]_a + u_a \dot{u}^b D_b f,$$
(5)

$$\nabla \times D_a f = 2\dot{f}\omega_a,\tag{6}$$

$$D^{a}\left[V,W\right]_{a} = W^{a}\nabla \times V_{a} - V^{a}\nabla \times W_{a},$$
(7)

and

$$D^{a}[A,B]_{a} = B^{ab}\nabla \times A_{ab} - A^{ab}\nabla \times B_{ab}, \qquad (8)$$

The Maxwellean equivalent of GR is based on the correspondence between the Weyl tensor  $C_{abcd}$  and the electromagnetic tensor  $F_{ab}$  for any  $u_a$ . Thus

$$E_{ab} = C_{abcd} u^{c} u^{d} = E_{ab}, \ H_{ab} = *C_{abcd} u^{c} u^{d} = H_{ab}$$
(9)

These gravito-electric/magnetic spatial tensors are in principle physically measurable in the frames of comoving observers, and together they are equivalent to the space-time Weyl tensor.

The gravito-electromagnetic version of the electromagnetic tensor is

$$C_{ab}^{\ cd} = 4 \left\{ u_{[a} u^{[c]} + h_{[a}^{\ [c]} \right\} E_{b]}^{\ d]} + 2\varepsilon_{abe} u^{[c} H^{d]e} + 2u_{[a} H_{b]e} \varepsilon^{cde}.$$
(10)

In the 1 + 3 covariant approach to GR, the source of gravity is a fluid in which the energy density, pressure and the gravito-electromagnetic tensors are the fundamental quantities and not the metric. These quantities are governed by the Bianchi identities and the Ricci identities for  $u^a$ . Einstein's equations incorporated through the algebraic definition of the Ricci tensor  $R_{ab}$  in terms of the energymomentum tensor  $T_{ab}$ . The Bianchi identities are

$$\nabla^a C_{abcd} = \nabla_{[a} \left( -R_{b]c} + \frac{1}{6} Rg_{b]c} \right). \tag{11}$$

here  $R = R_a^a$  and  $R_{ab} = \kappa \left( T_{ab} - \frac{1}{2} T_c^c g_{ab} \right)$  where  $\kappa$  is the Einstein constant. The gravitational equivalents of Maxwell's equations are therefore

$$D^{a}E_{ab} = -3\omega^{a}H_{ab} + \frac{1}{3}D_{a}\rho + [\sigma, H]_{a}, \qquad (12)$$

$$D^{a}H_{ab} = 3\omega^{a}E_{ab} - \frac{1}{3}(\rho + p)\omega_{a}\rho - [\sigma, E]_{a}, \qquad (13)$$

$$\dot{E}_{\langle ab \rangle} - \nabla \times H_{ab} = \theta E_{ab} + \sigma_{c \langle a} E_{b \rangle}^{\ c} - \omega^{c} \varepsilon_{cd(a} E_{b)}^{\ d} + 2\dot{u}^{c} \omega^{c} \varepsilon_{cd(a} H_{b)}^{\ d} - \frac{1}{2} (\rho + p) \sigma_{ab},$$

$$(14)$$

$$\dot{H}_{\langle ab\rangle} - \nabla \times E_{ab} = -\theta H_{ab} + \sigma_{c\langle a} H_{b\rangle}^{\ c} - \omega^{c} \varepsilon_{cd(a} H_{b)}^{\ d} - 2\dot{u}^{c} \omega^{c} \varepsilon_{cd(a} E_{b)}^{\ d}.$$
(15)

These are the full nonlinear gravito-electromagnetic equations in covariant form. In both electromagnetism and gravito-electromagnetism vorticity produces source terms however, gravity has additional sources from a tensor coupling of the shear to the field. Furthermore, in electromagnetism, there are no magnetic charge sources, but the gravito-magnetic field  $H_{ab}$  has the source  $(\rho + p)\omega_a$ . Since  $\rho + p$  is the relativistic inertial mass-energy density,  $(\rho + p)\omega_a$  is the "angular momentum density", which we identify as a gravito-magnetic "charge" density.

## 3. Modification of General Relativity

The electromagnetic analogy suggests a further interesting interpretation of the

vorticity. In flat spacetime, relative to inertial observers, the electric and magnetic vectors may be written as

$$\boldsymbol{E} = \boldsymbol{\nabla} \boldsymbol{V} - \hat{\partial}_t \boldsymbol{\alpha}, \ \boldsymbol{H} = \boldsymbol{\nabla} \times \boldsymbol{\alpha}, \tag{16}$$

where, V is the electric scalar potential and  $\alpha$  is the gravitomagnetic vector potential. We express the gravitomagnetic vector potential as  $\alpha = v = \mu(t)H_0 \times r$ . Here, the magnitude of  $H_0$  is the Hubble constant, r the radius vector, v the tangential velocity and  $\mu(t)$  the time dependent relative gravito-magnetic permeability. The curl of the tangential velocity for a given r results in an angular velocity about the z direction of

$$\omega = \mu(t)H_0. \tag{17}$$

Einstein's field Equations (EFE) need to be modified such that they include the contribution by the magnetic charge. We now proceed with a similar approach as in [24]. As stated earlier, in the 1 + 3 covariant approaches to GR, the source of gravity is a fluid in which the energy density, pressure and the gravito-electromagnetic tensors are the fundamental quantities and not the metric. This allows us to intuitively make the assumption that the gravito-magnetic field is a compact and expanding region of spacetime that can be described as a Ricci soliton of the form  $R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} = kg_{\mu\nu}$  with the Ricci tensor of the compact manifold ex-

pressed as a compact energy-momentum tensor field  $R_{ab} = \kappa \left( T_{ab} - \frac{1}{2} T g_{ab} \right) = k g_{ab}$ .

Here,  $\kappa$  is the Einstein gravitational constant k is a constant proportional to the cosmological constant. These expressions can only be self-consistent if and only if the traces are T = 0 and R = 0. Here, the compact stress momentum tensor field  $\left(T_{ab} - \frac{1}{2}Tg_{ab}\right)$  is considered as a barotropic fluid consisting of

gravito-electromagneto-energy when modeled at a sufficiently large scale. For example, the gravito-electromagneto-energy of the solar system, star and galactic clusters can be considered as fluid at sufficiently large scales relative to the intrinsic scale of the gravitationally bound system. By modelling the system as barotropic, it satisfies the condition of being homogeneous and isotropic in the volume it occupies. Einstein's field Equations (EFE) are then modified such that they include the GEM field modelled as 4-D compact Einstein manifold and thus expressing the EFE as follows:

$$G_{ab} - kg_{ab} + \Lambda g_{ab} = \kappa T_{ab}.$$
(18)

here,  $k = \mu^2 \Lambda = 3 \left(\frac{\mu H_0}{c}\right)^2$  with a trace R = 0. Here,  $\mu H_0$  is the angular velocity of a Ricci soliton of reduced radius  $r = \frac{c}{2\pi\mu H_0}$ . This implies a rotational velocity on the surface of the soliton of  $v = H_0 r = \frac{c}{2\pi\mu}$ . This circular motion on the surface of the Ricci soliton arises from the traceless condition T = 0. This condition demands marginally stable or zero energy orbits on the surface in which the kinetic energy is equal to the gravitational potential energy.

Equation (18) must therefore satisfy a metric solution of the form

$$-c^{2}\tau^{2} = -c^{2}t^{2} + \mu(t)^{2} d\Sigma^{2}$$
(19)

where  $d\Sigma^2 = \frac{dr^2}{1-kr^2} + r^2 (d\theta^2 + \sin^2\theta d\phi^2)$  in reduced-circumference polar coordinates.

The analytic solutions to Equation (19) in which the energy momentum tensor is isotropic and homogeneous are the following equations:

$$\left(\frac{\dot{\mu}}{\mu}\right)^2 + \frac{\mu^2 \Lambda c^2}{3\mu^2} - \frac{\Lambda c^2}{3} = \frac{\kappa c^4}{3} \rho \rightarrow \left(\frac{\dot{\mu}}{\mu}\right)^2 + \frac{2\Lambda c^2}{3} = \frac{\kappa c^4}{3} \rho , \qquad (20)$$

$$2\frac{\ddot{\mu}}{\mu} + \left(\frac{\dot{\mu}}{\mu}\right)^2 + \frac{\mu^2 \Lambda c^2}{\mu^2} - \Lambda c^2 = -\kappa c^2 p \rightarrow 2\frac{\ddot{\mu}}{\mu} + \left(\frac{\dot{\mu}}{\mu}\right)^2 = -\kappa c^2 p, \qquad (21)$$

$$R = -6 \left[ \frac{\ddot{\mu}}{\mu} + \left( \frac{\dot{\mu}}{\mu} \right)^2 + \Lambda c^2 \right] = 0.$$
 (22)

From Equation (22) we obtain

$$\frac{\ddot{\mu}}{\mu} = -\left(\frac{\dot{\mu}}{\mu}\right)^2 - \Lambda c^2.$$
(23)

Substituting Equation (23) into Equation (21) yields

$$\left(\frac{\dot{\mu}}{\mu}\right)^2 = \kappa c^2 p - 2\Lambda c^2. \tag{24}$$

Substituting Equation (24) into Equation (20) and taking  $p = -\rho c^2$  yields

$$\Lambda = -\kappa c^2 \rho \,. \tag{25}$$

Given a measured value of the cosmological constant of  $1.1\times10^{-52}\,m^{-2}~$  we compute a critical baryonic matter density for the Ricci soliton of

$$\rho_{cb} = \rho_{DE} = 5.6 \times 10^{-27} \text{ kg} \cdot \text{m}^{-3}$$

The Ricci soliton therefore expands exponentially when  $p = -\rho c^2$  with a scale factor  $\mu(t) = \mu_0 e^{H_0 t}$ . Here  $H_0 = \sqrt{\kappa c^2 p - 2\Lambda c^2}$  This expansion occurs when the covariant flatness condition of Equation (19) is satisfied. Under such conditions the centripetal acceleration on the surface of the soliton is given by the expression

$$\frac{v^2}{r} = \frac{(H_0 r)^2}{r} = H_0^2 r = \frac{H_0^2 c}{2\pi\alpha H_0} = \frac{H_0 c}{2\pi\alpha}.$$
 (26)

In the comoving reference frame  $\alpha = 1$  which implies a scale invariant acceleration

$$a_0 = \frac{H_0 c}{2\pi} \,. \tag{27}$$

Under such conditions the radius is computed from Equation (26) as

$$r = \frac{2\pi v^2}{H_0 c} \tag{28}$$

Substituting *r* in the expression  $\frac{GM(r)}{r} = v^2$  we obtain

$$v^{4} = \frac{GM(r)H_{0}c}{2\pi}.$$
 (29)

This is the baryonic Tully Fisher relation and  $a_0 = \frac{H_0 c}{2\pi} \approx 1.1 \times 10^{-10} \text{ m/s}^2$  is the empirically observed Milgrom's acceleration constant. The theoretically computed value is in agreement with observations.

Since  $r = r_0 e^{H_0 t}$  then the velocity evolves with the scale factor as follows:

$$v^{4} = \frac{GM(r)H_{0}c}{2\pi}e^{4H_{0}t}$$
(30)

From the above equations, we also obtain the following equations of galactic and galactic cluster evolution

$$r = \frac{1}{H_0} e^{H_0 t} \left( GM_B(r) \frac{H_0}{2\pi} c \right)^{\frac{1}{4}} = \frac{v_n}{H_0}, \qquad (31)$$

$$v = \frac{\mathrm{d}r}{\mathrm{d}t} = \mathrm{e}^{H_0 t} \left( GM_B(r) \frac{H_0}{2\pi} c \right)^{\frac{1}{4}} = H_0 r , \qquad (32)$$

$$a = \frac{\mathrm{d}v}{\mathrm{d}t} = H_0 \mathrm{e}^{H_0 t} \left( GM_B \frac{H_0}{2\pi} c \right)^{\frac{1}{4}} = H_0 v \,. \tag{33}$$

### 4. Discussion and Concluding Remarks

The gravito-electromagnetism formulation of Roy Maartens and Bruce Bassett draws parallels between the equations governing electromagnetism and gravity. This analogy helps in developing a unified theoretical framework that simplifies the understanding of gravitational phenomena using well-established electromagnetic principles. The concept of super-energy and the super-Poynting vector offers deeper insights into the dynamics of spacetime. In this paper we have taken insights from the Roy Maartens and Bruce Bassett formulation to model dark matter as a compact gravito-magneto field with the same properties as a Ricci soliton which was first discussed in Ref. [26]. This model of dark matter offers a unique perspective compared to traditional dark matter models. It provides a classical, geometric approach to dark matter, contrasting with the particle-based explanations of traditional models. This could lead to new insights and potentially broaden our understanding of dark matter and dark energy.

#### Acknowledgements

The authors gratefully appreciate the discussions, suggestions and constructive criticism from Molethanyi Tshipa and Christian Corda.

## **Data Availability**

Empirical data used in this research can be found in the cited articles.

#### **Conflicts of Interest**

We declare no conflict of interest.

#### References

- Bernabei, R., Belli, P., Cappella, F., Cerulli, R., Dai, C.J., d'Angelo, A., *et al.* (2008) First Results from DAMA/LIBRA and the Combined Results with DAMA/NaI. *The European Physical Journal C*, **56**, 333-355. https://doi.org/10.1140/epjc/s10052-008-0662-y
- [2] Aprile, E. (2017) Search for Electronic Recoil Event Rate Modulation with 4 Years of XENON100 Data. *Physical Review Letters*, **118**, Article 101101.
- Gaitskell, R.J. (2004) Direct Detection of Dark Matter. Annual Review of Nuclear and Particle Science, 54, 315-359. https://doi.org/10.1146/annurev.nucl.54.070103.181244
- [4] Drukier, A.K., Freese, K. and Spergel, D.N. (1986) Detecting Cold Dark-Matter Candidates. *Physical Review D*, 33, 3495-3508. <u>https://doi.org/10.1103/physrevd.33.3495</u>
- Behnke, E., Behnke, J., Brice, S.J., Broemmelsiek, D., *et al.* (2012) First Dark Matter Search Results from a 4-kg CF3I Bubble Chamber Operated in a Deep Underground Site. *Physical Review D*, 86, Article 052001. https://doi.org/10.1103/physrevd.86.052001
- [6] Aalbers, J., Akerib, D.S., Akerlof, C.W., Al Musalhi, A.K., *et al.* (2023) First Dark Matter Search Results from the LUX-ZEPLIN (LZ) Experiment. *Physical Review Letters*, **131**, Article 041002. <u>https://doi.org/10.1103/PhysRevLett.131.041002</u>
- [7] Profumo, S., Giani, L. and Piattella, O.F. (2019) Introduction to Particle Dark Matter.
- [8] Hooper, D. and Profumo, S. (2007) Dark Matter and Collider Phenomenology of Universal Extra Dimensions. *Physics Reports*, 453, 29-115. <u>https://doi.org/10.1016/j.physrep.2007.09.003</u>
- [9] Fan, J., Katz, A., Randall, L. and Reece, M. (2013) Double-Disk Dark Matter. *Physics of the Dark Universe*, 2, 139-156. <u>https://doi.org/10.1016/j.dark.2013.07.001</u>
- [10] Dodelson, S. and Widrow, L.M. (1994) Sterile Neutrinos as Dark Matter. *Physical Review Letters*, 72, 17-20. <u>https://doi.org/10.1103/physrevlett.72.17</u>
- [11] Chung, D.J.H., Kolb, E.W., Riotto, A. and Senatore, L. (2005) Isocurvature Constraints on Gravitationally Produced Superheavy Dark Matter. *Physical Review D*, 72, Article 023511. <u>https://doi.org/10.1103/physrevd.72.023511</u>
- Starobinsky, A.A. (1980) A New Type of Isotropic Cosmological Models without Singularity. *Physics Letters B*, **91**, 99-102. https://doi.org/10.1016/0370-2693(80)90670-x
- [13] Barausse, E., Sotiriou, T.P. and Miller, J.C. (2008) A No-Go Theorem for Polytropic Spheres in Palatini f(R) Gravity. *Classical and Quantum Gravity*, 25, Article 062001. <u>https://doi.org/10.1088/0264-9381/25/6/062001</u>
- [14] Cembramos, J.A.R. (2011) Dark Matter from R<sup>2</sup>-Gravity. Journal of Physics: Conference Series, 315, Article 012004. <u>https://doi.org/10.1088/1742-6596/315/1/012004</u>
- [15] Harko, T., Lobo, F.S.N., Nojiri, S. and Odintsov, S.D. (2011) f(R,T) Gravity. *Physical Review D*, **84**, Article 02420. <u>https://doi.org/10.1103/physrevd.84.024020</u>

- [16] Alvarenga, F.G., de la Cruz-Dombriz, A., Houndjo, M.J.S., Rodrigues, M.E. and Sáez-Gómez, D. (2013) Dynamics of Scalar Perturbations in f(R,T) Gravity. *Physical Review D*, **87**, Article 103526. <u>https://doi.org/10.1103/physrevd.87.103526</u>
- [17] Shamir, M.F. and Sadiq, M.A. (2018) Modified Gauss-Bonnet Gravity with Radiating Fluids. *The European Physical Journal C*, 78, Article No. 279. <u>https://doi.org/10.1140/epjc/s10052-018-5763-7</u>
- [18] Cord, C. (2009) Interferometric Detection of Gravitational Waves: The Definitive Test for General Relativity. *International Journal of Modern Physics D*, 18, 2275-2282.
- [19] Green, M.A., Moffat, J.W. and Toth, V.T. (2018) Modified Gravity (MOG), the Speed of Gravitational Radiation and the Event GW170817/grb170817A. *Physics Letters B*, 780, 300-302. <u>https://doi.org/10.1016/j.physletb.2018.03.015</u>
- [20] Maartens, R. and Bassett, B.A. (1998) Gravito-Electromagnetism. Classical and Quantum Gravity, 15, 705-717. <u>https://doi.org/10.1088/0264-9381/15/3/018</u>
- [21] Maartens, R., Ellis, G.F.R. and Siklos, S.T.C. (1997) Local Freedom in the Gravitational Field. *Classical and Quantum Gravity*, 14, 1927-1936. <u>https://doi.org/10.1088/0264-9381/14/7/025</u>
- [22] Maartens, R., Lesame, W.M. and Ellis, G.F.R. (1997) Consistency of Dust Solutions with div H = 0. *Physical Review D*, 55, 5219-5221. <u>https://doi.org/10.1103/physrevd.55.5219</u>
- [23] Ellis, G.F.R. (1971) Topology and Cosmology. *General Relativity and Gravitation*, 2, 7-21. <u>https://doi.org/10.1007/bf02450512</u>
- [24] Hogan, P.A. and Ellis, G.F.R. (1997) Propagation of Information by Electromagnetic and Gravitational Waves in Cosmology. *Classical and Quantum Gravity*, 14, A171-A188. <u>https://doi.org/10.1088/0264-9381/14/1a/015</u>
- [25] Jean, M. (1973) Cargese Lectures in Physics. Gordon and Breach.
- [26] Marongwe, S. and Kauffman, S.A. (2024) Dark Matter as a Ricci Soliton.