

# Spatio-Temporal Study of Toxic and Harmful Planktonic Taxa at Two Oceanic Sites Mehdia and Moulay Bouselham, Rabat Sale Kenitra Region, Morocco, between 2017 and 2018

A. Hal Aberrhaman, B. Bouhaddioui, L. A. Lrhorfi, R. Bengueddour

Biology Department, Laboratory Biochemistry, Biotechnology, Health and Environment, Faculty of Science, Ibn Tofail University, Kenitra, Morocco

Email: azza-20111@hotmail.fr, dr\_bouchra2010@yahoo.fr, alrhorfi\_com@yahoo.fr, rachidbengueddour@yahoo.fr

**How to cite this paper:** Aberrhaman, A.H., Bouhaddioui, B., Lrhorfi, L.A. and Bengueddour, R. (2020) Spatio-Temporal Study of Toxic and Harmful Planktonic Taxa at Two Oceanic Sites Mehdia and Moulay Bouselham, Rabat Sale Kenitra Region, Morocco, between 2017 and 2018. *Open Journal of Marine Science*, 10, 41-51. <https://doi.org/10.4236/ojms.2020.102004>

**Received:** December 22, 2019

**Accepted:** March 29, 2020

**Published:** April 2, 2020

Copyright © 2020 by author(s) and Scientific Research Publishing Inc.

This work is licensed under the Creative Commons Attribution International License (CC BY 4.0).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

## Abstract

Phytoplankton is all cyanobacteria and microalgae (microscopic plants) present in surface water that may be carried by water currents. The study was conducted at two oceanic sites Mehdia and Moulay Bouselham, Rabat sale Kenitra region, Morocco, between 2017 and 2018. The objective of this study is to compare the spatiotemporal evolution of certain toxic planktonic species between the two sites. The planktonic species identified all have toxic potency but to varying degrees such as Alexandrium, Dinophysis, Gymnodinium, Pseudonitzschia and Proocentrum. The comparison of the abundance of these taxa in the two sites confirms that the Moulay Bouselham site is characterized by the presence of the most toxic planktonic species represented mainly by Pseudonitzschia in contrast the Mehdia site experienced an abundance of the Proocentrum kind. This situation requires a more detailed study of the essential causes of the proliferation of these species.

## Keywords

Phytoplankton, Surface Water, Toxic Power, Spatiotemporal Evolution, Mehdia, Moulay Bouselham

## 1. Introduction

Phytoplankton is all the cyanobacteria and microalgae (microscopic plants) present in surface water that can be carried away by water currents [1] [2]. They are defined as unicellular or multicellular organisms, either eukaryotes or prokaryotes [3]. Microalgae occupy most ecological niches, mainly aquatic envi-

ronments; they live in salt marshes, in acidic or strongly alkaline environments [4]. According to Greenway and Munns (1980); Dabbadie (1992); Netondo *et al.* (2004); Perrine *et al.* (2012) [5] [6] [7] [8], the factors favoring a better microalgal production are the illumination, the temperature whose optimum is 18°C to 24°C affording to the species. The Ph must be between 8.2 and 8.7, nitrogen and phosphorus and other elements. Phytoplankton occupies the place of the first link in the food chain [9], it is the basis of ecosystems in marine environments [10]. The distribution, abundance, production and biodiversity of different species or groups of plankton are likely to be deeply affected by climatic changes and the physical and chemical properties of oceans [1]. Of the many species that make up phytoplankton, some produce toxins called phycotoxins. They belong mainly to the Dinophyceae class [11].

In Morocco, there are 84 listed wetlands according to the 1997 Protected Areas Study. Another study reports about 300 sites covering an area of 400,000 ha. Morocco's Gharb is rich in wetlands such as the Mehdia area and Moulay Bouselham, which are considered among the most populated areas by planktonic species.

The objective of this study is to compare the spatiotemporal evolution of certain toxic planktonic species between two wet sites Mehdia and Moulay Bouselham.

## 2. Material and Methods

### 2.1. Description of the Study Site

The samples were taken from two ocean sites, the Mehdia and Moulay Bouselham in the Rabat-Sale-Kenitra region. The first site is a picturesque coastal town near the Kenitra city, 30 km northeast of the capital Rabat. The average annual temperature is 18.6°C and the average rainfall is 283 mm. The second site is known for heavy rainfall in winter than in summer. The average annual temperature is 18.1°C, and the average annual precipitation is 622 mm.

### 2.2. Identification of Species

For the identification of phytoplankton, a volume of 500 ml of seawater is collected at 0.5 m depth using opaque glass bottles. The sample is then fixed with 2.5 ml of alkaline lugol and placed in a cool box protected from light. In the laboratory, samples are recorded and stored in the dark at 4°C.

After vigorous stirring, the samples are decanted for 24 hours and then subjected to observation under optical microscope morpho-anatomic characters (shape, size and color) representing the identification keys retained by Gregoire *et al.*, 1978; Bourrelly, 1985 [12] [13]. The cells are then counted per liter per vat.

### 2.3. Statistical Analyzes

The collected data are entered on an Excel support, the exploitation is carried out on a statistical processing software. The selected analyzes in this sense are

significance tests such as single-point analysis of variance, correlations, principal component analysis. The results are generally expressed as relative frequencies and/or means  $\pm$  standard deviation.

### 3. Results and Discussion

#### 3.1. Study of the Density Variation in the Two Sampling Sites

**Table 1** presents the results of the distribution of planktonic cells according to the years of sampling and according to the sampling sites. The results are expressed in mean and standard deviation. The comparison test and the student test. An Asterix marks significant differences.

\* In a first step, we compared the average densities between the two sites by year of sampling. The results obtained show that, in general, the abundance in 2017, except for the *Proocentrum* taxon where the density in the Mehdiya station exceeds that of Moulay Bouselham, the other planktonic species have been highly represented in the Moulay Bouselham station. The student test confirmed the effect of the site only for the *Dinophysis* taxon with a  $t = 3.41$  and  $p < 0.003$ .

\*In a second time, and in 2018, the average density of plankton in the Moulay Bouselham site exceeds that of the Mehdiya site, with the exception of *Gymnodinium* and *Pseudonitzschia* where it is the opposite. The student test shows a significant difference between the two stations for the *Alexandrium* taxon.

The planktonic species identified all have toxic potency but to varying degrees. Because of this, the abundance of these species in the Moulay Bouselham site makes the environment really too incriminated.

#### 3.2. Study of the Spatio-Temporal Variation of Toxic Taxa

##### 3.2.1. Variation of the Genus *Alexandrium*


*Alexandrium* is a nucleated unicellular living organism belonging to the Dinophyte group. Several *Alexandrium* species (*Gonyaulax* or *Protogonyaulax*) produce paralytic toxins, particularly in the Pacific Northwest, North East Coast of North America and Northern Europe. While in Morocco, paralytic poisoning is due to the efflorescence of *Gymnodinium catenatum* [14].

The distribution of *Alexandrium* densities according to years of sampling shows a very remarkable difference between 2017 and 2018, these averages are respectively  $217.83 \pm 470.19$  cells/liter (Min = 0, Max = 1990) and  $89.55 \pm 74.03$  cells/liter (Min = 0, Max = 200). This difference in density is largely felt between the two sampling sites Mehdiya and Moulay Bouselham, with mean values of  $61.39 \pm 7.98$  cells/liter (Min = 0, Max = 290) and  $251.57 \pm 46.17$  respectively. Cells/liter (Min = 0, Max = 1990), this difference is significant between the means of the two sampling sites (Fisher = 3.68,  $p < 0.048$ ). Analysis of variance did not show any interaction between the year factor and the sampling site factor (Fisher = 1.06,  $p < 0.31$ ). **Figure 1** shows the mean points of the monthly density in the two stations. The results show an abundance of the *Alexandrium* species in Moulay Bouselham station. Indeed, two maximum peaks characterize the two sites,

**Table 1.** Presentation of plankton species according to years and sampling sites.

species	Site	Year of sampling: 2017		Year of sampling: 2018		Significant
		Mean (cells/liter)	Standard deviation	Mean (cells/liter)	Standard deviation	
Alexandrium	Site I	74.33	95.51	47.27	50.81	NS
	Site II	361.33	638.92	131.82	70.68	NS
signification		t = 1.54; p < 0.14		t = 3.54; p < 0.004**		
Dinophysis	Site I	41.67	90.03	122.73	375.08	NS
	Site II	371.67	322.77	1036.36	2198.76	NS
signification		t = 3.41; p < 0.003**		t = 1.36; p < 0.194		
Gymnodinum	Site I	216.67	257.91	877.27	1680.98	NS
	Site II	466.67	502.87	172.73	134.84	t = 1.87; p < 0.037**
signification		t = 1.3; p < 0.14		t = 1.39; p < 0.19		
Pseudonitzschia	Site I	2741.67	4792.13	4727.27	5921.84	NS
	Site II	10,616.67	18,005.04	3259.09	2972.78	NS
signification		t = 1.46; p < 0.16		t = 0.73; p < 0.47		
Proocentrum	Site I	6166.67	9369.52	3000.00	1788.85	NS
	Site II	4250.00	4025.49	3727.27	1489.36	NS
signification		t = 0.65; p < 0.5		t = 1.04; p < 0.31		

NS: no significant; \*\*: significant at 5%.



**Figure 1.** Number of cells counted per month at both sites at Alexandria.

- For the Moulay Bouselham station, average densities are highest during the months of April and August, which correspond to the spring and summer seasons, with respectively more than 1000 cells/liter and more than 800 cells/liter.
- For the Mehdia site, the monthly distribution shows that the cell density peaks are recorded during the month of May (more than 180 cells/liter) and the month of October (more than 200 cells/liter) which correspond to the

seasons spring and autumn.

### 3.2.2. Variation of the Genus *Dinophysis*

The *Dinophysis* genus belongs to the class of dinophyceae. It includes many *Dinophysis* species, most of which are toxic.

The annual distribution of the average density of *Dinophysis* shows that in 2017, the average is  $206.67 \pm 58.49$  cells/liter (Min = 0, Max = 900) and in 2018 it reaches  $579.55 \pm 342.97$  cells/liter (Min = 0, Max = 7500). Analysis of variance showed a significant sampling site effect on the distribution of this plankton species (Fisher = 3.50,  $P < 0.044$ ). Indeed, the average density in the Moulay Bouselham station ( $689.57 \pm 320.66$  cells/liter) exceeds that of Mehdia ( $80.43 \pm 55.05$  cells/liter). The analysis of variance did not show any interaction between the year factor and the sampling site factor (Fisher = 0.80,  $p < 0.37$ ). **Figure 2** shows the mean points of the monthly density of the planktonic *Dinophysis* species at both stations. It appears that this species is very abundant in the Moulay Bouselham station. Indeed, two maximum peaks characterize the two sites:


- The monthly variation of the *Dinophysis* density in the Moulay Bouselham station has a peak exceeding 4000 cells/liter during the month of June, and to a lesser degree a second peak during the month of March (more than 170 cells/liter).
- For the Mehdia site, the monthly distribution shows that cell density peaks are recorded during the month of July (more than 500 cells/liter) and low during the other months of the year.

### 3.2.3. Variation of the Species *Gymnodinium*


The *Gymnodinium* sp., represent an important genus of the dinoflagellate protist group, the marine plankton without theca of the class Dinophyceae, order Gymnodinales. It is a species of red tide forming chains and producing toxins.

The average density of the cells of this planktonic species per liter per year of sampling shows that this average is  $341.67 \pm 83.93$  cells/liter during the year 2017, with a minimum of 0 cells/liter and a maximum of 1600 cell/liter, whereas for the year 2018, the average is  $525.00 \pm 259.74$  cells/liter (Min = 0, Max = 4500). Despite the difference in record between these two years, the Fisher test did not assert the direct effect of the year variation on density (Fisher = 0.48,  $p < 0.49$ ). On the other hand, the distribution of the number of planktonic cells per liter of water per sampling site shows that the Mehdia station has an average density of  $532.61 \pm 249.48$  cells/liter (min = 0, max = 4500), higher than that recorded at Moulay Bouselham (mean =  $326.09 \pm 82.68$  cells, Min = 0, Max = 1600). Given the large intra-station dispersion, the Fisher test did not show a significant difference between the two stations (Fisher = 0.62,  $p < 0.44$ ). The analysis of variance shows that the year factor and the station factor interact significantly with each other (Fisher = 3.44,  $p < 0.047$ ), thus having an effect on the density of this plankton.

**Figure 3** shows the mean points of the monthly density of the planktonic



**Figure 2.** Number of cells counted per month in the two sites at Dinophysis.



**Figure 3.** Number of cells counted per month at both sites at Gymnodinium.

Gymnodinium species at both stations. As a result, this taxon is very common in the Mehdia station, with two peaks in April and May; the density during these two months exceeds 2000 cells/liter. However, in the Moluay Bouselham station, there were three peaks in February (more than 500 cells/liter), in June (more than 600 cells/liter) and in October (more than 900 cells /liter).

### 3.2.4. Variation of the Species Pseudonitzschia

The micro-algae of the genus Pseudonitzschia are Diatoms of the family Bacillariaceae. Some species of this genus produce domoic acid, a neurotoxin responsible for food poisoning (ASP for Amnesic Shellfish Poisoning).

The mean densities of the Pseudonitzschia cells determined for the two years of study 2017 and 2018 are respectively  $6679.17 \pm 2755.33$  cells/liter (Min = 0, Max = 65,000) and  $3993.18 \pm 987.93$  cells/liter (Min = 0, Max = 19,000). The Fisher test did not show any effect of the year variation on the distribution of the cell density despite the difference between the average densities during the years

2017 and 2018, this is attributed to the great heterogeneity recorded in each group. The comparison of the mean *Pseudonitzschia* densities between the two sites, Mehdia and Moulay Bouselham, shows a great difference between these two stations, the corresponding averages recorded being  $3691.30 \pm 1112.202$  (Min = 0, Max = 19,000) in the Mehdia station and  $7097.83 \pm 2799.298$  (Min = 0; Max = 65,000) in the Moulay Bouselham station.


**Figure 4** shows the mean points of the monthly density of the planktonic species *Pseudonitzschia* in both stations. It appears from this figure that this taxon is very common in Moulay Bouselham station, with a peak in July and the density reaches more than 32,000 cells/liter. However, the distribution of this monthly taxon in the Mehdia station fluctuates around a density of 5000 cells/liter, especially in February, April, August and October.

### 3.2.5. Variation of the Proocentrum Species


Proocentrum is a dinoflagellate in formation, forming an armor, marine, planktonic. It is a cosmopolitan species in cold temperate waters with tropical waters. It is generally considered harmless [15] [16] [17].

The evolution of this taxon between 2017 and 2018 shows a decrease in average density, the rate of decline reaches 35.48%. The minimum number of cells determined is zero cells/liter and the maximum is displayed during the year 2017, it is 28,000 cells/liter. Regarding the distribution of this taxon in the two sampling sites, we did not notice a large difference between these two sites. The file test did not show a significant difference (Fisher = 0.17,  $p < 0.61$ ), between the mean density at the Mehdia site is 4652.17 cells/liter and the average density at the Moulay Bouselham site is 4000.00 cells/liter.

The monthly distribution at both stations is shown in **Figure 5**. The distribution of the Proocentrum at the Mehdia site is marked by a maximum in October (more than 14,000 cells/liter) and second peak in August but that does not exceed 8000 cells/liter. However, the distribution in the Moulay Bouselham station, a maximum is recorded in October at a density of more than 10,000 cells/liter, for other months, the density fluctuates between 2000 and 4000 cells/liter.



**Figure 4.** Number of cells counted per month in both sites at *Pseudonitzschia*.



**Figure 5.** Number of cells counted per month in both sites at proocentrum.


### 3.3. Global Analysis

In this part, we used a joint analysis of the five toxic taxa according to the cell density, the site of the site and the month of sampling. The two figures below represent hierarchical classification dendrograms of the five species identified. **Figure 6(a)** shows the projection results of the toxic taxa identified in the Mehdia site. This analysis reveals three distinct classes. The first group is composed of the Alexandria, Dinophysis and Gymnodinium taxa. The second group includes the Pseudonitzschia taxon and the third group is Proocentrum. In addition, **Figure 6(b)**, presents the results of the classification in the Moulay Bousselham site, this time, three groups are thus reformed, the first group is composed of the taxa Alexandrium, Gymnodinium and Dinophysis. A second group consisting of Proocentrum and a third group includes the Pseudonitzschia taxon.

The comparison of the abundance of these taxa in both sites confirms that the Moulay Bousselham site is characterized by the presence of genus with more toxic planktonic species represented mainly by Pseudonitzschia, in contrast to the Mehdia site known an abundance of the genus Proocentrum. This situation requires a more detailed study of the essential causes of the proliferation of these species.

## 4. Conclusion

There are five types of phytotoxins involved in human poisoning worldwide: • PSP or paralytic toxins • DSP or diarrheal toxins • NSP or neurotoxins • ASP or amnesic toxins • CFP or ciguatera toxins PSP toxins are the most common. The identification of the toxic species in the two Moroccan sites allowed us to mention five species known by their toxic power. Moreover, Dinophysis and Alexandrium are two planktonic taxa of which some species are capable of producing respectively diarrheal toxins and paralytic toxins. These two types of toxins accumulate in shells that feed on phytoplankton: this is the case of filtering shells



**Figure 6.** (a) Hierarchical presentation of planktonic taxa in the Mehdiya site. ALEX: Alexandrium; DINOPH: Dinophysis; gymm: Gymnodinium; pseudo: Pseudonitzschia; procc: Proocentrum; (b) Hierarchical presentation of planktonic taxa in the Moulay bousel-ham site. ALEX: Alexandrium; DINOPH: Dinophysis; gymm: Gymnodinium; pseudo: Pseudonitzschia; procc: Proocentrum.

that become toxic to consumers, without being themselves affected. According to Lassus and Bardouil (1991); Marcaillou *et al.* (1990) [18] [19], the genus Dinophysis is widely represented in almost all the seas of the globe and in particular in coastal areas including the coasts of the Atlantic. Dinophysis produces diarrheal toxins, called DSP (Diarrhetic Shellfish Poison) [20]. Another genus known by its toxic power that exists on the Moroccan coasts is Gymnodinium, which produces cytotoxic substances, hemolytic and aggressive for cell membranes [21]. However, the genus Proocentrum is not toxic to marine life, but some species of this genus may be toxic to the consumer. In fact, they produce ASP21 toxins or amnesic toxins, which can cause neurological disorders, including memory loss, in consumers of contaminated shellfish. The same assemblage of toxic planktonic species was found in a study conducted in Tunisia by Nejib *et al.* (2005) [22].

## Conflicts of Interest

The authors declare no conflicts of interest regarding the publication of this paper.

## References

- [1] Bougis, P. (1974) Ecology of Marine Plankton. II Zooplankton. Masson et Cie.
- [2] Dufour, P. and Durand, J.R. (1982) Plant Production in the Ivory Coast lagoons. *Revista de Biología Tropical*, **15**, 209-230.
- [3] Sialve, B. and Steyer, J.P. (2013) Microalgae, Promises and Challenges. *Innovations Agronomiques*, **26**, 25-39.
- [4] Cadoret, J.P., Bardor, M., Lerouge, P., Cabiglieria, M., Henriquez, V. and Carlier, A. (2008) Microalgae Cellular Plants Producing Recombinant Commercial Molecules. *Médecine/Sciences*, **24**, 375-382. <https://doi.org/10.1051/medsci/2008244375>
- [5] Greenway, H. and Munns, R. (1980) Mechanisms of Salt Tolerance in No Halophytes. *Annual Review of Plant Physiology*, **31**, 149-190. <https://doi.org/10.1146/annurev.pp.31.060180.001053>
- [6] Dabbadie, L. (1992) Intensive Cultures of Microalgae on PIG Manure: Performance, Constraints, Use of Biomass. Diploma in Advanced Agronomy. National Agronomic School of Montpellier, France, 123 p.
- [7] Netondo, G.W., Onyango, J.C. and Beck, E. (2004) Sorghum and Salinity: II. Gas Exchange and Chlorophyll Fluorescence of Sorghum under Salt Stress. *Crop Science*, **44**, 806-811. <https://doi.org/10.2135/cropsci2004.8060>
- [8] Perrine, Z., Negi, S. and Sayre, R.T. (2012) Optimization of Photosynthetic Light Energy Utilization by Microalgae. *Algal Research*, **1**, 134-142. <https://doi.org/10.1016/j.algal.2012.07.002>
- [9] Dhargalkar, V.K. and Ingoleb, S. (2004) Phytoplankton Identification Manual. Ed National Institute of Oceanography, 2 p.
- [10] Broutin, M., Caffier, G., Madi, M. and Artigas, L. (2011) Techniques for Monitoring the Abundance, Biomass and Diversity of Phytoplankton in Marine Waters. Ifremer, 2.
- [11] Sournia, A., Belin, C., Billard, C., Catherine, M., Erard-Le Denn, E., Fresnel, J., Lassus, P., Pastoureaud, A. and Soulard, R. (1992) The Repetitive and Expanding Occurrence of a Green Bloom-Forming Dinoflagellate (Dinophyceae) on the Coasts of France. *Cryptogamie Algologie*, **13**, 1-13.
- [12] le Grand, G. (1978) Dialogues. Volume I. Introduction, Bibliography and Maps by Adalbert de Vogué.
- [13] Bourrelly, P. (1985) Freshwater Algae. Introduction to Systematics. Volume III: Blue and Red Algae. The Euglenians, Peridinians and Cryptomonadines. Paris, 606 p.
- [14] Hall, S., Strichartz, G., Moczydlowski, E., Ravindran, A. and Reichardt, P.B. (1990) The Saxitoxins: Sources, Chemistry, and Pharmacology. In: *Marine Toxins: Origin, Structure, and Molecular Pharmacology*, American Chemical Society, New York, 29-65. <https://doi.org/10.1021/bk-1990-0418.ch003>
- [15] Taylor, D.L. and Seliger, H.H. (1979) Toxic Dinoflagellate Blooms. Elsevier, North Holland, New York, 505.
- [16] Anderson, D.M., White, A.W. and Baden, D.G. (1985) Toxic Dinoflagellates. Elsevier, New York, 561.

- [17] Graneli, E., Sundstrom, B., Edler, L. and Anderson, D.M. (1990) Toxic Marine Phytoplankton. Elsevier, New York, 554.
- [18] Lassus, P. and Bardouil, M. (1991) Le Complexe "Dinophysis Acuminata": Identification of Species along the French Coast. *Cryptogamie Algologie*, **12**, 1-9.
- [19] Marcaillou-Le Baut, C., Bardin, B., Bardouil, M., Bohec, M., Le Déan, L., Masselin, P. and Truquet, P. (1990) Study of the Decontamination of Toxic Mussels (Diarrheal Toxins) in the Laboratory and in the Wild. Internal Report IFREMER/ DERO-90.02-MR, 21 p.
- [20] UNESCO (1996) Design and Implementation of Some Harmful Algal Monitoring Systems. IOC Technical Series No. 44.
- [21] Sournia, A., Belin, C., Berland, B., Erard-le den, E., Gentien, P., Grzebyk, D., Marcaillou-le Baut, C., Lassus, P. and Partensky, F. (1991) The Harmful Phytoplankton of the Coasts of France. From Biology to Prevention. IFREMER, S.D.P., 154 p.
- [22] Nejib, M., Daly, Y., Souissi, S., Maamouri, F. and AISSAP (2005) Association Tintinnides (Ciliophora, Tintinnina) Dinoflagellés (Dinophyceae) Autotrophic Potentially Harmful at the Level of the Bay of Tunis, and Two Associated Lagoons: Ghar Elmelh and Tunis Sud. *La mer*, **43**, 19-32.